

U005-12

Room:IC

Time:May 26 11:55-12:10

Preliminary examination of Hayabusa asteroidal samples: Organic compound analyses

Hiroshi Naraoka^{1*}, Hajime Mita², Kenji Hamase³, Masashi Mita⁴, Hikaru Yabuta⁵, Kaori Saito⁶, Kazuhiko Fukushima⁶, Fumio Kitajima¹, Scott A. Sandford⁷, Tomoki Nakamura⁸, Takaaki Noguchi⁹, Ryuji Okazaki¹, Keisuke Nagao¹⁰, Mitsuru Ebihara¹¹, Hisayoshi Yurimoto¹², Akira Tsuchiyama⁵, Masanao Abe¹³, Toru Yada¹³, Yukihiro Ishibashi¹³, Kei Shirai¹³, Munetaka Ueno¹³, Tatsuaki Okada¹³, Akio Fujimura¹³, Toshifumi Mukai¹³, Makoto Yoshikawa¹³, Junichiro Kawaguchi¹³

¹Dept. Earth & Planet. Sci., Kyushu Univ., ²Fukuoka Inst. Technol., ³Grad. School Pharm. Sci., Kyushu Univ., ⁴Frontier Sci. Dev. Shiseido Co. Ltd., ⁵Grad. School Sci., Osaka Univ., ⁶Grad. School Agricul. Sci., Nagoya Univ., ⁷NASA-Ames Research Center, ⁸Fac. Sci., Tohoku Univ., ⁹Fac. Sci., Ibaraki Univ., ¹⁰Grad. School Sci., Univ. Tokyo, ¹¹Fac. Sci. & Tech., Tokyo Metro. Univ., ¹²Dept. Nat. History Sci., Hokkaido Univ., ¹³ISAS/JAXA

Many organic molecules have been observed in the interstellar medium as well as extraterrestrial materials. In particular, various organic compounds including amino acids are reported from carbonaceous chondrites, which may have connections to emergence of life on the primitive Earth. By the Hayabusa mission, the existence of organic matter at the surface of Asteroid 25143 Itokawa can be examined without terrestrial contaminants. Ordinary chondrites, which seem to have an origin of S-type asteroid including Itokawa [1] [2], are generally depleted in not only volatile organic compounds but also carbonaceous materials due to their relatively high formation temperature. However, the existence of organic compounds is possible at the surface of Itokawa, because indigenous amino acids are found in lunar soils by Harada et al. [3] and Brinton and Bada [4] and interplanetary dust particles (IDPs) by Brinton et al. [5]. In addition, polycyclic aromatic hydrocarbons (PAHs) are reported in a Martian igneous meteorite [6] and IDPs [7]. In this study, we will perform a preliminary organic compound analysis on particles from asteroid Itokawa returned by the Hayabusa mission.

The particles are rinsed with small amount of methanol/dichloromethane on the plate. The extract is hydrolyzed with HCl followed by separation into amino acid and other organic compound fractions. Amino acid analysis with enantiomeric separation is carried out using two-dimensional high-performance liquid chromatography with highly sensitive fluorescence detectors (2D-HPLC/FD) [8]. The other organic fraction including PAHs is subjected to time of flight-secondary ion mass spectrometry (ToF-SIMS) analysis. The ToF-SIMS analysis is also applied directly to the carbonaceous materials in the particles.

Detection of amino acids and PAHs is highly dependent on the concentrations of these compounds in particles as well as the sample amount available for the analyses. If the particles yield glycine, one of the abundant amino acids in extraterrestrial materials, as much as the concentration in carbonaceous chondrites (~10-500 fmol/micro gram) [9] [10], the 2D-HPLC/FD can reveal the amino acids distribution. If the particle contains glycine as the similar amount as lunar soils (~0.1 fmol/micro gram) [3] [4], the quantification competes against the detection limit. The ToF-SIMS analysis can identify various organic compounds including PAHs in a ppm level [12]. The compound distributions may clarify origins of organic compounds at the surface of Itokawa. If glycine is the most abundant as observed in lunar soils [3] [4] and cometary dusts [11], hydrogen cyanide (HCN) may contribute to the amino acid precursors. In the case of anhydrous minerals, the HCN may be implanted by solar wind. If alpha-aminoisobutyric acid (AIB) is abundant as observed in some CM chondrites, meteoritic source derived from aqueous altered parent body is considered after a Strecker-type reaction with ketones. In such a case, the organic compounds survived upon impact. The particle amount available for the preliminary organic analysis is very limited. Further investigations are necessary on the second stage of analysis with much sample by direct extraction with hot water.

References: [1] Abe M. et al. (2006) *Science*, 312, 1334. [2] Okada T. et al. (2006) *Science*, 312, 1338. [3] Harada K. et al. (1971) *Science*, 173, 433. [4] Brinton K. L. F. and Bada J. L. (1996) *GCA*, 60, 349. [5] Brinton K. L. F. et al. (1998) *OLEB*, 28, 413. [6] McKay D. S. et al. (1996) *Science*, 273, 924. [7] Clemett S. J. et al. (1993) *Science*, 262, 721. [8] Hamase K. et al. (2010) *J. Chromatogr. A*, 1217, 1056. [9] Ehrenfreund P. et al. (2001) *PNAS*, 98, 2138. [10] Martins Z. et al. (2007) *Meteoritics & Planet. Sci.*, 42, 2125. [11] Sandford S. A. et al. (2006) *Science*, 314, 1720.

Keywords: Hayabusa, Itokawa, Organic compounds