

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



U020-21

Room:304

Time:May 23 17:10-17:25

## Impact chemical evolution processes for biomolecules formed by oceanic impacts

Nao Fukunaga<sup>1</sup>, Toshimori Sekine<sup>1\*</sup>, Shunsuke Izumi<sup>2</sup>, Yoshihiro Furukawa<sup>3</sup>, Takeshi Kakegawa<sup>3</sup>, Takamichi Kobayashi<sup>4</sup>, Hiromoto Nakazawa<sup>4</sup>

<sup>1</sup>DEPSS, Hiroshima University, <sup>2</sup>Math. & Life Sci., Hiroshima Univ., <sup>3</sup>DEPMS, Tohoku University, <sup>4</sup>NIMS

The biomolecules on Earth are thought either to have come from the extraterrestrial parts carried with flying meteorites or to have been formed on Earth from the inorganic carbon through given energy. From the standpoint to address impact energy, we need to know possible process how the simple biomolecules formed at a violent impact have been evolved subsequently through several impacts at the time of the late heavy bombardment. In this study we investigated the simplest amino acid, glycine marked by <sup>13</sup>C, in order to understand how it will be evolved chemically when it is subjected to further impacts. The results indicate that some new molecules are formed and others are decomposed, and suggest not only that the impact-induced process is not so simple to proceed just to one way, but also that there are complicated and multi-process ways. It also must be taken into account the heterogeneous distribution of impact energy in an impact that may cause a significant effect on the chemical evolution.

Keywords: Biomolecules, Impact energy, Chemical evolution, Amino acids, Late heavy bombardment