

AEM002-01

会場:201A

時間:5月27日 08:30-08:45

Observation, Modelling of Lightning Activity in Hurricanes Observation, Modelling of Lightning Activity in Hurricanes

Colin Price¹, Yoav Yair^{2*}, Barry Lynn³, Na'ama Reicher¹
Colin Price¹, Yoav Yair^{2*}, Barry Lynn³, Na'ama Reicher¹

¹Tel-Aviv University, Tel-Aviv, Israel, ²The Open University, Ra'anana, Israel, ³Weather It Is, Efrat, Israel

¹Tel-Aviv University, Tel-Aviv, Israel, ²The Open University, Ra'anana, Israel, ³Weather It Is, Efrat, Israel

Modern global lightning detection networks enable us to monitor and investigate the lightning activity in areas where there is little or no coverage by local lightning location systems, such as the remote regions or the oceans. Hence, we are now better able to track and understand the complex connection between hurricane development and electrical activity in these dangerous storms. In a recent study (Price et al., 2009) it was shown that in category 4-5 hurricanes (typhoons), there is a clear increase of lightning activity approximately one day before the maximum intensity (lowest pressure, strongest winds) of the storm. Additional analysis confirms this initial finding, possibly allowing us to use real-time lightning observations to forecast the time of hurricane intensification. We have also started modeling lightning activity in hurricanes using the WRF mesoscale meteorological model. Numerical simulations of the electrical activity in hurricanes, employing microphysical parameterization of the charging potential (using the Lightning Potential Index, LPI; Yair et al., 2010) show a reasonable agreement with the observations of lightning for the few case studies we have analyzed thus far. Both observations and modeling results will be presented.

Price, C., M. Asfur and Y. Yair, 2009: Maximum hurricane intensity preceded by increase in lightning frequency, *Nature Geoscience*, doi:10.1038/NGEO477, 2, 329-332.

Y. Yair, B. Lynn, C. Price, V. Kotroni, K. Lagouvardos, E. Morin, A. Mugnai, and M. d. C. Llasat (2010). Predicting the potential for lightning activity in Mediterranean storms based on the Weather Research and Forecasting (WRF) model dynamic and microphysical fields, *J. Geophys. Res.*, 115, D04205, doi:10.1029/2008JD010868.

キーワード: Lightning, Hurricanes, Storm Intensity, Detection Networks, Modelling, WRF

Keywords: Lightning, Hurricanes, Storm Intensity, Detection Networks, Modelling, WRF

AEM002-02

会場:201A

時間:5月27日 08:45-09:00

ELF/VLF 帯空電観測に基づいた全球・アジア域における落雷活動の導出 Estimation of the global and Asian lightning activity based on the observation of ELF/VLF sferics

山下 幸三^{1*}, 高橋 幸弘², 佐藤 光輝², 土屋 史紀¹, 大矢 浩代³

Kozo Yamashita^{1*}, Yukihiro Takahashi², Mitsuteru Sato², Fuminori Tsuchiya¹, Hiroyo Ohya³

¹ 東北大・理・地球物理, ² 北大・理・宇宙理学, ³ 千葉大・工

¹Dept. of Geophysics, Tohoku Univ., ²Dept. of CosmoSciences, Tohoku Univ., ³Engineering, Chiba Univ

In the recent decade, many researchers have been attracted to the investigation of global lightning activity. One of the reasons is the progresses in the observation of global lightning discharges. Measurements of electromagnetic waves radiated from CGs with ground-based systems have been developed drastically.

Electromagnetic wave radiated from cloud-to-ground (CG) lightning discharge is observed as transient waveform called as atmospherics or sferics. Sferics in the Very Low Frequency (VLF: 3-30 kHz) and Extremely Low Frequency (ELF: 3-3000 Hz) bands can be detected at a receiver which is several mega-meters from CGs due to the propagation with quite low attenuation. This long propagation enables us to monitor the global CGs activity only with single or few receivers. Using observed waveform in ELF or VLF range, information not only about the location but also about the electrical properties such as polarity, peak current and charge moment (Qdl) of individual lightning stroke can be derived.

In this study, we developed new algorithm to estimate location and charge moments for relatively small Qdl events ($|Qdl| > 950$ C-km). This method is applied to the magnetic fields data observed in 1-100 Hz range obtained by global ELF observation network (GEON), Syowa station in Antarctica, Onagawa station in Japan, Esrange in Sweden, and Santa Cruz in U.S.A. Making use of the time-of-arrival method, the accuracy of geolocation and detection sensitivity is greatly modified. Thanks to this new algorithm, about a millions of CGs can be analyzed with one month (during January 2004). Global CGs distribution is derived with annual data (from October 2003 to July 2004) and about a million CGs whose Qdl are larger than 950 C-km are obtained. The results show the monthly and seasonal variation of global CGs distribution.

Furthermore, a new VLF observation network (Asian VLF observation network: AVON) is developed to monitor the activity of CGs in Southeast Asia. Observation sites of this system are located at Tainan in Taiwan, Saraburi in Thailand, and Pontianak in Indonesia. In this study, data observed at Tainan station and Pontianak station during three days (from October 13th, 2010 to October 15th, 2010) is used for the initial analysis. Using the waveforms obtained by this network, 1-3 sferics per seconds are geolocated. Median of geolocation error is estimated as 93 km comparing with WWLLN data using about 400 CGs located in the Maritime Continents. Furthermore, charge moment distribution for the CGs of > 200 C-km in the Maritime Continents is derived with a combination of CG locations obtained by AVON and the transient waveform observed by GEON.

In this presentation, we introduce the speculation of new observation network to monitor the lightning activity in Southeast Asia and demonstrate the efficiency of combination of the measurement of ELF sferics and that of VLF ones.

キーワード: 雷, 空電, ELF, VLF

Keywords: lightning, sferics, ELF, VLF

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-03

会場:201A

時間:5月27日 09:00-09:15

The characteristics of global lightning activities observed by ISUAL experiment The characteristics of global lightning activities observed by ISUAL experiment

Alfred Bing-Chih Chen^{1*}, Yen-Jung Wu², Chih-Yu Chiang², Yi-Jen Lee², Jung-Kuang Chou², Li-Jou Lee², Cheng-Ling Kuo², Han-Tzong Su², Rue-Ron Hsu², Lou-Chuang Lee³

Alfred Bing-Chih Chen^{1*}, Yen-Jung Wu², Chih-Yu Chiang², Yi-Jen Lee², Jung-Kuang Chou², Li-Jou Lee², Cheng-Ling Kuo², Han-Tzong Su², Rue-Ron Hsu², Lou-Chuang Lee³

¹ISAPS, National Cheng Kung Univ., Taiwan, ²Physics, National Cheng Kung Univ., Taiw, ³ISS, National Central Univ., Taiwan

¹ISAPS, National Cheng Kung Univ., Taiwan, ²Physics, National Cheng Kung Univ., Taiw, ³ISS, National Central Univ., Taiwan

Lightnings serve as an important charge transporter between cloud and ground, and emit sferics that propagate in the ground-ionosphere cavity. During the six-year observation of ISUAL/FORMOSAT-2, besides surveying upper atmospheric transient luminous events, more than 110,000 lightnings which exceeded the ISUAL trigger threshold were recorded. In this presentation, the distribution, the occurrence rate, distribution and seasonal variation of these lightnings at local time between 22:30 and 23:00 are reported and compared with those of the LIS experiment (Christian et al., 2003). The ocean-to-land ratio and geographic distributions suggests that the lighting recorded by ISUAL is averagely more energetic than the ones registered by LIS mission and intense lightning is more frequent over oceans. The anomaly of occurrence between warm and cold phases of ENSO will also be discussed in this presentation.

キーワード: Lightning, ENSO, ISUAL

Keywords: Lightning, ENSO, ISUAL

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-04

会場:201A

時間:5月27日 09:15-09:30

Energetic radiation associated with thunderstorm activity Energetic radiation associated with thunderstorm activity

鳥居 建男¹, 鴨川 仁^{2*}, 杉田 武志³, 渡辺 泰行²

Tatsuo Torii¹, Masashi Kamogawa^{2*}, Takeshi Sugita³, Yasuyuki Watanabe²

¹ 日本原子力研究開発機構, ² 東京学芸大学物理学科, ³ 科学システム研究所

¹Fugen Decommissioning Eng. Center, JAEA, ²Dpt. of Phys., Tokyo Gakugei Univ., ³SSL

Intense gamma rays likely associated with lightning and/or thunderstorm activities have been detected in recent years, at various altitudes ranging from the ground level to the ionosphere. However, neither the source of this radiation nor its nature has been clarified. We report gamma rays lasting for several minutes attributed to both winter and summer thunderstorms in Japan. Our findings in winter thunderstorm indicate that the gamma rays were emitted continuously from a downward hemispherical surface, the bottom of which was about 300 m above sea level, and this source of gamma rays moved from north to south above the observation site at a speed of about 7 m/s. The radiation source probably moved along with the charged region of the cloud at a height of around 1 km, because the estimated migration of the radiation source was consistent with the observed movement of atmospheric electric field variation between ground-based observation sites and with the wind speed and direction at about 1 km altitude. This movement implies that the intense electric field produced by the charged region in the thundercloud generated a radiation source beneath the charged region. On the other hand, gradual energetic radiations probably caused by a summer thunderstorm have been observed at the top of Mt. Fuji, Japan. The largest of such variation was gradual and lasted for about 20 minutes, and was found to be high-energy gamma rays having a continuous energy spectrum up to 10 MeV or more. Both the observations help explain probably the lightning initiation urged by the radiation.

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-05

会場:201A

時間:5月27日 09:30-09:45

ISUAL recorded halos brightness and their parent lightning emission ISUAL recorded halos brightness and their parent lightning emission

Cheng-Ling Kuo^{1*}, Earle E. Williams², Jozsef Bor³, Gabriella Satori³, Toru Adachi⁴, Alfred Chen¹, Han-Tzong Su¹, Rue-Rou Hsu¹, Mitsuteru Sato⁵, Yukihiro Takahashi⁵

Cheng-Ling Kuo^{1*}, Earle E. Williams², Jozsef Bor³, Gabriella Satori³, Toru Adachi⁴, Alfred Chen¹, Han-Tzong Su¹, Rue-Rou Hsu¹, Mitsuteru Sato⁵, Yukihiro Takahashi⁵

¹Physics, Natl Cheng Kung Univ, Taiwan, ²Parsons Laboratory, MIT, USA, ³Hungarian Academy of Sciences, Hungary, ⁴Stanford University, USA, ⁵Cosmoscience, Hokkaido University, Japan

¹Physics, Natl Cheng Kung Univ, Taiwan, ²Parsons Laboratory, MIT, USA, ³Hungarian Academy of Sciences, Hungary, ⁴Stanford University, USA, ⁵Cosmoscience, Hokkaido University, Japan

Halo, another type of transient luminous event, is a bright disk at altitude 80-85 km above the thunderstorm. Unlikely the fact that almost exclusively +CG triggered sprites, most of halos were induced by -CG that occurred exclusively over the open water [Frey et al., 2007]. In this presentation, we analyzed total 185 ISUAL recorded pure halo events from July 2004 to Dec 2007. Using 1PN2-filter Imager, the average brightness of halos is ~ 0.25 MR. We also derived the current moment using the 777.4 lighting emission of their parent lightning to [Adachi et al., 2009]. It is found that a relatively strong linear relationship between lightning peak current and lightning-induced halo emission. In total recognized 121 events by ELF radio emission at Nagycenk Observatory (NCK), Hungary, the polarities of their parent lightning for 23 halos are identified as +CG while 98 halos are for ?CG associated with NCK recorded ELF data. From NCK estimating CMC, we found a relatively weak correlation between NCK CMC and halo brightness. Furthermore, the extremely brightest halos over ocean were also found, and their lightning polarities were dominated by ?CG. The finding reflects the nature of intense peak current for oceanic lightning [Fullekrug et al., 2002]. It seems that lighting current may have more important effect on halos generation than charge moment.

キーワード: ISUAL, Halo

Keywords: ISUAL, Halo

AEM002-06

会場:201A

時間:5月27日 09:45-10:00

航空撮像されたエルブスの高速度画像 High-speed imagery of elves from airplane

島 侑奈^{1*}, 高橋 幸弘¹, NHK 宇宙の渚プロジェクト²

Yuuna Shima^{1*}, Yukihiro Takahashi¹, NHK's project "Beach of the Cosmos"²

¹ 北海道大学大学院理学院宇宙理学専攻, ² NHK制作局 科学環境番組部

¹Dept. CosmoSciences, Hokkaido University, ²NHK

NHK 宇宙の渚プロジェクト

エルブスとは、雷放電に伴って起こる高高度過渡発光現象の一つである。高度約 90 km の電離圏下部で発生し、水平方向約 300-600 km の広がりを持ちパンケーキ状に発光する。またその発光継続時間は約 0.1 ms で、約 2 ms の間に光速を越えるような位相速度で全体に広がる、極めて高速な現象である。

これまでは、鉛直方向に並んだマルチアノードフォトメーターを用いた観測 [Fukunishi et al., 1996] や、垂直方向に視野を持つフォトメーターを用いた観測 [Barrington-Leigh et al., 2001] によってエルブス発光の時間・空間的变化を観ていたが、その構造の変化を二次元画像によって捉えられたものはなかった。

今回 NHK の協力のもとで高度約 13 km を飛行するジェット航空機から、世界で初めて高速度カメラでのエルブスの撮像に成功した。カメラは航空機の窓に進行方向と水平に設置され、約 8000 コマ/秒のフレーム数でパングロマトミック撮影を行った。

2010 年 11 月 28 日 18 時頃から千葉県東方沖約 400 km で雷放電が群発しており、本講演で示すデータはその雷放電によって引き起こされたエルブスを撮像したものである。この日航空機は 20 時半頃関東沖を目指して飛び立ち、21 時 48 分から 22 時 45 分までの約 1 時間撮像を行い、21 イベントの過渡発光現象を捉えた。そのうち、初期解析の段階で少なくとも 3 イベントのエルブスが検出されている。

本研究では、撮影されたエルブス構造の時間変化とそれを引き起こす親雷放電の特質との関係を調べていく。

AEM002-P01

会場:コンベンションホール

時間:5月27日 10:30-13:00

The use of cloud classification and rainfall radar data to improve geostationary satellite based rainfall estimation

The use of cloud classification and rainfall radar data to improve geostationary satellite based rainfall estimation

Dwi Prabowo Yuga Suseno^{1*}, Tomohito J. YAMADA¹

Dwi Prabowo Yuga Suseno^{1*}, Tomohito J. YAMADA¹

¹Grad. Sch. Eng. Hokkaido University

¹Grad. Sch. Eng. Hokkaido University

The use of geostationary satellite dataset for rainfall estimation has several advantages that it has a hemisphere coverage and high temporal resolution. However, we can only use Visible/Infra Red (VIS/IR) sensor that carried by geostationary satellite. Because of the cloud is opaque in VIS/IR spectral band, an indirect approach is used for rainfall estimation, i.e. according to several top surface cloud characteristics such as shape, brightness, temperature etc. Another rainfall estimation approach is by using Passive Micro Wave (PMW) sensor. The microwave spectral band has characteristic that can penetrate the cloud and interact with the hydrometeor. Those of characteristics make the PMW method more direct in term of rainfall estimation. PMW sensor usually mounted on polar orbit satellite, so it has limitation on temporal resolution and coverage. This study combines the advantage of geostationary satellite and PMW satellite images for rainfall estimation. We use MTSAT datasets that is blended with TRMM 2A12 to estimate the rainfall over Japan. We make a statistical relationship between cloud top temperature from MTSAT and rainfall rate from TRMM 2A12, according to assumption that on the convective cloud situation lower cloud top temperature is associated with higher rain rate. In the actual situation such assumption sometimes cannot be fulfilled. The cloud top temperature of the cirrus cloud i.e.: cold but not produces rain and the nimbostratus cloud i.e.: produces rain but warm have disturbed such relationship. The cloud classification according to the cloud type and cloud height will be performed. We use several cloud classification methods such as segmentation method, split-window method and maximum likelihood method to classify the cloud type. We investigate the statistical relationship among cloud classes and height to the rain rate. A calibration with the C-band rainfall radar data will also be conducted. The estimation result will be validated with the measured rainfall (Automated Meteorological Data Acquisition System/AMeDAS System). We expected that cloud classification based on cloud type and height as well as C-band rainfall radar calibration will improve the rainfall estimation accuracy.

キーワード: geostationary satellite, rainfall estimation, MTSAT, TRMM 2A12, cloud classification

Keywords: geostationary satellite, rainfall estimation, MTSAT, TRMM 2A12, cloud classification

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-P02

会場:コンベンションホール

時間:5月27日 10:30-13:00

A warm season climatology of convective precipitation over the Korean Peninsula A warm season climatology of convective precipitation over the Korean Peninsula

Yu-Kyung Hyun^{1*}, Hee-Jeong Baek¹, ChunHo Cho¹

Yu-Kyung Hyun^{1*}, Hee-Jeong Baek¹, ChunHo Cho¹

¹NIMR

¹NIMR

The goal of this study is to investigate the variations in the spatial and temporal patterns of lightning activity over the Korean Peninsula in relation to precipitation during the summer monsoon months during 10 years (2000-2009) and to develop a better understanding of these two meteorological phenomena. In this study, we present the results of an analysis of lightning activity and associated monsoon rainfall over Korea. We obtained precipitation data from 98 synoptic stations and the lightning data were collected from a lightning detection network installed by the Korean Meteorological Administration (KMA).

This study will be of use in understanding the role of convective rain in the extreme precipitation over Korea, and this could eventually enhance skills for understanding the relationship between climate change and extreme precipitation.

Acknowledgement: This research is supported by a project, NIMR-2011-B-2.

キーワード: extreme precipitation, Korean Peninsula, lightning activity, climate change

Keywords: extreme precipitation, Korean Peninsula, lightning activity, climate change

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-P03

会場:コンベンションホール

時間:5月27日 10:30-13:00

雷雲の全球活動マップと太陽活動の関係

Global map of thunderstorm activity based on GEON and its relationship to the solar activity

高橋 幸弘^{1*}, 山下 幸三¹, 佐藤 光輝¹, 宮原 ひろ子², 星野 直哉³

Yukihiro Takahashi^{1*}, Kozo Yamashita¹, Mitsuteru Sato¹, Hiroko Miyahara², Naoya Hoshino³

¹北海道大学 宇宙理学専攻, ²東京大学 宇宙線研究所, ³東北大学 地球物理学専攻

¹Hokkaido University, ²University of Tokyo, ³Tohoku University

Hokkaido University developed a global ELF observation network, named GEON, which provides very unique information of each cloud-to-ground lightning discharge (CG), as well as Schumann resonance (SR) power, a proxy of global energy proxy of lightning discharge. From the standpoint of the relationship between the effect of solar activity to the climate of Earth, lightning activity estimated by data obtained by GEON and the outgoing longwave Radiation (OLR), an indicator of cloud amount, are examined for their periodicity and phase in the periodic range of about one month. SR power shows about 27-day periodicity in solar maximum years and it becomes elongated toward solar minimum. On the other hand, OLR shows same kind of 27-day periodicity in solar maximum years, but only in the Western Pacific Warm Pool area. Both the spectra of SR and OLR have a peak around 35-day in solar minimum years. The average spectrum of OLR in solar maximum years also shows an enhancement in the range of 50-60 days corresponding to the main MJO period. In this presentation the relationship between the thunderstorm activity inferred from global lightning distribution observed by GEON and OLR are discussed in detail, comparing the solar activity.

キーワード: 雷放電, 積乱雲, OLR, 太陽活動, GEON

Keywords: lightning, thunderstorm, OLR, solar activity, GEON

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-P04

会場:コンベンションホール

時間:5月27日 10:30-13:00

全球雷活動の周期的変動と地域依存性

Periodic Changes of Global Lightning Activities and Their Regional Dependences

佐藤 光輝^{1*}, 高橋 幸弘¹, 山下 幸三²

Mitsuteru Sato^{1*}, Yukihiro Takahashi¹, Kozo Yamashita²

¹ 北海道大学大学院理学院, ² 東北大学大学院理学研究科

¹Dept. of CosmoSciences, Hokkaido Univ., ²Dept. of Geophysics, Tohoku Univ.

In order to study the periodic changes of global lightning activity and their regional dependences, we have analyzed ELF magnetic field waveform data obtained at Syowa station in Antarctica, Onagawa observatory in Japan and Esrange in Sweden for the period between February 2000 and December 2009. We have estimated day-to-day amplitude variation of the global lightning activity derived from Schumann resonance (SR) spectral power. As a next step, we have calculated power spectrum of the SR spectral amplitude variation to estimate periodicities using MEM, FFT, and wavelet method. It is found that the periodogram showed steep spectral peak at ~28-day in 2000-2001 which is the solar maximum period. On the other hand, a peaked period of the SR spectral amplitude variation gradually increased and showed a steep spectral peak over 30-days after 2002. Using the transient SR waveform data and newly developed geolocation method, we have also estimated the occurrence locations of intense lightning discharges for the period between September 2003 and August 2003. We will discuss the relationship between the periodic changes of regional lightning occurrence numbers and periodic changes of the regional lightning activity.

キーワード: 雷放電, シューマン共鳴, 周期変動

Keywords: lightning, Schumann resonance, periodic change

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-P05

会場:コンベンションホール

時間:5月27日 10:30-13:00

高速II-CCDカメラおよびハイビジョンカメラを用いた航空機からのスプライト観測の初期結果報告 Initial Results of Sprite Observation from Aircraft Using High-speed II-CCD camera and High-Vision Camera

佐藤 光輝^{1*}, 高橋 幸弘², 島 侑奈¹, 山下 幸三¹, NHK 宇宙の渚プロジェクト²

Mitsuteru Sato^{1*}, Yukihiro Takahashi², Yunna Shima¹, Kozo Yamashita¹, NHK Uchu-no-Nagisa Project²

¹ 北海道大学, ²NHK 宇宙の渚プロジェクト

¹Hokkaido University, ²NHK Uchu-no-Nagisa Project

In order to study spatial and time evolution of Transient Luminous Events (TLEs), such as sprites, elves, and blue jets, we have carry out TLE observations from a jet-aircraft using high-speed Image-Intensified (II) CCD camera, Watec CCD camera, EM-CCD camera and high-vision camera. On November 28, we have carried out a first try of the observation and captured about 30 TLE events successfully. At the presentation, we will show the initial results of the spatial and time evolution of sprites measured by high-speed and high-vision cameras.

キーワード: 雷, スプライト, 高速カメラ, ハイビジョンカメラ

Keywords: lightning, sprite, high-speed camera, high-vision camera

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AEM002-P06

会場:コンベンションホール

時間:5月27日 10:30-13:00

国際宇宙ステーションからの雷放電とスプライトの観測計画 Global Lightning and Sprite Measurements (GLIMS) from International Space Station

牛尾 知雄^{1*}, 佐藤 光輝², 森本 健志¹, 鈴木 睦³, 山崎 敦³, 高橋 幸弘², 芳原 容英⁴, 石田 良平⁵, 菊池 雅行⁶, 河崎 善一郎¹
Tomoo Ushio^{1*}, Mitsuteru Sato², Takeshi Morimoto¹, Makoto Suzuki³, Atsushi Yamazaki³, Yukihiro Takahashi², Yasuhide Hobara⁴, Ryohei Ishida⁵, Masayuki Kikuchi⁶, Zen-Ichiro Kawasaki¹

¹ 大阪大学, ² 北海道大学, ³ 宇宙航空研究開発機構, ⁴ 電気通信大学, ⁵ 大阪府立大学, ⁶ 極地研究所

¹Osaka University, ²Hokkaido University, ³JAXA, ⁴University of Electro-Communications, ⁵Osaka Prefecture University, ⁶NIPR

The Global Lightning and sprIte MeasurementS (GLIMS) on the International Space Station (ISS) is a mission to detect and locate optical transient luminous events (TLEs) and its associated lightning simultaneously from the non-sun synchronous orbit, and is scheduled to be launch from Japan in January, 2012 as part of the multi-mission consolidated equipment on Japanese Exposure Module (JEM). Our mission goals are (1) to detect and locate lightning and sprite within storm scale resolution over a large region of the Earth's surface along the orbital track of the ISS without any bias, (2) to clarify the generation mechanism of sprite, and (3) to identify the occurrence conditions of TLEs. To achieve these goals, two CMOS cameras, six Photometers, VLF receiver, and VHF interferometer with two antennas, are installed at the bottom of the module to observe the TLEs as well as causative lightning discharges at nadir direction during day and night time. Though the luminous events so-called sprite, elves and jets have been investigated by numerous researchers all over the world based mainly on the ground observations, some important problems have not been fully understood yet such as generation mechanisms of columniform fine structure and horizontal offset of some sprites from the parent lightning discharges. In the JEM-GLIMS mission, observations from our synchronized sensors are going to shed light on above-mentioned unsolved problems regarding TLEs as well as causative lightning discharges. In this presentation, the scientific background, instrumentation, project summaries are given.

キーワード: 雷, スプライト, 宇宙ステーション

Keywords: Lightning, Sprite, ISS

AEM002-P07

会場:コンベンションホール

時間:5月27日 10:30-13:00

庄内平野において観測された冬季雷の3次元標定 3D mapping of winter lightning discharges observed in the Shonai area

西橋 政秀^{1*}, 下瀬 健一¹, 楠 研一², 林 修吾², 新井 健一郎³, 猪上 華子², 益子 渉², 鈴木 修², 足立 啓二³, 別所 康太郎⁴, 星野 俊介², 中里 真久², 山内 洋², 保野 聡裕³, 加藤 亘³, 楠目 雅子¹

Masahide Nishihashi^{1*}, Ken-ichi Shimose¹, Kenichi Kusunoki², Syugo Hayashi², Kenichiro Arai³, Hanako Inoue², Wataru Mashiko², Osamu Suzuki², Keiji Adachi³, Kotaro Bessho⁴, Shnsuke Hoshino², Masahisa Nakazato², Hiroshi Yamauchi², Yoshihiro Hono³, Wataru Kato³, Masako Kusume¹

¹ アルファ電子 / 気象研究所, ² 気象研究所, ³ 東日本旅客鉄道, ⁴ 気象庁観測部

¹Alpha-denshi/MRI, ²Meteorological Research Institute, ³East Japan Railway Company, ⁴Japan Meteorological Agency

The Shonai area railroad weather project has investigated fine-scale structure of wind gust using two X-band Doppler radars and the network of 26 surface weather stations since 2007, in order to develop an automatic strong gust detection system for railroad. In 2009, the project was expanded and started lightning observation to investigate the mechanism of winter lightning and the application to strong gust prediction. Lightning discharge is known to be related to microphysical and dynamical processes within storms. Many scientists have indicated that lightning activity is associated with severe weather. Therefore, integration of continuous three-dimensional (3D) lightning monitoring (intracloud and cloud-to-ground lightning) and comprehensive high-density meteorological observation can provide useful index for predicting strong gust.

We developed a lightning observation system. The azimuth and elevation of VHF radiation sources originated from lightning flashes are computed using arrival time difference of three VHF pulses. After operation test at Meteorological Research Institute (MRI), we installed this system in the north of Shonai area (Ohama, Sakata) in October 2009. Moreover, we constructed three lightning observation sites in the Shonai area in September 2010, in order to visualize lightning discharges in 3D.

Our sensors detected lightning discharges at 01:13:32 JST on 4 December 2010. Using the VHF waveform data, we conduct 3D lightning mapping. The duration of discharge is about 60 ms and divided into two stages. The locations of discharges are compared with the radar echo data observed with two X-band Doppler radars in the Shonai area. As a result, the distribution of lightning discharges is consistent with the strong echo region. The lightning flash was also recorded with the network cameras at each site. In this presentation, we will show the lightning discharge process in detail.

Keywords: Winter lightning, 3D mapping, VHF observation, X-band radar, Shonai area