

SCG008-01

会場:301A

時間:5月26日 08:30-08:45

## 高温変成帯における複変成作用を検知するためのツールとしてのCHIME モナズ石年代

### CHIME monazite dating as a tool to detect polymetamorphism in high-temperature metamorphic terrane

河上 哲生<sup>1\*</sup>, 鈴木和博<sup>2</sup>

Tetsuo Kawakami<sup>1\*</sup>, Kazuhiro Suzuki<sup>2</sup>

<sup>1</sup> 京都大学, <sup>2</sup> 名古屋大学

<sup>1</sup>Kyoto University, <sup>2</sup>Nagoya University

The chemical Th-U-total Pb isochron method (CHIME) monazite dating was carried out for pelitic-psammitic migmatites and the Ao granite (one of the Younger Ryoke granites) from the Aoyama area, Ryoke metamorphic belt, SW Japan. The Ao granite gives an unequivocal age of 79.8±3.9 Ma. The migmatites yield ca. 100 Ma monazite grains with younger domains of ca. 80 Ma. Some grains show ca. 80 Ma overgrowths on the older core of around ca. 100 Ma, and others show patchy rejuvenation of ca. 100 Ma grains.

Recent study on the Pb diffusion in monazite (Cherniak et al., 2004) concluded that Pb diffusivity in monazite is low and thus the closure temperature of monazite is high. However, reinterpretation of Cherniak et al. (2004) data by favoring the SIMS data rather than RBS data showed that results of Cherniak et al. (2004), Suzuki et al. (1994) and Smith & Giletti (1997) can define single array that gives the diffusivity similar to Suzuki et al. (1994). If this interpretation is accepted, the closure temperature of Pb diffusion in monazite becomes low, and about 600-750 °C provided that the cooling rate is ca.30 °C/Myr.

With this closure temperature, it is possible to reset monazite in the pelitic-psammitic migmatites from the Aoyama area by the thermal effect of the Ao granite intrusion alone. The patchy nature of the distribution of rejuvenated parts within monazite grains prefers, however, the rejuvenation mechanism other than volume diffusion, such as grain boundary diffusion along the microcracks developed in the monazite grains. Some of the monazite grains with young overgrowth rim are armored in biotite, supporting the possibility of the fluid activity affecting the rejuvenation of monazite grains. Therefore, ca. 80 Ma overprinting on migmatites over the Grt-Crd zone in the Aoyama area is probably caused by the combination of a thermal effect and a fluid activity caused by the Ao granite intrusion. Partial rejuvenation of the monazite grains implies that the thermal and fluidal effects were not too strong to completely reset the monazite grains.

Although the contact metamorphism by the Ao granite is hard to be detected through the field survey and petrographic examination, possibly because the migmatite already possessed the high-temperature mineral assemblage and was immune from the contact metamorphism in terms of major metamorphic mineral assemblage, the CHIME monazite dating reveals the presence of contact metamorphism clearly. The field mapping of the CHIME monazite age can be a powerful tool for recognition of polymetamorphism in high-temperature metamorphic terrains where later thermal effects cannot be easily detected by the growth of new metamorphic minerals.

キーワード: CHIME モナズ石年代, 複変成作用, 青山高原地域, 阿保花崗岩, 接触変成作用, 閉鎖温度

Keywords: CHIME monazite dating, polymetamorphism, Aoyama area, Ao granite, contact metamorphism, closure temperature

SCG008-02

会場:301A

時間:5月26日 08:45-09:00

## Accretionary complex related to Paleo-Tethys subduction in northern Thailand Accretionary complex related to Paleo-Tethys subduction in northern Thailand

原 英俊<sup>1\*</sup>, 國井 美幸<sup>4</sup>, 鎌田 祥仁<sup>2</sup>, 上野 勝美<sup>3</sup>, 脇田 浩二<sup>1</sup>, 久田 健一郎<sup>4</sup>, Punya Charusiri<sup>5</sup>

Hidetoshi Hara<sup>1\*</sup>, Miyuki Kunii<sup>4</sup>, Yoshihito Kamata<sup>2</sup>, Katsumi Ueno<sup>3</sup>, Koji Wakita<sup>1</sup>, Ken-ichiro Hisada<sup>4</sup>, Punya Charusiri<sup>5</sup>

<sup>1</sup> 産業技術総合研究所, <sup>2</sup> 山口大学, <sup>3</sup> 福岡大学, <sup>4</sup> 筑波大学, <sup>5</sup> チュラロンコン大学

<sup>1</sup> Geological Survey of Japan, <sup>2</sup> Yamaguchi University, <sup>3</sup> Fukuoka University, <sup>4</sup> University of Tsukuba, <sup>5</sup> Chulalongkorn University

The Paleo-Tethys, which opened in response to the Devonian separation of the North China, South China and Indochina blocks from Gondwana, occupied a large area around the equator from the Devonian to Triassic, where carbonates, chert, and basalt were deposited in a pelagic domain (e.g., Metcalfe, 1999). These Paleo-Tethyan rocks characterized by Ocean Plate Stratigraphy were subducted beneath the Indochina Block during the Permian to Triassic (Wakita and Metcalfe, 2005).

In the Inthanon Zone of northern Thailand, melanges occur in association with oceanic rocks of Paleo-Tethys origin. According to Hara et al. (2009), melange formation was characterized by hydrofracturing and cataclastic deformation, with mud injection under semi-lithified conditions followed by shear deformation and pressure solution. Illite crystallinity data suggest metamorphic temperatures below 250 degree during melange formation. The combined structural and metamorphic data indicate that during melange formation, the accretionary complex related to Paleo-Tethys subduction developed at shallow levels within an accretionary prism. Asymmetric shear fabrics in melange indicate top-to-south shear. After correction for rotation associated with collision between the Indian and Eurasian continents, the trend of the Paleo-Tethys subduction zone is estimated to have been N80E. We conclude that the Paleo-Tethys was subducted northward beneath the Indochina Block from the Permian to Triassic. Sandstones are dominant within melange, generally characterized by lithic greywacke. Based on the composition and geochemistry of sandstones collected from melange within the Inthanon Zone, provenance of sandstone is characterized by mostly continental arc, and slightly continental margin. We interpreted that the origins of sandstones are from both of continent (Indochina Block) and arc developed along the Indochina Block, which is possibly the Sukhothai Zone. Based on melange kinematics and sandstone provenance, we reconstruct island arc system by the Paleo-Tethys subduction during Permian to Triassic time.

キーワード: 付加体, パレオテチス, メランジュ, 砂岩化学組成, インタノン帯, タイ

Keywords: accretionary complex, Paleo-Tethys, melange, sandstone geochemistry, Inthanon Zone, Thailand

SCG008-03

会場:301A

時間:5月26日 09:00-09:15

## タイ国中央部の地体構造区分：ペルム系および三畳系サクセションからの展望 Geotectonic subdivision of the Central Plain of Thailand: A perspective from Permian and Triassic successions

上野 勝美<sup>1\*</sup>, 宮東 照<sup>1</sup>, 鎌田 祥仁<sup>2</sup>, 加藤 京都<sup>2</sup>, Thasinee Charoentitirat<sup>3</sup>, Seranee Limruk<sup>3</sup>  
Katsumi Ueno<sup>1\*</sup>, Akira Miyahigashi<sup>1</sup>, Yoshihito Kamata<sup>2</sup>, Miyako Kato<sup>2</sup>, Thasinee Charoentitirat<sup>3</sup>, Seranee Limruk<sup>3</sup>

<sup>1</sup> 福岡大, <sup>2</sup> 山口大, <sup>3</sup> タイ国・チュラロンコン大  
<sup>1</sup>Fukuoka Univ., <sup>2</sup>Yamaguchi Univ., <sup>3</sup>Chulalongkorn Univ., Thailand

In the last two decades, increasing geological evidence produced a number of substantial understandings on the geotectonic subdivision and evolution of mainland Thailand. Now, we have a consensus of opinion that the Sukhothai Zone (Permian-Triassic island arc system) to the east and the Sibumasu Block (fragment of the eastern Cimmerian continent) to the west are the two major geotectonic domains constituting Northern Thailand. However, their southward extensions to the Central Plain of Thailand are still less clear due to the poor information of basement rocks.

We have investigated several basement rock units distributed in Uthai Thani and Nakhon Sawan provinces of the central part of the Central Plain. Quaternary deposits widely cover the major parts of these provinces, but there are a number of monadnocks consisting of older rocks. In eastern Nakhon Sawan Province, the basements are composed mostly of Permian platform-type carbonates. They are interpreted as parts of the Saraburi Limestone of the Indochina margin.

In central Nakhon Sawan and eastern Uthai Thani provinces, there are some, N-S trending monadnock ranges. Of them, we newly studied a carbonate unit composed of several limestone mountains. It consists of massive, shallow-marine limestone, and bioclastic grainstone and packstone are the major microfacies. In places, boundstone made by skeletal metazoans is observed. Late Triassic foraminifers occur from it. Based on the foraminiferal contents and lithology, it can be correlated to a limestone member of the Lampang Group in the Sukhothai Zone of Northern Thailand. In the same area, intermediate-acidic volcanic/volcaniclastic rocks of possibly Permian age are distributed. In Northern Thailand, a similar rock unit widely underlies the Sukhothai Zone.

Slightly to the east of the Late Triassic limestone, there are ranges of small mountains consisting mainly of fine-grained siliciclastics. Characteristically this unit contains chert successions in some parts; one of which distributed around Nakhon Sawan city is called the Khao Gob Chert. Permian radiolarians are found from some of the chert successions. Their lithological characters and age suggest that they can be correlated to the Permian Khanu Chert exposed to the north in the Sukhothai Zone. Moreover, granitoids in the same mountain ranges are of I-type characterizing the Eastern Granitoid Belt, which corresponds in Northern Thailand to the Sukhothai Zone.

In western Uthai Thani Province, there is a remarkable range consisting of karstified limestone, which is called the Uthai Thani Limestone. It is generally weakly metamorphosed and deformed, but Permian age is evident based on the occurrence of rare and poorly preserved fusulines from some localities. The Uthai Thani Limestone can be distinguished from the Late Triassic limestone just to the east by its distinctive NNW-SSE extension and by having different deformation and metamorphic records. It is more reasonable to correlate it with the Sai Yok and the Ratburi limestones widely distributed in the Sibumasu Block of Western and Peninsular Thailand. Moreover, granite in western Uthai Thani Province exhibits an S-type affinity of the Central Granitoid Belt, which corresponds to the eastern part of the Sibumasu Block.

All these lines of evidence suggest that central Nakhon Sawan and eastern Uthai Thani provinces belong to the southern extension of the Sukhothai Zone of Northern Thailand. In contrast, western Uthai Thani Province where the Uthai Thani Limestone and the S-Type granitoids crop out can be labeled as a part of the Sibumasu Block. Thus, we conclude that, in the Nakhon Sawan-Uthai Thani area there is a remarkable gap in terms of geotectonic properties of basement rocks between the Late Triassic limestone and the Uthai Thani Limestone. It corresponds to the southern extension of the Chiang Rai Tectonic Line in Northern Thailand, which separates the Sukhothai Zone to the east and the Sibumasu Block to the west.

キーワード: タイ国中央部, 地体構造区分, スコタイ帯, シブマス地塊, ペルム系, 三畳系

Keywords: Central Thailand, geotectonic subdivision, Sukhothai Zone, Sibumasu Block, Permian, Triassic

SCG008-04

会場:301A

時間:5月26日 09:15-09:30

## Evolution of Magmatism and Metallogeny along the Western Edge of the Indochina Terrane Evolution of Magmatism and Metallogeny along the Western Edge of the Indochina Terrane

チャルシリ パンニャ<sup>1\*</sup>

Punya Charusiri<sup>1\*</sup>

<sup>1</sup> チュラロンコーン大学

<sup>1</sup>Chulalongkorn University

K. Kampawong (MIH, Laos), S. Meffree (Univ. Tasmania), A. Salum (Univ. Tasmania), T. Y. Lee (NTNU, Taiwan), C. H. Lo (NTU, Taiwan), K. Hisada (Univ. Tsukuba), W. Lunwongsa (Issara Min. Com.)

Igneous rocks and their associated mineral deposits form north-south trending belt with the dimension of about 1700 x 100 km<sup>2</sup> along the edge of the Khorat Plateau from central Lao PDR to southeastern Thailand. These igneous rocks are predominantly felsic to intermediate plutonic and volcanic rocks with minority of mafic and ultramafic exposures. The latter which are usually Late Paleozoic in ages are regarded as to be temporally and spatially associated with the Loei suture which subdivided Indochina terrane from the Paleotethys Ocean. Based on U-Pb zircon, Rb-Sr whole rock, and <sup>40</sup>Ar-<sup>39</sup>Ar dating, magmatism and associated mineralization are inferred to have a prolong history of tectonism from compression to extension and ranging in ages from Middle Paleozoic to Miocene. The first stage of tectono-magmatic episode includes calc-alkaline magmatism during Middle Paleozoic (ca. 405 ? 399 Ma) ages and associated with their mild Kuroko type mineralization of Fe, Cu, Pb, Zn, Ba and Mn minerals. They are mainly in the northern part of the belt and may have been part of Caledonian Orogeny. This igneous activity was formed as a result of calc-alkaline magmatism due to subduction of paleothethystic oceanic slab beneath the western edge of Indochina terrane. The second stage of magmatism took place within the early Indosinian Orogeny (ca. 260 ? 240 Ma). The igneous rocks are of calc-alkaline I-type affinity and thought to have been responsible for the late stage mineralization including porphyry copper, volcanogenic sulfide, skarn type, and hydrothermal Au-Ag vein styles which are mainly occurred in the northern belt. Such the strongly mineralized magmatism is believed to have formed in response to the continuous oceanic subduction beneath the Indochina terrane. Though associated unexplained Sn mineralization was reported in Lao PDR, it was virtually absent in Thailand. Associated accretionary complex and weak Cyprus type Cr and Ni mineralization may also have formed at this stage. The third stage of magmatism may have occurred in the very late stage of Indosinian Orogeny (ca. 190-210 Ma), and it was thought to have occurred in the extension environment giving rise to mainly alkaline magmatism, particularly along the suture zone. Stitching plutons with weak poly metallic vein-style mineralization may have formed due to the relaxation of the crust after intense compression. The fourth stage of magmatism may have formed in association with reactivation of major strike ? slip northwest ? trending faulting in response to early Himalaya Orogeny. Mylonitization of earlier granitoid rocks may have been occurred at this stage along with the post tectonic minor intrusions of water poor pegmatites and aplites with weak vein-style mineralization of Sb + Au-Ag hydrothermal ores. The fifth stage of magmatism is found in the central belt, and it was formed as a result of crustal relaxation during Miocene time (18-25 Ma). This rift related alkaline magmatism may have formed due to crustal thinning, and mantle upwelling by high heat flow activity may be responsible for partial melting of the lower crustal materials. Poly-metallic mineralization and Au-Ag veins of high sulfidation systems may have took place along the north-south ? trending open fractures, particularly in the central to eastern Thailand. Continental rifting event may have been continued to the last episode of magmatism during ca. 0.5 to 7 Ma along the Loei suture in response to high heat flow by upwelling mantle materials. Tectonic activity is geomorphologically and geochronologically evident to continue to the present time and gives rise to the presence of reactivated northwest ? trending active faults in the central belt. Only non-mineralized hydrothermal veins are encountered along these faults.

Keywords: Indochina, Metallogeny, Tectonic, Magmatism, Thailand, Laos

SCG008-05

会場:301A

時間:5月26日 09:30-09:45

## 更新された北ベトナム地質；古期の付加体を突き破った中新世変成岩ドーム，デタッチメント正断層としての紅河断層 Renewed geology of northern Vietnam: Miocene metamorphic domes extruded the older accretionary complexes

遅沢 壮一<sup>1\*</sup>, Nguyen Van Vuong<sup>2</sup>, Vu Van Tich<sup>2</sup>  
Soichi Osozawa<sup>1\*</sup>, Nguyen Van Vuong<sup>2</sup>, Vu Van Tich<sup>2</sup>

<sup>1</sup> 東北大学理学研究科地学専攻, <sup>2</sup>Hanoi University of Science  
<sup>1</sup>Tohoku University, <sup>2</sup>Hanoi University of Science


北ベトナムの基本地質構造は、背斜軸に変成岩や古期岩があり、それを取り巻いて、順次、整然とより新しい地層が分布するマップスケールの背斜として、地質図に図示されている。また、紅河断層など大小の横ずれ断層が多数、地層境界を含めて、図示されている。しかし、ここ数年の現地地質調査から、これらは空想であることが明らかになった。

紅河断層とソンチャイ断層に挟まれた変成岩ドームには、いくつかの、互いに見解が異なる研究があるが、筆者らはそれらとセンスが異なる右ずれ成分を伴うトランスプレッションの場での、軸面フォリエーションやそのセンスの伸長線構造を伴うD1ドーム（大規模背斜）と解析した。これのミニチュア褶曲が観察される大露頭を、カバー堆積物であるが、発見、記載している。D2は変成岩ドーム縁に観察される脆性断層で、結局、紅河断層とソンチャイ断層は変成岩エクシユージョンの最終段階を表すデタッチメント正断層として機能している。大規模横ずれ断層である根拠は、露頭観察からは得られていない。

ドームや正断層の上盤にあるカバー堆積物は、カンブリア紀から白亜紀まで順次重なる正常層ではない。基底不整合も保存されている白亜紀の赤色層を除いて、すべて付加体由来と考えられる。膨大な量の石灰岩や玄武岩の分布は例えば、三疊系として、マッピングされているが、遠洋性チャート、アンバー、半遠洋性多色泥岩など、付加体を特徴付ける岩石を、ベトナムで初めて発見した。例えばデボン系とされているタービダイトは海溝充填堆積物で、多くは海洋性岩石からなるブロックの基質をなすと考えられる。ただし、付加帯形成時の構造はほぼ消失していて、これらカバー堆積物の構造はドーム変成岩の構造と、大差ない。つまり、中新世の変形は、古期のカバー堆積物にも強く及んでいる。なお、化石産地は従来の地質図に多数記されているが、筆者らは1地点でペルム紀フズリナ化石を発見したに留まっていて、これは付加体であることに加えて、強烈な中新世変成作用を反映している。



Moetamorphic dome model



Miniture of metamorphic dome

Anticlinal fold accompanies axial planar cleavages, dextral stretching lineations, and parasitic folds

キーワード: 北ベトナム, 変成岩ドーム, D1 トランスプレッション, 右ずれ, D2 紅河断層, 付加体起源

Keywords: Northern Vietnam, metamorphic dome, D1 trans pression, dextral, D2 Red River normal fault, accretionary prism origin

SCG008-06

会場:301A

時間:5月26日 09:45-10:00

## Insight into the tectonic deformation patterns along eastern Taiwan from seismological observations

## Insight into the tectonic deformation patterns along eastern Taiwan from seismological observations

Wen-Tzong Liang<sup>1\*</sup>

Wen-Tzong Liang<sup>1\*</sup>

<sup>1</sup>Inst. Earth Sciences, Academia Sinica

<sup>1</sup>Inst. Earth Sciences, Academia Sinica

To decipher the seismotectonic structures in the Taiwan region, we have obtained more than 1,900 centroid moment tensor (CMT) solutions by inverting the broadband waveforms recorded at the BATS network. This dataset reveals the characteristics of faulting types along the eastern coast of Taiwan that is associated with the interaction between the continental margin and the NW moving Philippine Sea plate (PSP). In general, to the south of Hoping (24.3°N), the faulting type is dominated by thrusting and the sigma-1 direction is almost perpendicular to the strike of Coastal Range. However, between Hopin and Lanyang river, the strike-slip faulting type is prevailing along some EW trending geological boundaries in the crust with sigma-1 directed in NW-SE. Some low-angle dip-slip events at depth around 20 km were observed in the vicinity of Hopin. We infer that they may correspond to the interface earthquakes, where the underneath subducted slab moves westward. To the north of Lanyang river, the crust is dominated by normal faulting, which is believed to be caused by the spreading of the Okinawa Trough. On the other hand, for those intermediate depth earthquakes occurred beneath Ilan plain, the determined focal mechanisms show thrusting pattern with sigma-1 sub-parallel to the trench. Therefore, Kao and Jian(2001) suggested that the subducted PSP slab is colliding with the mantle lithosphere of EUP. Chou et al.(2006) also discovered the folding at the western edge of slab. Based on seismicity and 3-D velocity structures, Wu et al.(2009) proposed that the intersection of the Ryukyu Trench with Taiwan is placed at the latitude of 23.7°N, placing the northern part of the Coastal Range on EUP. This is consistent with the uplifting rate along the Coastal Range and the modeled deformation pattern. According to these observations, we concluded that the PSP starts subducting at 23.7°N and the upper plate boundary approaches the bottom of the brittle seismogenic layer at about 24.3°N, where very low-angle dip-slip events may occur. Once the slab subducts completely underneath the seismogenic layer, the faulting type in the crust would be dominated by pre-existed boundaries subjected to a torque that may be imposed by the trench retreat and the opening of back-arc basin. Behind the Lanyang river, the Okinawa Trough plays an important role for the occurrence of shallow normal faulting events.

キーワード: CMT, Focal Mechanism, Seismotectonics, Stress Tensor, Collision, Subduction

Keywords: CMT, Focal Mechanism, Seismotectonics, Stress Tensor, Collision, Subduction

SCG008-07

会場:301A

時間:5月26日 10:00-10:15

## Petrology of eclogites from the Orobashy area, Aktyuz region, northern Kyrgyz Tien-Shan

## Petrology of eclogites from the Orobashy area, Aktyuz region, northern Kyrgyz Tien-Shan

Orozbaev Rustam<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Apas Bakirov<sup>2</sup>, Kadyrbek Sakiev<sup>2</sup>, Michio Tagiri<sup>3</sup>

Orozbaev Rustam<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Apas Bakirov<sup>2</sup>, Kadyrbek Sakiev<sup>2</sup>, Michio Tagiri<sup>3</sup>

<sup>1</sup>Dept. of Geosci., Shimane University, <sup>2</sup>Inst. Geol. of Kyrgyz Academy of Science, <sup>3</sup>Inst. Res. Off., Ibaraki University

<sup>1</sup>Dept. of Geosci., Shimane University, <sup>2</sup>Inst. Geol. of Kyrgyz Academy of Science, <sup>3</sup>Inst. Res. Off., Ibaraki University

The Aktyuz region is located in the Zaili Range of the Northern Kyrgyz Tien-Shan. Here, eclogites have been reported from two areas, i.e. Aktyuz area and Orobashy area (Orozbaev et al., 2010). Lens-shaped serpentinite bodies of varying size occur in the Orobashy area along NNE-SSW trending faults located within Kapchygai and Kokbulak migmatites. The largest serpentinite body contains 20 cm to 30 m across blocks of eclogites, garnet amphibolites, garnetites and gneisses. Eclogite, garnetite and garnet amphibolite blocks are massive, and range from 40 cm to 10 m in diameter. The marginal parts of the eclogite and garnetite blocks are amphibolitized. Both blocks and serpentinite matrix are intruded by granitic dykes.

Eclogites consist mainly of garnet, clinopyroxene, amphibole, clinozoisite and rutile with minor plagioclase, epidote, quartz, chlorite, biotite, paragonite, phengite, ilmenite, titanite, apatite and zircon. Garnets and clinopyroxene show coexisting texture and thus define the peak metamorphic assemblage. Garnets contain inclusions of amphibole (tschermakite, pargasite), quartz, epidote, rutile, ilmenite, apatite, plagioclase, biotite and rare paragonite and phengite. Garnets show a prograde zoning and they have compositions of 36.1-61.3 mol% of almandine, 13.3-21.9 mol% of grossular, 10.8-43.4 mol% of pyrope and 0.1-4.9 mol% of spessartine. Clinopyroxenes are Na-rich diopside in composition and their jadeite content varies from 0 to 19 mol%. The symplectite of Amp+Qtz and/or Amp+Pl is developed after clinopyroxene suggesting decompressional decomposition. Amphiboles occur as inclusions in garnet and clinopyroxene, and also in the matrix. Amphibole in the matrix is Mg-hornblende and contains inclusions of garnet, clinopyroxene (Jd9-12), rutile and quartz. Amphiboles in garnets are tschermakite and pargasite, whereas those in clinopyroxene are Mg-hornblende. Amphiboles in symplectites and in the veins are also classified as Mg-hornblende. Clinopyroxene, garnet and amphibole are cross cut by veins. Feldspars in the eclogites are plagioclase (An4-50) and K-feldspar. Paragonite occurs only as inclusions in garnets. Phengite occurs in the matrix and as rare inclusions in garnet. Epidote occurs as inclusions in garnets, clinopyroxenes and in the veins. Occasionally, clinozoisite develops in the matrix of some eclogites. Veins are mainly composed of plagioclase, epidote, amphibole and apatite.

The texture and the mineral chemistry of the Orobashy eclogites suggest three metamorphic stages i.e., prograde, peak and decompression stages. The mineral inclusions in garnet such as amphibole (tschermakite and pargasite), quartz, plagioclase (An13-50), epidote, biotite, paragonite, phengite, rutile, ilmenite and apatite may suggest mineral assemblage of the amphibolite facies conditions for the prograde stage. The coexistence of garnet and Na-rich diopside together with rutile and quartz in the matrix suggests the peak metamorphic assemblage. The garnet-clinopyroxene geothermometer of Ravna (2000) and Nakamura (2009) yielded  $T = 590-710$  C at 13 kb, minimum pressure based on Jd=19 content in clinopyroxene. The symplectite of Amp+Qtz and Amp+Pl suggest decompression stage for the eclogites. Hbl-Pl geothermometer (Holland & Blundy, 1994) yielded 610-670 °C for this stage, and the minimum Jd content (>5) may suggest about 5 kbar for pressure conditions.

There are three different types of metamorphic events for the HP rocks of the Aktyuz Formation. Here, for the Orobashy eclogites single metamorphic event is recognized, which is HT metamorphism from amphibolite facies up to eclogite facies conditions and then decompression again to amphibolite facies conditions. Exhumation of the Orobashy eclogites to relatively shallow crustal levels was accompanied by re-equilibration under HT amphibolite facies conditions, probably during a period of continental collision, correlative to the third metamorphic event in the Aktyuz area.

キーワード: Orobashy, Aktyuz, eclogite, Tien-Shan, Kyrgyzstan

Keywords: Orobashy, Aktyuz, eclogite, Tien-Shan, Kyrgyzstan

## モンゴル中北部ハラホリン地域のペルム紀周氷河堆積物 Permian peri-glacial deposits in the Harahorin area, north-central Mongolia

大藤 茂<sup>1\*</sup>, 藤本 辰弥<sup>1</sup>, 東田 和弘<sup>2</sup>, 下條 将徳<sup>1</sup>, セルスマ・ゴンチクドルジ<sup>3</sup>  
Shigeru Otoh<sup>1\*</sup>, Tatsuya Fujimoto<sup>1</sup>, Kazuhiro Tsukada<sup>2</sup>, Masanori Shimojo<sup>1</sup>, Sersmaa Gonchigdorj<sup>3</sup>

<sup>1</sup> 富山大学大学院理工学研究部, <sup>2</sup> 名古屋大学博物館, <sup>3</sup> モンゴル科学技術大学

<sup>1</sup>University of Toyama, <sup>2</sup>The Nagoya University Museum, <sup>3</sup>Mongolian Univ. Sci. & Tech.

はじめに モンゴル中北部のハラホリン地域は、前期石炭紀付加体が分布するハンガイ - ヘンテイ帯 (Kurihara et al., 2009) に位置する。当地域で付加体を被覆するウルメグテイ層には、ドロップストーンを含む周氷河堆積物が含まれる。この度、この周氷河堆積物に関する地質学的・年代学的研究をしたので、ここにその成果を発表する。

地質概説 研究対象地域は、主に以下の地質単元からなる。

ヤシル層：ハンガイ - ヘンテイ帯の前期石炭紀付加体。

ウルメグテイ層：ヤシル層を被覆すると見られる地層。下位より礫岩層、砂岩層、花崗岩ドロップストーンを含むヴァーヴ状砂岩泥岩互層と、上方細粒化する。全体的に変形を受けており、下位ほど顕著な構造性面構造と伸長性線構造を有する。構造性面構造は北西走向・南西傾斜で、伸長性線構造は南北トレンドである。剪断センス指示物から、これらの変形構造は右横すべり成分をもつ北方への衝上性剪断により形成されたものと考えられる。

珪長質岩脈：ウルメグテイ層に貫入し、一部、ウルメグテイ層と共に上記の剪断変形を蒙っている。

花崗岩バソリス：ウルメグテイ層の変形構造を切って貫入する、非変形斑状花崗岩。

年代測定 京都大学平田研究室および東京大学地震研究所設置のレーザー照射型誘導結合プラズマ質量分析計 (LA-ICP-MS) を用いて、以下の試料より抽出したジルコンを分析し、ウラン - 鉛年代を求めた。

ウルメグテイ層中の花崗岩ドロップストーン：多くのジルコンは包有物を含み、細かい破断面をもち、分析に適さなかった。そこで、例外的にきれいなジルコン 3 粒子を分析し、<sup>206</sup>Pb/<sup>238</sup>U 年代の加重平均値をとった結果、273.0 ± 4.9 Ma という年代値が得られた。

ウルメグテイ層に貫入する変形した珪長質岩脈：ジルコン 17 粒子から 246.5 ± 4.0 Ma の <sup>206</sup>Pb/<sup>238</sup>U 加重平均年代を得た。

花崗岩バソリス：ジルコン 10 粒子から、217.5 ± 8.7 Ma の <sup>206</sup>Pb/<sup>238</sup>U 加重平均年代を得た。

考察

地史概要：ウルメグテイ層中の花崗岩ドロップストーンおよびウルメグテイ層を切る珪長質岩脈の年代から、ウルメグテイ層の堆積年代は 278-242 Ma (前期ペルム紀アーティンスク期～中期三畳紀アニサス期に相当；ICS, 2009) に拘束される。上記珪長質岩脈 (246.5 ± 4.0 Ma) は、ウルメグテイ層と共に剪断変形を受けている。従って、この剪断変形は 242 Ma より前に始まっており、251 Ma 以降まで継続したことがわかる。更に 217.5 ± 8.7 Ma の花崗岩バソリスが変形していないことから、ウルメグテイ層の剪断変形は、251 Ma (前期三畳紀インダス期；ICS, 2009) 以降まで継続し 208 Ma (後期三畳紀ノール期；ICS, 2009) には終わっていたと考えられる。

ウルメグテイ層周氷河堆積物の堆積年代と意義：後期石炭紀からペルム紀初頭にかけては、汎世界的な氷河時代で、南半球の Gondwana 大陸は広く氷河に覆われた。前期ペルム紀末には Gondwana の氷河も縮小し、以後は中生代に向けて地球は徐々に温暖化したとされる。アジアでも、278-242 Ma の氷河堆積物は、ベルホヤンスク、コリマ、オモロン地域など、シベリアの北東方でのみ知られている (Ustritskiy, 1973)。例えば、オモロン地域ルースカヤ川に分布する上部ペルム系ウーチャピン階の周氷河堆積物が分布し、その中のドロップストーンの産状および礫種構成 (Mikhaylov et al, 1969) は、ウルメグテイ層と類似する。また、主要な礫種は異なるが、類似する産状を示す周氷河堆積物は、ベルホヤンスク地域およびコリマ地域の中部ペルム系ワード階～上部ペルム系ウーチャピン階の中にも見られる。従って、今回得られた年代値およびアジアの周氷河堆積物の分布・年代から考えて、ウルメグテイ層は中期ペルム紀ワード期から後期ペルム紀ウーチャピン期 (268.0-253.8 Ma；ICS, 2009) に堆積したと判断される。ウルメグテイ層に覆われるハンガイ - ヘンテイ付加体は、これら中期～後期ペルム紀に氷河の発達した大陸または島弧と連続するシベリア東縁に付加し、中期ペルム紀ワード期から後期ペルム紀ウーチャピン期には現シベリア北東部のオモロン地域付近に位置していた可能性が大きくなる。ハラホリン地域とオモロン地域は現在約 2,000 km 離れているため、ハンガイ - ヘンテイ付加体とウルメグテイ層は、ワード期以降に再配列を受けて、後期三畳紀ノール期までに現在の位置に移動してきたと考えられる。ウルメグテイ層に見られる、前期三畳紀インダス期前後の右横すべり成分をもつ衝上性剪断運動は、この再配列の際にできた可能性がある。



キーワード: ペルム紀, 周氷河堆積物, ウラン - 鉛年代, テクトニクス, ハンガイ - ヘンテイ帯, モンゴル  
Keywords: Permian, peri-glacial deposit, U-Pb age, tectonics, Khangai-Khentei belt, Mongolia

SCG008-09

会場:301A

時間:5月26日 10:45-11:00

## Significance of metacarbonate rocks in unraveling the formation of Central Asian Orogenic Belt

### Significance of metacarbonate rocks in unraveling the formation of Central Asian Orogenic Belt

Satish-Kumar Madhusoodhan<sup>1\*</sup>, Yasuhito Osanai<sup>2</sup>, Nobuhiko Nakano<sup>2</sup>, S. Jargalan<sup>3</sup>, C. Boldbaatar<sup>4</sup>, Masaaki Owada<sup>5</sup>, Tatsuro Adachi<sup>2</sup>, Kazuhiro Yonemura<sup>2</sup>, Aya Yoshimoto<sup>2</sup>  
Satish-Kumar Madhusoodhan<sup>1\*</sup>, Yasuhito Osanai<sup>2</sup>, Nobuhiko Nakano<sup>2</sup>, S. Jargalan<sup>3</sup>, C. Boldbaatar<sup>4</sup>, Masaaki Owada<sup>5</sup>, Tatsuro Adachi<sup>2</sup>, Kazuhiro Yonemura<sup>2</sup>, Aya Yoshimoto<sup>2</sup>

<sup>1</sup>Shizuoka University, <sup>2</sup>Kyushu University, <sup>3</sup>Mongolian Univ. Sci. Tech., <sup>4</sup>Mineral Resources Authority of Mongolia, <sup>5</sup>Yamaguchi University

<sup>1</sup>Shizuoka University, <sup>2</sup>Kyushu University, <sup>3</sup>Mongolian Univ. Sci. Tech., <sup>4</sup>Mineral Resources Authority of Mongolia, <sup>5</sup>Yamaguchi University

Metacarbonate rocks are key rock units in orogenic belts because they represent possible remnants of oceanic sediments. Furthermore, they provide important information regarding the fluid-rock history during metamorphism. Central Asian Orogenic Belt (CAOB) consists of several subduction-accretion complexes formed between 1000 Ma and 250 Ma. Metamorphic rocks in this region are important not only in understanding the tectonic evolution of the CAOB, but also can give important clues in understanding the formation of Asian continent. During the Japan-Mongolia Joint Geological Research we have collected several key rock units from the western and northwestern regions of Mongolia. Here we present carbon and oxygen stable isotope characteristics of metacarbonate rocks from the several localities of Central Asian Orogenic Belt of Mongolia. Metacarbonate rocks are common in the CAOB and occur as comparatively thin layers intercalated with pelitic and psammitic gneisses.

Carbon and oxygen isotope studies on metacarbonate rocks shows a large spread in isotope values among different belts within the CAOB. However, the coupled C-O isotope trend can be broadly grouped into two categories. (1) Metacarbonate rocks that preserve pre-metamorphic stable isotope signatures, and (2) those affected by fluid-rock interaction processes during metamorphism. In the first case, we were able to obtain carbon and oxygen isotopic composition that might be representative of sedimentary signatures, although isotopic shifts caused by the exchange of carbon isotopes with graphite derived from organic material is present in some samples. The equilibrium between calcite and graphite in such samples were used to understand the temperature of metamorphism in such samples. In the second case, we have got large coupled carbon and oxygen isotope shifts which are indicative of fluid-rock interaction during metamorphism. Such samples helped to understand the role of fluids during subduction-accretion processes during the formation of CAOB.

キーワード: metacarbonate rocks, carbon isotopes, oxygen isotopes, Central Asian Orogenic Belt, Mongolia  
Keywords: metacarbonate rocks, carbon isotopes, oxygen isotopes, Central Asian Orogenic Belt, Mongolia

SCG008-10

会場:301A

時間:5月26日 11:00-11:15

## Geodynamics and ore mineralization in Bayankhongor metallogenic belt, Central Mongolia Geodynamics and ore mineralization in Bayankhongor metallogenic belt, Central Mongolia

Sereenen Jargalan<sup>1\*</sup>, 小山内 康人<sup>2</sup>, 中野 伸彦<sup>2</sup>, 足立 達朗<sup>2</sup>

Jargalan Sereenen<sup>1\*</sup>, Yasuhito Osanai<sup>2</sup>, Nobuhiko Nakano<sup>2</sup>, Tatsuro Adachi<sup>2</sup>

<sup>1</sup> モンゴル科学技術大学, <sup>2</sup> 九州大学

<sup>1</sup>Mongolian University of Science & Techno, <sup>2</sup>Kyushu University

Bayankhongor metallogenic belt is one of the main gold producing area in Mongolia. In the regional geotectonic setting the Bayankhongor metallogenic belt is north-west trending tectonic zone, which consists of several tectonic units as Archean Baidrag terrane, Burd metamorphic unit, Bayankhongor ophiolite belt, Zag metamorphic belt and southern part of Hangai intra-continental basin. The tectonic zones of Bayankhongor belt has several episodes magmatic activity starting from Proterozoic continuing to Cambrian, Early Paleozoic, Late Paleozoic, and Mesozoic.

The Bayankhongor belt is known by its gold mineralization since ancient time, and there are remnants of old mining activity at the several places. Ore mineralization is closely associated with long history of geodynamic setting of the belt starting from old oceanic crust through accretion, collision and obduction as well as intra-continental magmatic activity.

Results of various stage geological stud, indicate that there are many deposits and occurrences of gold, copper, tungsten, iron, nickel and PGE etc. Copper, nickel and PGE mineralization thought to be related with mafic and ultramafic units of ophiolite belt, and formed during oceanic setting. Gold bearing quartz vein zones, and cold copper bearing scarn deposits, which must be primary source of placer deposits, are mainly distributed along the south and south west part of the belt.

Placer deposits mainly found along the along the mountain Jargalant, which is obducted ophiolite zone, and Burd passive continental margin terrane and Bombogor Proterozoic metamorph terrane. Age of placer deposits ranges from Mesozoic to Recent.

We study age and geochemistry of magmatic rocks which are distributed in and around the ore mineralization occurrences, in order to make clear its genetic relation and as well as reconstruction of tectonic evolution of Bayankhongor metallogenic belt. Result petrochemistry and of age dating analyses will be discussed and presented.

Keywords: Bayankhongor, ophiolite, gold, placer, PGE, magmatic rocks

SCG008-11

会場:301A

時間:5月26日 11:15-11:30

## 伊豆弧衝突帯における大陸地殻形成 Formation of continental crust at the Izu-Honshu collision zone

田村 芳彦<sup>1\*</sup>

Yoshihiko Tamura<sup>1\*</sup>

<sup>1</sup> 海洋研究開発機構地球内部ダイナミクス領域

<sup>1</sup>IFREE, JAMSTEC

The tectonic setting of arc-arc collision and arc accretion in the Izu-collision zone is similar to that of Archean orogenic belts (e.g. Taira et al., 1992). Understanding the petrological processes of granite formation in the Izu-collision zone, where the geodynamic information is not modified by polyphase deformation and metamorphism, may contribute to understanding ancient orogenic belts, especially those related to collisional settings. The Pacific plate began subducting beneath the Philippine Sea plate about 50 million years ago to produce the currently active Izu-Bonin-Mariana (IBM) arc. The collision between the northern IBM arc system and the Honshu arc of the Eurasia plate has been occurring since the middle Miocene (c. 15 Ma) as a consequence of the northwestward migration of the Philippine Sea plate. The collision has led to the accretion of IBM arc crust to the Honshu arc, associated with a southward migration of the plate boundary and trench system. Neogene granite plutons are widely exhumed by tectonic uplift associated with arc collision. Tamura et al. (2010) integrated new geochemical results with recent geophysical imaging of the arc and concluded that Miocene plutonic rocks in the Izu collision zone were derived from the Eocene-Oligocene middle crust, which was partially melted, remobilized and rejuvenated during the collision. Similar rejuvenating processes may have operated in other collision zones. Moreover, (1) the mafic arc lower crust is missing at the collision zone (Kitamura et al., 2003) and (2) the aseismic Philippine Sea plate, which is subducting to depths of 130-140 km without evidence of a tear or other gap, has been detected even beneath areas 120 km NW of the collision zone (Nakajima et al., 2009). Thus, collisional coalescence accompanies delamination of arc mafic lower crust. Both processes are inevitable at the collision and necessary to yield continental crust.

キーワード: かこう岩, 中部地殻, 地殻構造, 島弧マグマ, 安山岩

Keywords: granites, middle crust, crustal structure, arc magmas, andesite

SCG008-12

会場:301A

時間:5月26日 11:30-11:45

## 西南日本白亜紀 古第三紀花崗岩に関する年代学的、岩石学的研究および数値計算 Chronology, Petrology and Numerical Simulation for Cretaceous to Paleogene Granitic Rocks, SW Japan

飯田 和也<sup>1\*</sup>, 岩森光<sup>2</sup>, Park Taeho<sup>2</sup>, 折橋祐二<sup>3</sup>, Yong-Joo Jwa<sup>4</sup>, Sung-Tack Kwon<sup>5</sup>, 檀原 徹<sup>6</sup>, 岩野 英樹<sup>6</sup>  
Kazuya Iida<sup>1\*</sup>, Hikaru Iwamori<sup>2</sup>, Park Taeho<sup>2</sup>, Yuji Orihashi<sup>3</sup>, Yong-Joo Jwa<sup>4</sup>, Sung-Tack Kwon<sup>5</sup>, Tohru Danhara<sup>6</sup>, Hideki  
Iwano<sup>6</sup>

<sup>1</sup> 東京大学, <sup>2</sup> 東京工業大学, <sup>3</sup> 東京大学地震研究所, <sup>4</sup> Gyeongsang Nat'l University, <sup>5</sup> University of Yonsei, <sup>6</sup> 京都フィッション  
トラック

<sup>1</sup> Tokyo University, <sup>2</sup> Tokyo Institute of Technology, <sup>3</sup> Earthquake Research Institute, <sup>4</sup> Gyeongsang Nat'l University, <sup>5</sup> University  
of Yonsei, <sup>6</sup> Kyoto Fission-Track

Granitoid is one of the main components of the continental upper crust, and is thought to provide key information on evolution of the continental crust yet the origin of granitoid remains enigmatic. In order to address this important and unresolved problem, we have studied the origin of the Cretaceous to Paleogene granitic rocks in SW Japan. We especially focus on (1) spatial-temporal variation of granitoid, (2) temporal variation of petrological signature of granitoid, and (3) tectonic and dynamic setting that caused the observed spatial-temporal variation and provided heat for the melt generation, based on the U-Pb zircon age, whole rock chemistry, and numerical simulation for the thermal field of subduction zone. As a result, systematic spatial-temporal variations of magmatism and the spatial variations of petrological signatures in SW Japan have been observed. Numerical modeling suggests that ridge subduction model can explain spatio-temporal variation of granitic rocks in SW Japan. Based on these results, we discuss the origin of these variations.

U-Pb zircon age determinations using LA-ICPMS was performed on total 91 rock samples. The obtained age range from 95 Ma to 30 Ma, with a possible temporal gap between 60 Ma and 50 Ma. During 95-60 Ma, the systematic migration of granitoid magmatism from the south to the north occurred. Temporal variation of petrological signatures is also observed: with time, (1) initial ratio of Sr isotopes ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) decreased from enriched characters (0.7090-0.7065) to depleted ones (0.7065-0.7050), and (2) rock types of granitoid changed from ilmenite-series to magnetite-series.

In order to investigate the mechanism of temporal changes observed, ridge subduction model has been tested with numerical simulation, especially in terms of thermal impact of the subducted ridge as a function of subduction velocity and angle with or without slab window. The numerical results suggest the following points: (1) thermal impact of ridge subduction is potentially large especially when subduction velocity changes from high to low (i.e., the leading plate subducts faster), creating a slab window, (2) after the ridge subduction, melting region, mainly of slab melting, shrinks from the trench side to the rear arc side, and (3) subduction angle together with subduction velocity is important to control the overall duration and spatial range of magmatism, required to explain the observed wide horizontal range of the magmatism.

Finally, the numerical results have been compared with the observed spatio-temporal variation of the granitoid magmatism in SW Japan. The comparison suggests that the observed features (e.g., long duration (greater than 35 m.y.), a wide across-arc distance (~150 km) of the magmatism, and northward (toward backarc side) migration of the magmatism) require a shallow subduction angle (~10 degrees) with velocity change of the plates from high to low is required. With these specific conditions, it is inferred that the ridge subduction could have been the primary cause of Cretaceous to Paleogene granitoid magmatism in SW Japan.

SCG008-13

会場:301A

時間:5月26日 11:45-12:00

## Correlations of the blueschist facies metamorphism from the Heilongjiang Complex, NE China and the Suo belt, SW Japan Correlations of the blueschist facies metamorphism from the Heilongjiang Complex, NE China and the Suo belt, SW Japan

Weimin Li<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Yongjiang Liu<sup>2</sup>, Yingli Zhao<sup>2</sup>, Limin Zhao<sup>1</sup>  
Weimin Li<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Yongjiang Liu<sup>2</sup>, Yingli Zhao<sup>2</sup>, Limin Zhao<sup>1</sup>

<sup>1</sup>Shimane University, <sup>2</sup>Jilin University, P. R. China

<sup>1</sup>Shimane University, <sup>2</sup>Jilin University, P. R. China

The Jurassic accretionary terranes, which are characterized by high-P/T blueschist facies metamorphism, have been documented from the eastern margin of the Asian continent. Such high-P/T metamorphic evidence occurs from northeastern Russia via northeastern China to southwestern Japanese islands (Sengor et al., 1993; Isozaki, 1997; Taira, 2001; Wu et al., 2007).

High-P/T blueschists of the Heilongjiang Complex are exposed in northeastern China, and they consist mainly of epidote-glaucophane schists with minor amounts of garnet-barroisite schists and glaucophane aegirine-augite schists. The peak metamorphic conditions obtained for the Heilongjiang high-P/T metamorphic rocks vary from the epidote blueschist facies (320-550 C, 6-15 kbar; e.g. Bai et al., 1988; Zhou et al., 2009; Li et al., 2010a) to the epidote amphibolite facies (500-540 C, 10-12 kbar; Li et al., 2010b). U-Pb ages of 190-255 Ma for detrital zircons suggest that the protoliths of the Heilongjiang Complex are the early Mesozoic granitic batholiths (e.g. Zhou et al., 2009; Li et al., 2010c). Furthermore, the high-P/T Heilongjiang metamorphic rocks were developed by the subduction and collision of the Jiamusi Massif to the eastern border of Central Asian Orogenic Belt at the time of Jurassic period (145-190 Ma; Wu et al., 2007; Li et al., 2010c).

In the Japanese islands, Jurassic accretionary terranes occur as large-scale nappes that are tectonically sandwiched between overlying pre-Jurassic and underlying post-Jurassic sequences (e.g. Isozaki, 1997). The Suo metamorphic belt in southwestern Japan is characterized by blueschist facies metamorphism of 160-230 Ma in metamorphic age (Nishimura, 1998). The blueschists are predominantly made up of epidote-glaucophane schists with subordinate epidote-barroisite schists and sodic clinopyroxene-bearing phengite-epidote schists in the Suo belt, especially in the Gotsu and Masuda areas (e.g. Sengan, 1985). The peak metamorphic conditions preserved in the epidote-glaucophane schists are defined by the crystallization of porphyroblastic epidote, Na-amphibole (glaucophane and Mg-riebeckite), phengite (Si = 6.6-7.0 pfu), chlorite, hematite and titanites, suggesting the epidote-blueschist facies metamorphic conditions (430-530 C, 12-15.5 kbar). Additionally, the metamorphic facies series of the Suo schists of pumpellyite-actinolite facies through glaucophane schist facies to epidote amphibolite facies has been described by previous studies (e.g. Nishimura et al., 1998).

These facts suggest the similarities of both high-P/T blueschists from the Heilongjiang Complex, NE China and the Suo metamorphic belt, SW Japan, not only in the timing of metamorphism, but also in the metamorphic conditions. In this study, we therefore propose that the Heilongjiang high-P/T metamorphic rocks continued to the south, and the Suo metamorphic belt in southwest Japan is thus possible southern extension of the Heilongjiang Complex. The results reported here will contribute toward a better understanding of the Mesozoic tectono-metamorphic development of the eastern margin of the Eurasian continent.

### References:

Bai, J.W. et al., 1988, *Acta Petrologica et Mineralogica* 7, 298-308; Isozaki, Y. 1997, *The Island Arc*, 6, 25-51; Li, W.M. et al., 2010a, *Abst. 2010 Annual Meeting of the JAMS*; Li, W.M. et al., 2010b, *J. Miner. Petrol. Sci.*, 105, 86-91; Li, W.M. et al., 2010c, *Abst. 117th Annual Meet Geol. Soc. Japan*; Nishimura, Y. 1998, *J. Metamorphic Geol.*, 16, 129-140; Sengan, H. 1985, *Geol. Rep. of Shimane Univ.*, 4, 41-59; Sengor A. M. C. et al., 1993, *Nature*, 364, 299-307; Taira, A. 2001, *Annu. Rev. Earth Planet. Sci.*, 29, 109-34; Wu, F.Y. et al., 2007, *Island Arc*, 16, 156-172; Zhou, J.B. et al., 2009, *Tectonophysics*, 478, 230-246.

キーワード: high-P/T, blueschist, Heilongjiang Complex, Suo metamorphic belt, NE China, SW Japan

Keywords: high-P/T, blueschist, Heilongjiang Complex, Suo metamorphic belt, NE China, SW Japan

SCG008-14

会場:301A

時間:5月26日 12:00-12:15

## アジア大陸成長期の大陸衝突変成作用と黒瀬川帯の形成

## Multiple collision metamorphism during Asian continental growth with relation to the formation of Kurosegawa tectonic zone

小山内 康人<sup>1\*</sup>, 中野 伸彦<sup>1</sup>, 大和田正明<sup>2</sup>, サティッシュ クマール エム<sup>3</sup>, 河上哲生<sup>4</sup>, 柚原雅樹<sup>5</sup>, 足立達郎<sup>1</sup>, 米村和紘<sup>1</sup>, 吉本 紋<sup>1</sup>

Yasuhito Osanai<sup>1\*</sup>, Nobuhiko Nakano<sup>1</sup>, Owada Masaaki<sup>2</sup>, Satish-Kumar, M.<sup>3</sup>, Kawakami Tetsuo<sup>4</sup>, Yuhara Masaki<sup>5</sup>, Adachi Tatsuro<sup>1</sup>, Yonemura Kazuhiro<sup>1</sup>, Yoshimoto Aya<sup>1</sup>

<sup>1</sup>九州大学, <sup>2</sup>山口大学, <sup>3</sup>静岡大学, <sup>4</sup>京都大学, <sup>5</sup>福岡大学

<sup>1</sup>Kyushu University, <sup>2</sup>Yamaguchi University, <sup>3</sup>Shizuoka University, <sup>4</sup>Kyoto University, <sup>5</sup>Fukuoka University

In E- and SE-Asia, there are at least seven micro-continental blocks of Mongolia, North China, South China, Indochina, Shanthai, Sibumasu and West Burma from N-NE to SW. Geological Research program to realize the tectono-metamorphic processes during continental collision orogen of E- and SE-Asia has been done for the last decade.

Multiple collision boundaries in Asian continent are identified and they are subdivided into following six;

- 1) Siberian craton/Mongolian: 480 ? 500 Ma,
- 2) Mongolian/North China: 280 ? 300 Ma (so called Central Asian Orogenic Belt),
- 3) North China/South China: 240 ? 250 Ma,
- 4) South China/Indochina: 230 ? 250 Ma (Trans Vietnam Orogenic Belt),
- 5) Indochina-Shanthai/Sibumasu: ca. 210 Ma,
- 6) Sibumasu-West Burma/Indian subcontinent: ca. 45 Ma.

Characteristics of constituent rock types and metamorphic processes in each collision boundary were reported by many detailed works (e.g. Osanai et al., 2004, 2008; Nakano et al., 2008, 2009, 2010).

Especially in the collision boundary between the North and the South China blocks (e.g. north Dabie terrane, Imjingang belt), various kinds of Permo-Triassic collision zone metamorphic rocks are identified. The Higo metamorphic complex as well as the Hida-Oki terrane in Japan would also have belonged to this type of collision zone and which experienced a top-to-the-south displacement under extrusion tectonics with forming a regional nappe structure before the Cretaceous granitic activities. The basal portion of this regional nappe extrusion would be considered as a serpentinite diapir derived from relatively ductile mantle beneath the North China craton. The diapir intruded along the collision boundary and included various kinds of metamorphic rocks of HT granulite and amphibolite from the lower crustal portion of obducted North China craton and HP blueschist and eclogite from the shallower portion of subducted South China craton. The Kurosegawa tectonic zone as the serpentinite melange corresponds with this diapir.

In the presentation, we will make a detailed talk on this tectono-metamorphic evolution in eastern Asia.

SCG008-15

会場:301A

時間:5月26日 12:15-12:30

## Permian sequences and faunas in the peri-Gondwanan region and their palaeogeographical and tectonic implications Permian sequences and faunas in the peri-Gondwanan region and their palaeogeographical and tectonic implications

Shuzhong Shen<sup>1\*</sup>, Yichun Zhang<sup>1</sup>, Yue Wang<sup>1</sup>, Katsumi Ueno<sup>2</sup>  
Shuzhong Shen<sup>1\*</sup>, Yichun Zhang<sup>1</sup>, Yue Wang<sup>1</sup>, Katsumi Ueno<sup>2</sup>

<sup>1</sup>Nanjing Inst Geol Palaeont., <sup>2</sup>Fukuoka University

<sup>1</sup>Nanjing Inst Geol Palaeont., <sup>2</sup>Fukuoka University

The rifting and drifting of different tectonic blocks in the peri-Gondwanan region has long been one of the most hotly-discussed issues among geoscientists. Extensive investigations during the past decade indicate that numerous sections with carbonate and clastic deposits containing abundant Permian faunas are distributed in the Himalayan region and the Indus-Tsangbo Suture Zone in southern Tibet. A detailed comparison of faunal affinities among different tectonic blocks in different stages in the Permian Period suggests that nearly all blocks in the peri-Gondwanan region including the Qiangtang, Baoshan, Tengchong, Lasha and the Himalaya Tethys Zone are characterized by containing diamictites and typical cold-water faunas in the pre-Artinskian time, therefore were probably in a relatively high-latitude area and attached to the northern margin of Gondwanaland. By the end of late Sakmarian or early Artinskian time, warm-water faunas first occur in the Baoshan and Tengchong blocks in western Yunnan, which probably implies that those blocks began to drift away from the peri-Gondwanan margin and they moved to a relatively warm temperate zone in the Late Guadalupian (Middle Permian), as indicated by widespread distribution of warm-water faunas and carbonate deposits. The earliest warm-water faunas occurred in the Midian in the Lasha Block, which suggest a slightly later rifting and/or climatic amelioration than the Tengchong and Baoshan blocks. Numerous exotic blocks between the Lhasa Block and the Himalaya Tethys Zone containing abundant Middle Permian fusuline faunas and compound rugose corals also indicate a different palaeontological contents and palaeobiogeographical affinities between the Lhasa Block in the north and the Himalaya Tethys Zone in the south. By the Late Permian (Lopingian) time, these palaeontological and palaeobiogeographical disparities became more evident, therefore strongly suggest that the Lhasa Block probably rifted away from the peri-Gondwanan region in the early Middle Permian, a much earlier open time of the Neotethys, not Triassic as previously suggested.

キーワード: Permian, tectonic blocks, peri-Gondwanan region, faunas, rifting, drifting  
Keywords: Permian, tectonic blocks, peri-Gondwanan region, faunas, rifting, drifting



SCG008-16

会場:301A

時間:5月26日 12:30-12:45

## 東アジア地域の地震テクトニクス、火山の起源とマントルダイナミクス East Asia: Seismotectonics, Volcanism and Mantle Dynamics

趙 大鵬<sup>1\*</sup>

Dapeng Zhao<sup>1\*</sup>

<sup>1</sup> 東北大学・理

<sup>1</sup>Geophysics, Tohoku University

We introduce the significant recent results of geophysical studies and discuss their implications on seismotectonics, magmatism, and mantle dynamics in East Asia. High-resolution geophysical imaging revealed structural heterogeneities in the source areas of large crustal earthquakes, which may reflect magma and fluids that affected the rupture nucleation of large earthquakes. In subduction zone regions, the crustal fluids originate from the dehydration of the subducting slab. Magmatism in arc and back-arc areas is caused by the corner flow in the mantle wedge and dehydration of the subducting slab. The intraplate magmatism has different origins. The continental volcanoes in Northeast Asia (such as Changbai and Wudalianchi) seem to be caused by the corner flow in the big mantle wedge (BMW) above the stagnant slab in the mantle transition zone and the deep dehydration of the stagnant slab as well. The Tengchong volcano in Southwest China is possibly caused by a similar process in BMW above the subducting Burma microplate (or Indian plate). The Hainan volcano in southernmost China seems to be a hotspot fed by a lower-mantle plume associated with the Pacific and Philippine Sea slabs' deep subduction in the east and the Indian slab's deep subduction in the west down to the lower mantle. The occurrence of deep earthquakes under the Japan Sea and the East Asia margin may be related to a metastable olivine wedge in the subducting Pacific slab. The stagnant slab finally collapses down to the bottom of the mantle, which may trigger upwelling of hot mantle materials from the lower mantle to the shallow mantle beneath the subducting slabs and cause the slab-plume interactions. Some of these issues, such as the origin of intraplate magmatism, are still controversial, and so further detailed studies are needed from now.

キーワード: 東アジア, 地震テクトニクス, プレート内部火山, マントルダイナミクス, 沈み込むスラブ

Keywords: East Asia, seismotectonics, intraplate volcanism, mantle dynamics, subducting slab

SCG008-P01

会場:コンベンションホール

時間:5月26日 14:00-16:30

## Thermal isostasy below the arabian shield and platform: implications caused by red sea spreading Thermal isostasy below the arabian shield and platform: implications caused by red sea spreading

Mogren Saad<sup>1\*</sup>, Manoj Mukhopadhyay<sup>1</sup>, Abdulaziz Al-jasser<sup>1</sup>  
Saad Mogren<sup>1\*</sup>, Manoj Mukhopadhyay<sup>1</sup>, Abdulaziz Al-jasser<sup>1</sup>

<sup>1</sup>King Saud University

<sup>1</sup>King Saud University

Thermal isostatic compensation for the Arabian Shield and Platform is investigated here by an analysis of gravity field, deep seismic refraction profile and heat-flow data. Regression relationships between elevation and gravity anomalies found for the geologic provinces of Arabian Shield and Platform are examined first using both point-values as well as the averaged anomalies over 1 degree and 1 degree areas. These suggest for grossly different isostatic status prevailing for the geologic provinces in response to crustal dynamics. The composite terrain-corrected Bouguer anomaly and Airy-Heiskanen Isostatic anomaly maps for Arabian Shield and Platform are prepared using these data sets. The gravitational attraction of the Moho was determined from the 30 arc-seconds topography data; calculation of isostatic correction was made assuming 2.67 g/cm<sup>3</sup> and 3.3 g/cm<sup>3</sup> densities for the crust and top mantle respectively. A mean thickness of 40 km is used for isostatic data reduction. Mass/unit area under the crustal columns at the assumed level of compensation at 40 km depth for Arabian Shield and Platform is constructed and compared with those for other shield regions in the world. Thermal isostatic compensating model for ASP is constructed using the available deep crustal seismic refraction results and heat-flow data for the Arabian Shield. For modeling, the digitized elevation data are examined in details to look for geological mass anomalies in upper crust. Once these effects are isolated, the long wavelength isostatic anomalies are ascribed to crust-mantle configuration below the Arabian Shield and Platform. The results suggest that the elevated Asir Igneous Province (that initially formed as a magmatic arc but later accreted to the Arabian Shield) is largely uncompensated, whereas, the isostatic compensation for the Arabian Shield and Platform is more complete. Tectonic implications for the moving Arabian lithosphere due to Red Sea spreading for the last 26 M.Y. in influencing the thermal isostatic compensation status is discussed.

キーワード: thermal isostasy, arabian shield, red sea, gravity, Saudi Arabia, isostatic anomalies  
Keywords: thermal isostasy, arabian shield, red sea, gravity, Saudi Arabia, isostatic anomalies

## タイ王国中央部に分布する放射虫層状チャート：地質年代と対比 Radiolaria-bearing bedded chert in the Central Plain of Thailand: its geologic age and correlation.

鎌田 祥仁<sup>1\*</sup>, 加藤京都<sup>1</sup>, 上野勝美<sup>2</sup>, 宮東 照<sup>2</sup>, Thasinee Charoentitirat<sup>3</sup>, Seranee Limruk<sup>3</sup>  
Yoshihito Kamata<sup>1\*</sup>, Miyako Kato<sup>1</sup>, Katsumi Ueno<sup>2</sup>, Akira Miyahigashi<sup>2</sup>, Thasinee Charoentitirat<sup>3</sup>, Seranee Limruk<sup>3</sup>

<sup>1</sup> 山口大学, <sup>2</sup> 福岡大学, <sup>3</sup> チュラロンコン大学

<sup>1</sup> Yamaguchi Univ., <sup>2</sup> Fukuoka Univ., <sup>3</sup> Chulalongkorn Univ.

Geotectonic division of Thailand has been established based on fundamental differences of regional geological and stratigraphical features, including the origin of granitoids series, distribution of ultra-mafic rocks which represent position of tectonic suture lines, and fusuline and radiolarian paleontological paleontological data. In the Central Plain of Thailand, however, the tectonic division is vague due to the scarceness of Paleozoic and Mesozoic information by the covering of thick Quaternary sediments. Recently, occurrence of Permian radiolarians from siliceous rocks in the Nakhon Sawan and Uthai Thani areas in the Central Plain has been reported (Saesaengseerung et al., 2007). However, tectonic implication of Paleozoic and Mesozoic rocks distributed in the Central Plain, in other word, extension to the Central Plain of the tectonic divisions such as the Sibumas Block, Inthanon Zone, and Sukhothai Zone which are well established in Northern Thailand, has not been well known.

Recently, we made field survey in Sukhothai, Nakhon Sawan, Uthai Thani, and Kanchanaburi provinces in the Central Plain. Chert beds distributed in the Sukhothai, Nakhon Sawan and Uthai Thani areas are gray or red in color. It is generally recrystallized by contact metamorphism. Under the microscope, the chert is composed mainly of a microcrystalline quartz matrix with radiolarian tests and quartz veins. Poorly preserved radiolarians have been detected from those exposed in eight localities of these areas. Sakmarian (middle Cisuralian) to Capitanian (late Guadalupian) radiolarians such as *Albaillella asymmetrica*, *Pseudoal-baillella fusiformis*, *Ps. globosa*, *Follicucullus scholasticus* are obtained.

At Bo Phloi in the Kanchanaburi area, we examined green to greenish gray chert. Under the microscope, radiolarian tests are observed to be embedded within a clay-rich cryptocrystalline quartz matrix. Calcareous foraminiferal tests have often been obtained from the chert by acidic treatment. Moreover, a lot of thin-shelled bivalves, probably *Daonella* or *Halobia*, are found in several horizons of the chert. It also yields well-preserved Middle Triassic (Anisian) radiolarians such as *Eptingium manfredi*, *Pseudostylosphaera japonica*, *Triassocampe coronata*, and others.

Based on the radiolarian occurrence and lithology, the Permian chert in the Nakhon Sawan and Uthai Thani areas are correlated with the Khanu Chert distributed in the Sukhothai area of the Sukhothai Zone in Northern Thailand.. On the other hand, the Triassic chert exposed at Bo Phloi in the Kanchanaburi area clearly exhibits hemipelagic deposition on the eastern margin of the Sibumasu Block, based on its lithology, radiolarian age, and faunal content characterized by possessing calcareous tests of thin-shelled bivalves and foraminifers.

キーワード: タイ王国中央部, シブマス地塊, スコタイ帯, 放射虫, ペルム系, 三畳系

Keywords: Central Thailand, Sibumasu Block, Sukhothai Zone, radiolarians, Permian, Triassic

SCG008-P03

会場:コンベンションホール

時間:5月26日 14:00-16:30

## タイ中部 Hua Hin 地域に分布する花崗岩質岩の Rb-Sr 年代 Rb-Sr ages of granitic rocks from the Hua Hin area, Thailand

柚原 雅樹<sup>1\*</sup>, 亀井淳志<sup>2</sup>, 吉本 紋<sup>3</sup>, 河上哲生<sup>4</sup>, 上久保 寛<sup>5</sup>, 中野伸彦<sup>3</sup>, 小山内康人<sup>3</sup>, Punya Charusiri<sup>6</sup>  
Masaki Yuhara<sup>1\*</sup>, Atsushi Kamei<sup>2</sup>, Aya Yoshimoto<sup>3</sup>, Tetsuo Kawakami<sup>4</sup>, Hiroshi Kamikubo<sup>5</sup>, Nobuhiko Nakano<sup>3</sup>, Yasuhito Osanai<sup>3</sup>, Punya Charusiri<sup>6</sup>

<sup>1</sup> 福岡大学, <sup>2</sup> 島根大学, <sup>3</sup> 九州大学, <sup>4</sup> 京都大学, <sup>5</sup> JOGMEC, <sup>6</sup> Chulalongkorn 大学

<sup>1</sup> Fukuoka University, <sup>2</sup> Shimane University, <sup>3</sup> Kyushu University, <sup>4</sup> Kyoto University, <sup>5</sup> JOGMEC, <sup>6</sup> Chulalongkorn University

The granitic rocks are widely distributed in the Hua Hin area, Thailand. This area is located in the Main-Rang Province (Cobbing et al., 1986) consists mainly of S-type granitic rocks, whose ages range from early Late Triassic to late Early Jurassic (ca. 230-180 Ma) (Sone and Metcalfe, 2008). The petrogenesis of these granitic rocks is explained by partial melting of the Sibumasu crust subducted beneath the Palaeo-Tethys accretionary complex (Sone and Metcalfe, 2008). However, the timing of magma activity and characteristic of source material of granitic rocks in the Hua Hin area are poorly understood. In this paper, we address Rb-Sr isotopic study of granitic rocks from this area.

The granitic rocks in the Hua Hin area are composed of foliated granitic rocks and non-foliated granitic rocks. The formers are the Hub Kapong Gneissic Granite, Hua Hin Gneissic Granite and Pran Buri Gneissic Granite. The Hub Kapong and Hua Hin Gneissic Granites are partly weakly mylonitized K-feldspar porphyritic biotite granite. Some of the Hub Kapong Gneissic Granite includes tourmaline. Parts of Hua Hin Gneissic Granite include garnet and tourmaline. Sillimanite is also contained in some of them (Kawakami et al., 2010). The Pran Buri Gneissic Granite is mylonitic biotite granite. Non-foliated granitic rocks are stock bodies intruded into the Hub Kapong Gneissic Granite, and composed of biotite to two-mica granite. These granitic rocks have peraluminous chemical composition. The Hub Kapong and Hua Hin Gneissic Granites and non-foliated granitic rocks indicate low magnetic susceptibility ( $SI=1.3-21.5 \times 10^{-5}$ ), but the Pran Buri Gneissic Granite indicates high value ( $SI=185-455 \times 10^{-5}$ ) (Yoshimoto et al., 2010).

The Hub Kapong and Hua Hin Gneissic Granites give Rb-Sr whole-rock isochron ages of  $202 \pm 22$  Ma ( $SrI=0.7259 \pm 0.0024; N=13$ ) and  $209 \pm 14$  Ma ( $SrI=0.7258 \pm 0.0034; N=10$ ), respectively. Because of dispersion, errors of age and SrI are slightly big. The dispersion might be due to effects of deformation and metamorphism. The Pran Buri Gneissic Granite does not give significant isochron age. A body of non-foliated granitic rocks gives a Rb-Sr whole-rock isochron ages of  $84 \pm 13$  Ma ( $SrI=0.7356 \pm 0.0015; N=4$ ). These ages indicate timing of magmatic activity in the Hua Hin area, and accord with U-Th-Pb monazite ages (213 Ma (core), 83 Ma (rim)) of metamorphic rocks from this area (Nakano et al., 2010).

Keywords: Rb-Sr whole-rock isochron age, granitic rocks, Hua Hin, Thailand

SCG008-P04

会場:コンベンションホール

時間:5月26日 14:00-16:30

## 地球化学的にみたパレオテチス沈み込み帯における砂岩の後背地：タイ北部インタノン帯の例

### Geochemistry of sandstone related to Paleo-Tethys subduction zone in the Inthanon Zone, northern Thailand

國井 美幸<sup>1\*</sup>, 原 英俊<sup>2</sup>, 足立 佳子<sup>3</sup>, 久田 健一郎<sup>1</sup>, チャルシリ パンニャ<sup>4</sup>, Wiwegwin Weerachat<sup>1</sup>

Miyuki Kunii<sup>1\*</sup>, Hidetoshi Hara<sup>2</sup>, Yoshiko Adachi<sup>3</sup>, Ken-ichiro Hisada<sup>1</sup>, Punya Charusiri<sup>4</sup>, Weerachat Wiwegwin<sup>1</sup>

<sup>1</sup> 筑波大学生命環境科学研究科, <sup>2</sup> 産業技術総合研究所地質情報部門, <sup>3</sup> 新潟大学超域研究機構, <sup>4</sup> チュラロンコン大学  
<sup>1</sup>University of Tsukuba, <sup>2</sup>Geological survey of Japan, <sup>3</sup>Niigata University, <sup>4</sup>Chulalongkorn University

パレオテチスは、デボン紀に、ゴンドワナ大陸からインドチャイナ地塊を含むカタイシア地塊群がリフティングしたことにより形成された。そして、三畳紀にシンメリアン地塊群の一部であるシブマス地塊と、インドチャイナ地塊の、2つの微小大陸が衝突したことにより閉塞した (Metcalf, 1999)。タイ北部のインタノン帯は、パレオテチスの収束域とされており、その形成やテクトニック・セッティングは、東南アジア半島部の形成史にとって重要な意味を持つ。

上野・久田 (1999) によれば、インタノン帯には、デボン系から三畳系の放散虫リボンチャートや石炭系からベルム系の海山起源炭酸塩岩類が大量に分布しており、また、シブマス地塊の構成要素とみなすことのできる、オルドビス系石灰岩や変成岩及び堆積岩が分布している。そのため、インタノン帯は、構造的にシブマス地塊を基盤に持ち、その上に、パレオテチスの要素が、巨大なスラストシートとして定置しているとした。さらに、Hara et al. (2009) では、インタノン帯から、メランジュを記載し、パレオテチス沈み込みに伴う付加体の形成を報告した。そこで、本研究では、インタノン帯の基盤であるシブマス地塊に相当する石炭系砂岩と、メランジュ中の砂岩について、その後背地を検討するため、岩石学的・地球化学的な分析を行った。

シブマス地塊に相当する石炭系砂岩と、メランジュ中の砂岩について、Gazzi-Dickinson 法に基づき、モード組成の測定を行った。Dickinson ダイアグラム (Dickinson et al., 1983) によれば、シブマス地塊の砂岩は、well-rounded の石英粒子を含む石英質砂岩であり、大陸性砂岩であるといえる。また、ダイアグラム上では、安定地塊内部から石英質再循環にプロットされた。一方、メランジュ中の砂岩は、火山岩片を多く含む石質砂岩であり、ダイアグラム上では、石英質から石質再循環にプロットされた。

インタノン帯の両砂岩を、主要元素および微量元素・希土類元素について地球化学的に分析した。主要元素については、産総研の XRF (PANalytical Axios PW4400/40) を、微量元素・希土類元素については、新潟大学理学部の XRF (Rigaku RIX3000) および ICP-MS (Agilent 7500a) を使用し、分析を行った。砂岩の全岩分析結果を、主要元素は Bhatia (1983) の、微量元素・希土類元素は Bhatia and Crook (1986) のグラフ上にプロットした。主要元素の分析結果について、 $Fe_2O_3 + MgO \% - TiO_2 \%$ ,  $Fe_2O_3 + MgO \% - Al_2O_3 / SiO_2$ ,  $Fe_2O_3 + MgO \% - K_2O / Na_2O$ ,  $Fe_2O_3 + MgO \% - Al_2O_3 / (CaO + Na_2O)$  から、シブマス地塊の石英質砂岩は、Passive Margins 中またはその周辺にプロットされる傾向を示した。また、微量元素・希土類元素の分析結果について、La - Th - Sc, Th - Sc - Zr, La / Sc - Ti / Zr, Sc / Cr - La / Y から、シブマス地塊の石英質砂岩は Passive Margins, メランジュ中の砂岩は Continental Island Arc 中またはその周辺にプロットされた。

以上の結果より、シブマス地塊の石英質砂岩は大陸要素が強く、島弧の影響を受けていないと考えられる。一方、メランジュ中の砂岩は、島弧の影響が強く、その島弧から砕屑物が供給されていたと考えられ、シブマス地塊の石英質砂岩とメランジュ中の砂岩は、全く異なる後背地の特徴を示すといえる。すなわち、後期ベルム紀から三畳紀にかけて、インドチャイナ地塊西縁から南縁で島弧 (スコタイ帯) が発達した。メランジュ中の砂岩は、この島弧を起源とすると考えられる。また、パレオテチスの沈み込みによる付加体は、この島弧からの砕屑物の供給によって発達したと考えられる。

キーワード: パレオテチス, インタノン帯, 砂岩, 後背地, 地球化学

Keywords: Paleo-Tethys, Inthanon Zone, sandstone, provenance, geochemistry

SCG008-P05

会場:コンベンションホール

時間:5月26日 14:00-16:30

## タイ王国 Pranburi-Hub Kapong 地域に産する変花崗岩中のジルコンの産状 Mode of occurrence of zircon in metagranites from the Pranburi-Hub Kapong area, Thailand

河上 哲生<sup>1\*</sup>, 柚原 雅樹<sup>2</sup>, 横山 隆臣<sup>1</sup>, 牧 賢志<sup>1</sup>, 小山内 康人<sup>3</sup>, 平田 岳史<sup>1</sup>, 上久保 寛<sup>4</sup>, チャルシリ パンチャ<sup>5</sup>  
Tetsuo Kawakami<sup>1\*</sup>, Masaki Yuhara<sup>2</sup>, Yokoyama D. Takaomi<sup>1</sup>, Kenshi Maki<sup>1</sup>, Yasuhito Osanai<sup>3</sup>, Takafumi Hirata<sup>1</sup>, Hiroshi Kamikubo<sup>4</sup>, Punya Charusiri<sup>5</sup>

<sup>1</sup> 京都大学大学院理学研究科, <sup>2</sup> 福岡大学理学部, <sup>3</sup> 九州大学大学院比較社会文化研究院, <sup>4</sup> 石油天然ガス・金属鉱物資源機構, <sup>5</sup> チュラロンコーン大学

<sup>1</sup>Kyoto University, <sup>2</sup>Fukuoka University, <sup>3</sup>Kyushu University, <sup>4</sup>JOGMEC, <sup>5</sup>Chulalongkorn University

The metagranites and pelitic-psammitic metamorphic rocks are widely exposed from Pranburi to Hub Kapong, central Thailand. The EPMA dating of monazite (so-called CHIME monazite dating, Suzuki et al., 1994) in a high-grade pelitic metamorphic rock from the south of the study area revealed that the monazite core is 213 Ma and the rim is 83 Ma (Nakano et al., 2010). Yuhara et al. (2011) dated the metagranites by the Rb-Sr whole-rock isochron method. The Hub Kapong gneissic granite that crops out in the north gave 202±22 Ma, and the Hua Hin gneissic granite at the central area gave 209±14 Ma. The non-foliated granitic rocks in the central area gave 84±13Ma. The large error is attributed to the later metamorphism and deformation that postdated granite crystallization. This study aims to constrain the timing of granite crystallization and metamorphism through textural observation and U-Pb dating of zircon by LA-ICPMS.

The foliation of the Hua Hin gneissic granite is defined by the arrangement of Bt +/- Si, and this body is metamorphosed. Zircon in this metagranite has euhedral oscillatory zoning and likely magmatic in origin. Under the cathode luminescence (CL) image, detrital cores and dark overgrowths on them are commonly recognized.

The Hub Kapong gneissic granite is a two-mica granite and muscovite can be metamorphic in origin (Kawakami et al., 2010). This metagranite characteristically contains xenotime-zircon intergrowth of magmatic origin (e.g., Viskupic et al., 2005). Xenotime shows mosaic-like zoning and includes a lot of bright Th-bearing phases under the back scattered electron (BSE) image. It also has many tiny holes and shows spongy texture. Zircon shows idiomorphic oscillatory zoning under BSE image, while it is dark and shows mosaic-like texture under CL image. Rarely, zircon as well includes bright Th-bearing phases under BSE image, and shows spongy texture. This kind of spongy zircon is considered to form by a dissolution-precipitation process in a fluid (Geisler et al., 2007). Therefore, zircon in the Hub Kapong gneissic granite would be originally magmatic, but some of them may have experienced the dissolution-precipitation process. Yuhara et al. (2011) interpreted that 83 Ma age preserved in the monazite from the pelitic metamorphic rock is likely due to intrusion of the non-foliated granitic rocks. Since the non-foliated granitic rocks are scattered around the Hub Kapong gneissic granite, it is probable that the intrusion of the non-foliated granitic rocks are responsible for the formation of the spongy zircon in the Hub Kapong gneissic granite. These possibilities will be tested through the zircon U-Pb dating by the LA-ICPMS.

キーワード: 花崗岩, タイ, ジルコン, 変成作用, 流体  
Keywords: granite, Thailand, zircon, metamorphism, fluid

SCG008-P06

会場:コンベンションホール

時間:5月26日 14:00-16:30

## 中国南西部およびタイ西部のシャンタイーシブマス地塊境界に分布する苦鉄質変成岩類の地球化学的特徴

### Geochemical character of metamorphosed mafic rocks from the collision boundary between Shan-Thai and Sibumasu craton in

米村 和紘<sup>1\*</sup>, 小山内 康人<sup>1</sup>, 中野 伸彦<sup>1</sup>, 大和田 正明<sup>2</sup>, 馬場 壮太郎<sup>3</sup>, チャルシリ パンニャ<sup>4</sup>, 加々島 慎一<sup>5</sup>, 野原 (今中) 里華子<sup>6</sup>

Kazuhiro Yonemura<sup>1\*</sup>, Yasuhito Osanai<sup>1</sup>, Nobuhiko Nakano<sup>1</sup>, Masaaki Owada<sup>2</sup>, Sotaro Baba<sup>3</sup>, Punya Charusiri<sup>4</sup>, Shin-ichi Kagashima<sup>5</sup>, Nohara (Imanaka) Rikako<sup>6</sup>

<sup>1</sup>九州大学比較社会文化, <sup>2</sup>山口大学理工学研究科, <sup>3</sup>琉球大学教育学部, <sup>4</sup>チュラロンコーン大学, <sup>5</sup>山形大学理学部, <sup>6</sup>新潟大学自然科学研究科

<sup>1</sup>Kyushu Univ., <sup>2</sup>Yamaguchi Univ., <sup>3</sup>Univ. Ryukyus, <sup>4</sup>Chulalongkorn Univ., <sup>5</sup>Yamagata Univ., <sup>6</sup>Niigata Univ.

The collision boundary between Shan-Thai craton and Sibumasu craton is distributed from the Nujiang area in Yunnan province of China to the Khanom area in western Thailand, through Myanmar and The Inthanon area of Thailand (Osanai et al., 2010). Especially in the Inthanon area, the metamorphic ages were determined as 210 Ma (Nakano et al., 2010). This boundary is important for understanding the tectonic evolution of multiple collision orogeny during Permian-Triassic Asian continent growth. This study focus to metamorphosed mafic rocks that consider to have distributed between of those cratons before collision.

The metamorphosed mafic rocks from the collision boundary between the Shan-Thai and The Sibumasu cratons appear as blocks or Layers in granitic gneiss and pelitic gneiss. In the Nujiang area, the metamorphic rocks indicate greenschist? to amphibolite?facies conditions. Variations of the metamorphosed mafic rocks are clinopyroxene amphibolite, amphibolite and epidote?hornblende schist. The metamorphic rocks from the Inthanon area and the Khanom area suggest amphibolite?facies conditions. Major mineral assemblages of the metamorphosed mafic rocks from the Inthanon area are garnet + hornblende, clinopyroxene + hornblende and hornblende. Hornblendes are usually green?brownish color and show graonblastic texture. On the other hand, in the Khanom area, mineral assemblages of metamorphosed mafic rocks are clinopyroxene + hornblende, hornblende and epidote + hornblende.

The bulk chemical compositions of amphibolite and mafic schist from the Nujiang area show SiO<sub>2</sub> = 46.2-51.7 wt. %, XMg = 0.45-0.72 and various Nb/Y ratios ranging from 0.09 to 0.82. They are plotted within the fields between with-in plate basalts and MORB in some discrimination diagrams. Furthermore, the chondrite normalized REE patterns show two patterns; LREE-enriched pattern (OIB type) and flat pattern (E- to T-MORB type). These geochemical characters and mode of occurrence indicate that they would be derived from oceanic crust, which are distributed between the Shan-Thai craton and the Sibumasu craton before collision.

Keywords: Inthanon, Khanom, Nujiang, Metamorphosed mafic rock, Geochemistry, Collision boundary

SCG008-P07

会場:コンベンションホール

時間:5月26日 14:00-16:30

## 北部ベトナム・レッドリバー剪断帯のペルム紀-トリアス紀の高度変成作用 Permo-Triassic high-grade metamorphism recorded in metamorphosed bauxites from the Red River shear zone, Vietnam

中野 伸彦<sup>1\*</sup>, 小山内 康人<sup>1</sup>  
Nobuhiko Nakano<sup>1\*</sup>, Yasuhito Osanai<sup>1</sup>

<sup>1</sup>九州大学  
<sup>1</sup>Kyushu Univ.

The Red River shear zone is left-lateral shear zone caused by collision of the Indian subcontinent to the Eurasian continent at the Tertiary. Although most metamorphic rocks were reset by the deformation, recent chronological works have revealed presence of the former Permo-Triassic thermal event. However, the details of such thermal event are obscure. Here, we report metamorphic evolution and the age of aluminous granulite from the shear zone.

The whole rock chemistry of the granulites is extremely rich in  $\text{Al}_2\text{O}_3$  (36.7-50.9 wt%),  $\text{TiO}_2$  (5.5-13.0 wt%), and  $\text{Fe}_2\text{O}_3^{\text{total}}$  (24.9-35.3 wt%) and poor in  $\text{SiO}_2$  (7.9-24.1 wt%),  $\text{MgO}$  (0.5-1.0 wt%), Na (75-258 ppm analyzed by ICP-MS), and K (mostly lower than 100 ppm analyzed by ICP-MS). HFSEs and REEs also show high concentrations (e.g., Ga, 37-74 ppm; Y, 41-102 ppm; Zr, 367-724 ppm; Nb, 44-99 ppm; La, 43-187 ppm; Ce, 90-357 ppm; Pr, 10-38 ppm; Nd, 45-172 ppm; Sm, 10-37 ppm) whereas LILEs are depleted (Rb, 0.3-4.6 ppm; Sr, mostly <30 ppm; Ba, mostly <15 ppm). These features strongly suggest that the protolith of the aluminous granulites is lateritic bauxite.

The granulites commonly contain garnet, corundum, sillimanite, hercynite, and ilmenite with minor monazite, apatite, and zircon. Garnet includes kyanite, staurolite, ilmenite, siderite, and rutile. Kyanite, staurolite, siderite, and rutile are not observed in the matrix and they appear only as inclusion phases. Ilmenite occurring in the matrix contains sillimanite inclusions. Based on the petrographical observations and isochemical phase diagram using whole rock chemistry suggests former eclogite-facies (>1.8 GPa at 790 C) metamorphism and subsequent decompression under granulite-facies condition (>1000 C at 1.5 GPa).

In-situ U-Pb zircon dating was carried out using LA-ICP-MS system, Kyushu University. Based on the CL image, zircon is commonly zoned and sometimes contains dark core portion. Preliminary result is that there are several  $^{206}\text{Pb}/^{238}\text{U}$  ages (including discordant ages) from the Permian to Tertiary but the Tertiary ages were obtained only from zircon occurring in the matrix. Permian-Triassic ages (265-230 Ma) were mostly detected from the dark core and some are from zircons included in garnet. Only the dark core includes large number of high-density (Ave.  $1.15 \text{ g/cm}^3$ )  $\text{CO}_2$  fluid inclusions. Garnet, corundum and staurolite also contain  $\text{CO}_2$  fluid inclusions and their densities are also high (Ave.  $1.00 \text{ g/cm}^3$ ,  $1.07 \text{ g/cm}^3$  and  $1.09 \text{ g/cm}^3$ , respectively). These features strongly suggest that the dark cores had been in equilibrium with the garnet, corundum and staurolite that had formed under high-grade (eclogite- to granulite-facies) metamorphic conditions mentioned above.

Although further chronological works are now in progress, present study will suggest that the Permo-Triassic thermal event in the Red River shear zone is high-grade metamorphic event and it should be related with collision between the South China and Indochina cratons.

キーワード: 変成ボーキサイト, 変成作用, ペルム紀-トリアス紀, レッドリバー剪断帯, ベトナム  
Keywords: metamorphosed bauxite, metamorphic evolution, Permo-Triassic, Red River shear zone, Vietnam



SCG008-P08

会場:コンベンションホール

時間:5月26日 14:00-16:30

## A Preliminary View of Metamorphic Rocks in Indonesia Region A Preliminary View of Metamorphic Rocks in Indonesia Region

セティアワン ヌグロホ<sup>1\*</sup>, 小山内 康人<sup>1</sup>, Joko Wahyudiono<sup>2</sup>  
Nugroho Setiawan<sup>1\*</sup>, Yasuhito Osanai<sup>1</sup>, Joko Wahyudiono<sup>2</sup>

<sup>1</sup>九州大学大学院比較社会文化, <sup>2</sup>インドネシア地質調査所  
<sup>1</sup>Kyushu University, <sup>2</sup>GRDC Indonesia

Indonesian archipelago formed by reassembly of fragment rifted from Gondwana that arrived at the Eurasian subduction margin. A Cretaceous active margin is interpreted along Sumatra into western Java continued through southeastern Kalimantan into western Sulawesi, as suggested by the distribution of high pressure-low temperature subduction-related metamorphic rocks. Present-day, geology of Indonesia is the result of Cenozoic subduction and collision and situated at the boundaries of Eurasia, Indo-Australia, and Pacific-Philippine Sea plate.

Previous study by Hall (2002) suggests that in western Sumatra, Paleozoic sediments and volcanic rocks interpreted part of Indochina-East Malaya block that separated from Gondwana. In eastern Sumatra, there are Carboniferous sediments belong to Sibumasu block separated from Gondwana. Permian and Triassic granites of Thai-Malay tin belt extend into western Indonesia are the products of subduction and post-collisional magmatism in Triassic. In the southwestern Kalimantan, the Paleozoic age is represented by Carboniferous to Permian metamorphic rocks. Cretaceous granitoid plutons intrude the metamorphic rocks in Schwaner Mountains. Northwestern Kalimantan may mark as a subduction margin continuing south from East Asia.

On the east side of Indonesia, geology of Sulawesi caused by convergence between Australian, Pacific, and Eurasia plate. In eastern Sulawesi, collision resulted in thrusting of Australian continental rocks. The Banda Arc including Timor-Tanimbar-Seram islands, situated between Indo-Australian, Eurasian and Caroline plate. From the inner to outer side of the following belts, have been distinguished: ophiolite, metamorphic, and thrust-fold belt. The geology of Papua is involving interaction between Australian and Pacific plate. Lithotectonic of Papua can be divided from north to south are collided arc, ophiolite and metamorphic belt, fold-thrust belt, and foreland basin.

Purpose of this study is to understanding in detail Indonesian tectonic regime from the study of metamorphic rocks. In this preliminary view, 16 samples taken by GRDC (Geological Research and Development Centre) Indonesia have been observed petrographically and five samples from those have been analyzed by EPMA.

Seven samples from West Papua including Kaimana, Ransiki, Maar and Enarotali area are slate with the composition of quartz, muscovite, and chlorite. On the Steenkool area are biotite gneiss and white schist suggest greenschist to amphibolite facies. Biotite gneiss has composition of biotite, quartz, muscovite, plagioclase and k-feldspar. White schist has composition of quartz, muscovite and chlorite. Four samples from Kendari - southeast Sulawesi are mica schist with composition of quartz, muscovite, chloritoid and chlorite suggest greenschist facies. One sample amphibolite from Central Java shows composition of hornblende (XMg 0.62-0.70), garnet (almandine), sphene, epidote, quartz and minor minerals of plagioclase and k-feldspar.

Four samples from Schwaner Mountains, West Kalimantan, have been observed in petrography and EPMA analyses. The samples can be divided into three groups by name, which are metatonalite, metagranitoid and biotite schist. Mineral assemblages for metatonalite are quartz, plagioclase (An36-50), k-feldspar, hornblende (XMg 0.53-0.64), biotite, sphene, apatite and clinopyroxene. While on metagranitoid is shows assemblages of quartz, plagioclase (An22-23), k-feldspar, biotite, sphene, epidote and apatite. Biotite schist mineral assemblages show quartz, plagioclase (An83-85), k-feldspar, biotite, epidote and muscovite.

Detailed field observation and new collecting samples with EPMA, bulk chemistry analyses and age dating determinations are needed to understanding in details geological evidence constrain from metamorphic petrology on the Indonesian region. These steps will be held during March 2011 on the South Sulawesi and Central Java and the result will be presented on the conference.  
Keywords: metamorphic rock, Indonesia, tectonic, South Sulawesi, Central Java, Schwaner Mountains

SCG008-P09

会場:コンベンションホール

時間:5月26日 14:00-16:30

## Blueschists along the suture zone between the Ergun block and the Xing'an block in Inner Mongolia, northeastern China Blueschists along the suture zone between the Ergun block and the Xing'an block in Inner Mongolia, northeastern China

Limin Zhao<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Yongjiang Liu<sup>2</sup>, Weimin Li<sup>1</sup>  
Limin Zhao<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Yongjiang Liu<sup>2</sup>, Weimin Li<sup>1</sup>

<sup>1</sup>Shimane University, <sup>2</sup>Jilin University, P.R.China

<sup>1</sup>Shimane University, <sup>2</sup>Jilin University, P.R.China

Blueschists and related metamorphic rocks are exposed in the Toudaoqiao area of the Honghuaerji-Yimin district in the Inner Mongolia, northeastern China. These HP metamorphic rocks occur along the Tayuan-Xiguitu fault, separating the Ergun block and the Xing'an block. The HP metamorphic area belongs to the eastern segment of the Xingmeng Orogenic Belt (Li and Oyang, 1998), and it is located in the eastern parts of the Central Asian Orogenic Belt (Sengor et al., 1993). The blueschists and the related metamorphic rocks have been found in 1980's, and preliminary geological study was performed (Ye et al., 1994). According to them, there are phyllites, glaucophane-albite-chlorite-epidote schists, chlorite-quartz schists, and albite-chlorite-glaucophane schists. The mineral assemblages suggest typical LT-HP conditions.

We have had an opportunity to have a field survey in the Toudaoqiao area, and collected blueschists and other metamorphic rocks. The blueschists consist mainly of chlorite, epidote, glaucophane, with subordinate amounts of white mica, quartz, albite, hematite, apatite and titanite. A schistosity is well defined by orientation of glaucophane and chlorite. Glaucophane is of euhedral to subhedral prismatic crystal with size up to 0.3mm long. It is partly replaced by chlorite along the rim and the cleavage. The chlorite is euhedral to subhedral, and its size is up to 0.3mm across. Epidote is of anhedral rounded grain with size up to 0.1mm across. White mica is of anhedral platy crystal, and its size up to 0.2mm across. Texture and the mineral assemblage of the blueschists suggest the metamorphic conditions of typical epidote-blueschist facies.

The HP metamorphic rocks in the Toudaoqiao area is located along the Tayuan-Xiguitu fault, that is supposed to be the boundary between the Ergun block and the Xing'an block (Ge et al., 2005; Sui et al., 2006). In the eastern side of the fault, there occur significant Xinlin ophiolite (Li, 1991) and the Tahe granite (Ge et al., 2005), and these indicate a suture zone. The collision is supposed to take place in the Early Paleozoic time (Ge et al., 2005). The precise study of the other parts of the fault have never been done so far. The present study on the metamorphic rocks, in the western part of the Tayuan-Xiguitu fault will contribute to better understanding the collision tectonics between the Ergun block and the Xing'an block.

### Reference:

Li and Oyang, 1998, *Marine Geology & Quaternary Geology*, 18, 45-54; Sengor et al., 1993, *Nature*, 364, 299-307; Ye, H.W. et al., 1994, *M-SGT Geology Issue*, 73-83; Ge, W.C. et al., 2005, *Chinese Science Bulletin*, 50, 1239-1247; Sui, Z.M. et al., 2006, *Global Geology*, 25, 229-236; Li, R.S., 1991, *Heilongjiang Geology*, 2, 19-32.

キーワード: high P/T metamorphism, blueschist, Tayuan-Xiguitu fault, northeastern China

Keywords: high P/T metamorphism, blueschist, Tayuan-Xiguitu fault, northeastern China

SCG008-P10

会場:コンベンションホール

時間:5月26日 14:00-16:30

## モンゴルにおける多重衝突・沈み込みシステム Multiple subduction and collision system in Mongolia

大和田 正明<sup>1\*</sup>, 小山内康人<sup>2</sup>, 中野伸彦<sup>2</sup>, 足立達朗<sup>2</sup>, 米村和紘<sup>2</sup>, 吉本 紋<sup>2</sup>, ジャルガラン<sup>3</sup>, ボルドバートル<sup>4</sup>, サティッシュクマール<sup>5</sup>

Masaaki Owada<sup>1\*</sup>, Yasuhito Osanai<sup>2</sup>, Nobuhiko Nakano<sup>2</sup>, Tatsuro Adachi<sup>2</sup>, Kazuhiro Yonemura<sup>2</sup>, Aya Yoshimoto<sup>2</sup>, S. Jargalan<sup>3</sup>, C. Boldbaatar<sup>4</sup>, M. Satish-Kumar<sup>5</sup>

<sup>1</sup> 山口大学, <sup>2</sup> 九州大学, <sup>3</sup> モンゴル科学技術大学, <sup>4</sup> モンゴル鉱産局, <sup>5</sup> 静岡大学

<sup>1</sup>Yamaguchi University, <sup>2</sup>Kyushu University, <sup>3</sup>Mongolian Univ. Sci. & Tech., <sup>4</sup>Mine. Res. Auth. Mongolia, <sup>5</sup>Shizuoka University

The collage tectonics of micro-continents formed the Central Asia during Proterozoic to Mesozoic time in terms of the multiple accretion and collision. Mongolia is made up of the four micro-continents; i.e, Baltica, Siberia, North China and Tarim. There are four geological division; continental blocks, subduction complexes, sedimentary basins and Cenozoic cover sequences. The petrological study including geochronology and isotopic geochemistry of various types of igneous rocks provides us some information for tectonic processes during the assembly of microcontinents. Here, we address geochemistry and geochronology of igneous rocks, mainly granitic rocks and discuss magmatic history of the western part of Mongolia.

The study area, western part of Mongolia, is underlain by low- to high-grade metamorphic rocks accompanied by sedimentary rocks and intrusive rocks. The granitic rocks mainly intrude metamorphic rocks with various metamorphic grades. The metamorphic rocks are composed mainly of pelitic gneisses and amphibolites in the northern part of Bulgan, western Mongolia, the Mongolian Altai Mountains. The pelitic gneiss locally contains Grt, St and Ky as porphyroblasts, suggesting medium-pressure type amphibolite-facies conditions (Nakano et al., 2010). Grt-Ms granite intrudes Grt-St-Ky gneiss as post-kinematic dikes or stocks. Grt-Ms granite is free from any deformations. The chemical compositions of Grt-Ms granite show 71-75 wt% in SiO<sub>2</sub> and peraluminous (alumino-saturation index: A/CNK=1.02-1.29). Other geochemical characters indicate enrichment of K<sub>2</sub>O, Rb and depression of Nb, Y. On the other hand, syn-kinematic deformed granitic rocks containing garnet grains are exposed on the Hanhohiyn Mountains, north-west Mongolia. Bt-Ms granite and Hbl-Bt diorite occur as stocks around Tsagaanhayrhan. The granitic rocks show granoblastic texture with or without foliation, and tonalite to granodiorite in compositions. SiO<sub>2</sub> contents range from 63 to 66 wt% and alumino-saturation index is more than 1.1. These granitic rocks, high-Sr granite are characterized by high-Ba and -Sr contents similar to some collision related granitic rocks. Hbl-Bt diorite possesses 52 wt% in SiO<sub>2</sub> and A/CNK=0.82 and the chemical composition of the diorite is similar to that of within-plate basalt. In addition to these granitic rocks, pink-colored granites occur probably as stocks in the southeastern part of the Hanhohiyn Mountains. These granites contain pink-colored K-feldspar and show no signs of deformation features. Their chemical compositions are consistent with alkali granite having 70-73 wt% in SiO<sub>2</sub> and 8-11 wt% in total alkali, and the pink-colored granites possess low-Ba and -Sr contents.

We conducted monazite EMP dating for selected samples from both regions. Grt-Ms granite from the northern part of Bulgan gave an age of 262 Ma, whereas the syn-kinematic granite and the high-Sr granite from the Hanhohiyn Mountains show ages of 506 Ma and 493 Ma, respectively. It is considered that the magma activities of each region occurred at different ages or events.

Various types of intrusive rocks are exposed on the Mongolian orogenic belt. Most of granitic rocks from both regions are plotted within the fields between volcanic arc granite and syn-collisional granite in some discrimination diagrams. Therefore, the magmatic processes combined with metamorphic evolution in the western part of Mongolia were formed by multiple subduction and collision events during formation of Central Asian Orogenic Belt.

キーワード: モンゴル, 中央アジア造山帯, 大陸衝突帯, 沈み込み帯, EMP モナザイト年代

Keywords: Mongolia, Central Asia Orogenic Belt, Continental collision zone, Subduction zone, EMP monazite ages

SCG008-P11

会場:コンベンションホール

時間:5月26日 14:00-16:30

## Petrology and geochronology of metamorphic rocks in the Hanhohiyn Mountains, the northwestern part of Mongolia.

## Petrology and geochronology of metamorphic rocks in the Hanhohiyn Mountains, the northwestern part of Mongolia.

足立 達朗<sup>1\*</sup>, 小山内 康人<sup>1</sup>, 中野 伸彦<sup>1</sup>, 大和田 正明<sup>2</sup>, Madhusoodhan Satish-Kumar<sup>3</sup>, Sereenen Jargalan<sup>4</sup>, C. Boldbaatar<sup>5</sup>, 米村 和紘<sup>1</sup>, 吉本 紋<sup>1</sup>

Tatsuro Adachi<sup>1\*</sup>, Yasuhito Osanai<sup>1</sup>, Nobuhiko Nakano<sup>1</sup>, Masaaki Owada<sup>2</sup>, Satish-Kumar Madhusoodhan<sup>3</sup>, Jargalan Sereenen<sup>4</sup>, C. Boldbaatar<sup>5</sup>, Kazuhiro Yonemura<sup>1</sup>, Aya Yoshimoto<sup>1</sup>

<sup>1</sup>九州大学, <sup>2</sup>山口大学, <sup>3</sup>静岡大学, <sup>4</sup>モンゴル科学技術大学, <sup>5</sup>モンゴル鉱物資源石油管理庁

<sup>1</sup>Kyushu University, <sup>2</sup>Yamaguchi University, <sup>3</sup>Shizuoka University, <sup>4</sup>Mongolian Univ. Sci. Tech., <sup>5</sup>Miner. Res. Auth. Mongolia

Mongolia is situated in the Central Asian Orogenic Belt (e.g. Mossakovsky et al., 1993), which extends between the Siberian craton, Baltica craton, North China craton and Tarim craton. It is composed of subduction-accretion complexes and developed from c. 1000 Ma to c. 250 Ma (Windley et al., 2007). Metamorphic rocks are partly distributed in the western part of Mongolia. Metamorphic history would be a key for understanding the tectonic evolution of this area, however detailed study on metamorphic rocks has not been well-documented, including geochronology.

Recent fieldwork of the Japan-Mongolia Joint Geological Research has identified several occurrences of high-grade metamorphic rocks in the Mongol-Altai Mountains and Hanhohiyn Mountains (e.g. Nakano et al., 2010). In this study, we report petrology and monazite U-Th-Pb EPMA ages on metamorphic rocks occurring in the Hanhohiyn Mountains.

Hanhohiyn Mountains is located in the northwestern part of Mongolia. The mountains are mainly composed of marble, meta-quartzite and pelitic gneisses intercalating with lenticular garnet-amphibolite, garnet-orthopyroxene gneiss and so on.

Pelitic gneiss is mainly composed of quartz, plagioclase (An=31), garnet (Alm68Sps8Prp20Gr4), biotite (XMg=0.53) and sillimanite. Kyanite occurs only as inclusions in garnet and plagioclase, which indicates that kyanite is prograde relic. Most minerals have almost homogeneous chemical composition. Metamorphic condition for quartz+plagioclase+garnet+biotite+sillimanite assemblage is estimated as 650-700 degree of Celsius and 5.5-6 kbar based on the garnet-biotite thermometer and the GASP barometer. Monazite grains in pelitic gneiss show slight chemical heterogeneity and occur in matrix and as inclusions in the major constitute minerals. The monazites yield 500-510 Ma which would be interpreted as the timing of metamorphism within the sillimanite stability field.

In the Central Asian Orogenic Belt, at least two distinct age clusters are recognized: 250-300 Ma and 480-510 Ma. Ca. 500 Ma ages have been recognized at the marginal region of the Siberia Craton including the Hanhohiyn Mountains. This suggests that the Hanhohiyn Mountains would record the early stage event related to the formation of the Asia continent.

### References

Mossakovsky et al. (1993) Central Asian fold belt: geodynamic evolution and history of formation. *Geotectonics*, 6, 3-33.

Nakano et al. (2010) Metamorphic rocks from the Mongol-Altai: A first report of Japan-Mongolia Joint Geological Research (2009-2013). *Journal of Geological Society of Japan*, 116, XI-XII.

Windley et al. (2007) Tectonic models for accretion of the Central Asian Orogenic Belt. *Journal of the Geological Society*, London, 164, 31-47.

Keywords: Hanhohiyn Mountains, Mongolia, metamorphic rocks, monazite EPMA dating