

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-01

Room:201B

Time:May 26 08:30-08:45

Introduction to the Joint session

Hiroshi Kitazato^{1*}

¹JAMSTEC

Principal goal of the session is to discuss proper approaches to reconstruct the Earth climate system. Many of marine calcified organisms are used as biological proxies for climate change researches at present and past. For progressing the field of this science, we should precisely understand the biocalcification mechanisms and the incorporation of proxy signals. Thus, we have invited contributions related to the biocalcification, calibration and validation of marine proxies.

Keywords: biogeosciences, proxy development, culture, biology, chemical characters, biomineralization

BPO003-02

Room:201B

Time:May 26 08:45-09:00

Vesicle Dynamics In Foraminifera

Nina Keul^{1*}, Lennart Jan de Nooijer², Gerald Langer¹, Jelle Bijma¹

¹Alfred Wegener Institute, ²Utrecht University

The calcium carbonate produced by foraminifera contains a wealth of information (so called "proxies") for paleoceanographical studies. The environmental conditions, under which they calcified, such as temperature, salinity, productivity and the ocean carbonate chemistry, are recorded in the geochemical composition of their shells (isotopic signature, trace metals concentrations). However, foraminifera maintain their own distinct (trace) metal homeostasis, which results in characteristic and species specific elemental ratios as well as in particular isotope fractionation ("vital effect"), which is offset from inorganic-thermodynamic equilibrium. In order to improve the predictive capability of the proxies it is important to identify the physiological processes controlling the chemical and isotopic composition of foraminiferal calcite.

Foraminifera precipitate calcite from modified seawater vacuoles, which are incorporated via the process of endocytosis (Erez 2003). We studied this vacuolisation process on juvenile foraminifera using fluorescent dyes (FITC-Dextran and Calcein) and confocal imagery. Specimens of the shallow, benthic foraminifer *Ammonia tepida* were incubated for various periods (ranging from 2-10 hrs) in seawater labelled with fluorescent dyes. After incubation the Petri dishes containing the foraminifera were carefully washed with seawater to remove the dye. Once the dishes were filled with fresh, unlabelled seawater the dynamics of the formed endocytosis vesicles were followed by means of confocal microscopy until chamber formation occurred. Depending on the length of the incubation period and the physiological status of the cell (prior/ after chamber formation) different patterns of vesicle cycling could be observed.

Keywords: foraminifera, cytology, biomineralization

BPO003-03

Room:201B

Time:May 26 09:00-09:15

Characterization of the shell matrix proteins of calcareous foraminifera

Isao Sarashina^{1*}, Takashi Toyofuku², Kazuhiko Fujita³, Masashi Tsuchiya², Kazuyoshi Endo¹, Hiroshi Kitazato²

¹Earth and Planetary Sci., Tokyo Univ., ²JAMSTEC, ³University of the Ryukyus

Foraminifera are unicellular marine organism. Many of them have calcitic shells and are among the major calcium carbonate producers in the oceans. Their calcareous shells are widely used for stratigraphic and paleoenvironmental analyses. Like the case of skeletons of many other organisms, foraminiferan shell formation is thought to be controlled to a large extent by organic macromolecules such as proteins. But none of them have been identified so far.

Baculogypsina sphaerulata and *Calcarina gaudichaud* are common calcareous foraminiferan species, and were collected from Okinawa, Japan.

Both of the soluble and insoluble organic shell materials of *B. sphaerulata* and *C. gaudichaudii* were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). For the soluble shell materials of the two foraminifera, no signal was detected when stained with Coomassie Brilliant Blue (CBB), but strong smear bands were seen when stained with silver. CBB is known to stain most proteins, but it does not usually stain very acidic proteins. Thus we also stained the soluble shell materials with Stains-all which stains cation-binding proteins blue and the other proteins pink. When stained with Stains-all, the soluble shell material of *B. sphaerulata* appeared as a blue smear, and that of *C. gaudichaudii* showed a blue protein band of 66kDa as well as a blue smear along the lane. These results suggest that the shells of both *B. sphaerulata* and *C. gaudichaudii* contain soluble acidic proteins.

For the insoluble shell materials of the two foraminifera, no signal was detected when stained with CBB. But blue smears were seen for both species when stained with Stains-all. The signals did not appear so blue as that of soluble materials. Thus, insoluble materials are unlikely to be so very acidic, and perhaps they play a different role from that of the soluble materials.

In order to clarify the process and the underlying mechanism of foraminiferan shell formation, it is necessary to understand the function of shell matrix proteins. The results of this study provide a basis for those experiments in future.

BPO003-04

Room:201B

Time:May 26 09:15-09:30

Foraminiferal cellular pH control under low pH environment in the laboratory

Takashi Toyofuku^{1*}, Lennart Jan de Nooijer², Hidetaka Nomaki¹, Masashi Tsuchiya¹, Hiroshi Kitazato¹

¹JAMSTEC, ²University of Utrecht

Foraminifera have been considered one of the major carbonate producers in marine environments. Their calcareous tests are broadly utilized as paleo-environmental indicators in various studies of earth science because their tests act as geochemical archives. Although foraminifers are unicellular organisms, they precipitate a highly complex and fine-decorated test. Their calcification mechanisms must thereby be strongly controlled by biological activity and knowledge about the cytological control on carbonate precipitation is rapidly accumulating. In particular, mechanisms of calcium and carbonate ion uptake into foraminiferal cells from ambient seawater are of great interest. Our previous studies showed the potential to understanding the biomineralization of foraminifera by the application of fluorescent indicators. Development of fluorescent indicators allow us to visualize the spacial distributions of cytological chemical environment (e.g. pH and calcium concentrations) and organelles in living cell. Observed results show that foraminifera operate their biomineralization by controlling their intracellular environments. Acidified oceanic condition by antigenic high pCO₂ becomes global environmental problem in the near future. Therefore, we have extended our approach in order to observe foraminiferal cellular environment under a range of environmental pH's through laboratory culture experiments.

Keywords: foraminifera, calcium carbonate, ocean acidification

BPO003-05

Room:201B

Time:May 26 09:30-09:45

Improvement of culturing experiment of planktic foraminifera using the fluorescent indicator calcein

Daiki Kawata^{1*}, Minoru Ikehara²

¹Dept. Natural Environ.Sci.,Kocchi Univ., ²Center Adv. Marine Core Res.

To reconstruct the paleoenvironmental variations using trace elements and stable isotopes of planktic foraminiferal calcite, it is necessary to determine the coefficient by vital effect based on culturing experiment. In the conventional culturing method, however, it was necessary to conduct the culturing experiment every one individual to distinguish the newly formed chambers. In that case, a lot of time was spent to obtain enough number of cultured specimens. Therefore, we attempted to construct a new culturing method of planktic foraminifera using the fluorescent indicator calcein.

Living planktic foraminifera were collected using a plankton net from the Tosa Bay, south of Kochi in Japan. Specimens of planktic foraminifera were cultured in calcein solution, which was adjusted at a concentration of 10 mg/L. We cultured individuals in calcein solution under two conditions for 48 hours and 72 hours temporarily to determine time necessary for the dye. Cultured individuals were moved to natural seawater condition after the culturing experiment in calcein solution.

We observed the calcein labeled chambers using a fluorescence microscope. As a result, fluorescence was observed in not only the newly formed final chamber but also the chamber that had been added before. Therefore, it is suggested that the surface of chambers, which were added before started culture, were also calcified in the experiment. On the other hand, it has been observed that a final chamber, which was added in natural seawater after culturing in calcein solution for 48 hours did not fluorescence. Based on our culturing experiment, the fluorescent indicator calcein is a good tool to distinguish the newly formed chamber. In addition, a temporarily culturing in the calcein solution is effective to culture more individuals of planktic foraminifera. Time necessary for temporary culturing in calcein solution is enough in 48 hours.

Keywords: planktic foraminifera, culture, fluorescent indicator calcein

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-06

Room:201B

Time:May 26 09:45-10:00

Effects of thermal and salinity stresses on coral calcification: approach by aposymbiotic and symbiotic primary polyps

Mayuri Inoue^{1*}, Akira Iguchi², Shinmen Kotaro¹, Sakai Kazuhiko², Atsushi Suzuki³, hodaka kawahata¹

¹AORI, the Univ. Tokyo, ²Univ the Ryukyus, ³GSJ, AIST

In order to better understand the effects of high thermal and low salinity stresses on coral calcification from the aspect of coral-algal symbiosis, aposymbiotic (lacking symbionts) and symbiotic coral primary polyps were experimentally exposed to several seawater temperatures (27 ~ 33 °C) and salinities (26 ~ 34 psu). Symbiotic polyps showed non-linear calcification responses to thermal stresses whereas aposymbiotic demonstrated linear increase of calcification responses according to the increase of temperature. Both aposymbiotic and symbiotic polyps showed the linear decreases of calcification rates according to the decrease of salinity. Our results suggest that future global warming might have positive and negative impact on coral calcification, and low salinity stress, which would be caused by increase of the frequency of local floods related to future climate change, would certainly decrease coral calcification despite the existence of symbiotic algae.

Keywords: coral polyp, growth rate, environmental stress, algal symbiosis

BPO003-07

Room:201B

Time:May 26 10:00-10:15

Interindividual differences in stable isotopes of benthic foraminifera: the profile of isotopic disequilibrium

Toyoho Ishimura^{1*}, Urumu Tsunogai², Takeshi Oi³, Shiro Hasegawa³, Fumiko Nakagawa²

¹Geological Survey of Japan, AIST, ²Hokkaido University, ³Kumamoto University

For more than 50 years, variations in stable carbon and oxygen isotopic compositions of calcium carbonate, especially foraminiferal shells, have been used for estimating paleoenvironments, such as global sea-level changes, paleotemperature, global deep-sea circulation, and huge methane release events from the seafloor. In particular, the stable isotopic compositions of calcareous benthic foraminiferal shells have been used as tracers to determine the paleoenvironment at the seafloor.

Several factors determine the isotopic values of benthic foraminiferal shells, including the isotopic composition and temperature of the bottom water, the organic carbon flux, and "vital effects". Major isotopic variations in some species of benthic foraminifera are already being utilized as paleoindicators of bottom water conditions, but recently researchers have begun to quantify in detail the relationship between the isotopic composition of benthic foraminiferal shells and ecological characteristics such as species' microhabitat and the organic carbon flux. Such precise calibration and validation of isotopic indicators in benthic foraminifera will broaden the range of their application as paleoenvironmental tracers. However, until recently, it has not been possible to analyze the stable isotopic compositions of small carbonate samples of less than 20 micrograms and obtain results with an acceptable error range, thus, each sample used for analysis included multiple individuals, resulting in isotopic values averaged across individuals. This limitation prevented precise association of shell compositions with microenvironments.

In this study, by using a custom-made analytical system to determine stable carbon and oxygen isotopes in specimens with submicrograms calcium carbonate; the quantity required by the system is less than 1/100 of that required by conventional analytical methods, we determined the stable isotopic values of small individual shells of deep sea benthic foraminifera from core-top samples from three sites in marginal seas of the northwestern Pacific to characterize the magnitude of interindividual variation in their stable isotope ratios. We expect the results to be useful for exploring which species are most appropriate to use as paleoindicators in paleoenvironmental studies.

Keywords: stable isotope, foraminifera, vital effect, proxy, microscale analysis, calcification

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-08

Room:201B

Time:May 26 10:15-10:30

Effects of carbonate leaching on foraminifer stable isotopes ratios

Stephen Obrochta^{1*}, Saburo Sakai², Toyoho Ishimura³, Yusuke Yokoyama¹, Atsushi Suzuki³

¹AORI, the Univ. Tokyo, ²IFREE, JAMSTEC, ³GSJ, AIST

Stable carbon and oxygen isotope ratios were measured on 125 individual epifaunal and infaunal benthic foraminifers from two discrete intervals in a shallow-water sediment core (~ 450 m) from the Timor Sea. Methane seeps are common in the area, likely resulting in significant precipitation of secondary calcite, the effects of which were assessed by subjecting foraminifers to varying degrees of pretreatment. All foraminifers received standard cleaning with ethanol and brief sonication. A subset were further cleaned and sonicated in a dilute HCl solution (~ 0.003 M). Foraminifer tests were photographed using both reflected light and scanning electron microscopes during the course of treatment to monitor the changing degree of secondary calcite contamination as increasingly aggressive cleaning methods were employed. While foraminifers subjected to treatment with HCl exhibit a lower relative standard deviation, the variance remains high relative to expected results. Therefore, a similar experiment is being conducted on living individuals (stained with Rose Bengal) taken from a nearby multicore with complementary isotope ratios of overlying seawater, porewater, and DIC.

Keywords: stable isotopes ratios, foraminifera, carbonate leaching

BPO003-09

Room:201B

Time:May 26 10:45-11:00

A Holocene diatom oxygen isotopes record from the Indian Sector of the Southern Ocean

Masako Yamane^{1*}, Yusuke Okazaki², Akira Ijiri¹, Minoru Ikehara³, Yusuke Yokoyama⁴

¹Grad. Sch. Sci., Univ. Tokyo, ²JAMSTEC, ³KCC, Kochi Univ., ⁴AORI, Univ. Tokyo

The oxygen isotope ratio ($d^{18}O$) of diatom frustules are promising as a quantitative proxy of past seawater temperature and seawater $d^{18}O$, analogous to foraminifer $d^{18}O$. Therefore, diatom $d^{18}O$ is expected to be used for paleoceanographic studies in polar regions where foraminifer shells are hard to be preserved in sediments. However, conventional methods for measurements on diatom $d^{18}O$ have various limitations, including difficulty of processing and necessity for large amount of samples. Here we present diatom $d^{18}O$ records from the Southern Ocean in order to reconstruct paleoceanographic change in the Southern Ocean and the Antarctic Ice Sheet fluctuations during the Holocene. Our analytical system is composed of inductive high temperature carbon reduction and continuous-flow isotope mass spectrometry. In this system, silica is reduced by carbon at 1600 degree C to produce carbon monoxide for isotope analysis. The carbon monoxide gas is directly introduced into mass spectrometry by helium gas. Our system is capable to measure ~100 microgram of diatom $d^{18}O$ safely without any cumbersome procedures. A piston core COR-1PC (54°16.04'S, 39°46.00'E; 2,864 m water depth; 408 cm core length) was collected from the Conrad Rise, Indian sector of Southern Ocean in January 2008 by R/V Hakuho-Maru. Sediments were mostly composed of diatom ooze. The age model was established using AMS ¹⁴C dating on planktonic foraminifera (*Neogloboquadrina pachyderma*, sinistral) with core top and bottom ages estimated as 813 cal BP and 10,192 cal BP, respectively. We measured diatom $d^{18}O$ from 43 samples. We found that; i) diatom $d^{18}O$ in the Conrad Rise fluctuates between 38 per mil and 43 per mil through the Holocene, ii) diatom $d^{18}O$ show periodic variation during early to middle Holocene, iii) fluctuations of diatom $d^{18}O$ are different from those of foraminifer $d^{18}O$. We interpret fluctuations in diatom $d^{18}O$ to be dominated by a sea water $d^{18}O$ signal because the diatom $d^{18}O$ fluctuations exhibit more volatility than is expected for sea surface temperature. The periodic variation of early to middle Holocene seen in the Conrad Rise is also seen in other parts of the Southern Ocean, which coincides with the Holocene Thermal Maximum. This suggests that the Antarctic Ice Sheet underwent periodic melting associated with warming since the end of the last ice age.

Keywords: diatom $d^{18}O$, foraminifer $d^{18}O$, Southern Ocean, Holocene, Holocene Thermal Maximum

BPO003-10

Room:201B

Time:May 26 11:00-11:15

Temperature dependence of Mg isotope fractionation in deep-sea coral: paleoceanographic implications as a new proxy

Toshihiro Yoshimura^{1*}, Masaharu Tanimizu², Mayuri Inoue³, Atsushi Suzuki⁴, Nozomu Isawaki⁵, Hodaka Kawahata³

¹GSFS&AORI, the University of Tokyo, ²KCC, JAMSTEC, ³AORI, the University of Tokyo, ⁴IGG, AIST, ⁵UMBI, Kochi University

This study presents magnesium isotopic composition and its temperature dependence of high Mg biogenic calcium carbonates to evaluate their potential proxy of paleo seawater temperature. Degrees of Mg isotope fractionation compared to present seawater were measured in deep-sea coral. The mean $\delta^{26}\text{Mg}$ value of deep-sea corals was -2.5 permil. Moreover, Mg isotope fractionation in deep-sea coral showed a clear temperature dependence from 2.5 to 19.5 degree. The observed temperature dependence of Mg isotope fractionation in deep-sea coral skeletons implies that a combination of proxy developments and further high-precision isotope analysis allows potential application of Mg isotopes of high-Mg calcite to an environmental proxy for water temperature. The mean Mg isotope value of large benthic foraminifera which are also composed of high-Mg calcite was -2.34 permil. Even though the precipitation rates of deep-sea coral, benthic foraminifera were several order of magnitude different, they both plot on the same regression line within uncertainty. This result suggests that kinetic isotope fractionation may not be a major controlling factor, and indicate a possible further application of Mg isotope values as temperature proxy. Deep-sea corals and benthic foraminifera also showed similar Mg isotope fractionation factor to inorganically precipitated calcite, and the slope of temperature dependence in Mg isotope fractionation is similar to that for an inorganically precipitated calcite speleothem. Moreover, Mg concentrations and the relationship between Mg/Ca and temperature were also similar between deep-sea corals and inorganically precipitated calcite. Taking into account elemental partitioning and the calcification rate of biogenic CaCO_3 , the similarity among inorganic minerals, deep-sea corals and benthic foraminiferas may indicate a strong mineralogical control on Mg isotope fractionation for high-Mg calcite.

Keywords: magnesium isotope, precious coral, MC-ICP-MS, temperature dependence, proxy

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-11

Room:201B

Time:May 26 11:15-11:30

Assessment of foraminiferal richness from deep-sea benthos using Illumina massive sequencing technology

Beatrice Lecroq^{1*}, Franck Lejzerowicz², Jan Pawlowski², Philippe Esling³, Loic Baerlocher⁴, Laurent Farinelli⁴, Magne Osteras⁴

¹JAMSTEC, ²University of Geneva, ³IRCAM, ⁴FASTERIS SA

New massive sequencing technologies are especially relevant to assess the diversity of environmental samples from the Deep Sea since they allow retrieving extensive genetic information from limited amount of material uneasy to collect at such remote places.

We collected unsieved sediment samples from the Deep Sea of five distinct geographic regions and sequenced a very short (36 nt) fragment of the foraminiferal SSU rDNA hypervariable region.

Among the numerous phylotypes resulting from our analyses, some of them have been identified to the genus or species level and most of identified OTUs were assigned to monothalamous (single chambered) taxa. Our results, in which multi-chambered orders account for only a minority part of the richness, contrast with the classical view of foraminiferal diversity based on micropaleontologically oriented study of fossilized species.

This study emphasizes the usefulness of such technology for environmental biomonitoring perspectives regarding climate changing and human activities impact on deep-sea environment.

Keywords: benthic foraminifera, deep sea, massive sequencing

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-12

Room:201B

Time:May 26 11:30-11:45

Genetic diversity of planktic foraminifera and the bipolarity of genotypes in the Pacific

Atsushi Kurasawa^{1*}, Masashi Tsuchiya², Hiroshi Kitazato², Hiroshi Nishi²

¹Tohoku University, ²JAMSTEC

Molecular phylogenetic analyses have revealed high genetic diversity within planktic foraminifer morphospecies. Molecular studies of planktic foraminifera suggest these genotypes exhibit distinct ecological preferences. Moreover, their potential differences of their ecology and habitats could affect their chemical and isotopic composition of the test. However, the phylogeography of planktic foraminifera in the South Pacific is yet to be revealed. This study shows the phylogeography of *Globigerina bulloides* in the South Pacific. Living planktic foraminifera specimens were collected during R/V Mirai cruise (MR08-06). Molecular phylogeny and Identification of genotypes were based on partial small subunit ribosomal RNA gene (rDNA). We confirmed that one bipolar genotype in the Atlantic (type IIa) also exhibits bipolar distribution in the Pacific. Our results also suggest that trans-equatorial dispersal occurred in the East Pacific genotype.

Keywords: Planktic foraminifera, Genetic diversity

BPO003-13

Room:201B

Time:May 26 11:45-12:00

Chlorophyll derivatives as proxies of the marine photosynthetic production and succeeding biogeochemical processes

Yuichiro Kashiyama^{1*}, Hisami Suga⁴, Daiki Fujinuma², Shinya Akutsu², Masami Kobayashi², Satoshi Ohkubo³, Hideaki Miyashita³, Shinya Nomoto¹

¹Dept. Chemistry, University of Tsukuba, ²Materials Sci., University of Tsukuba, ³Human Environ Studies, Kyoto University, ⁴Institute of Biogeosciences, JAMSTEC

Chlorophylls produced in the oceanic environment undergo potentially both abiotic and biotic decompositions. A majority of chlorophylls is to be destroyed through so-called Type II process that cleaves tetrapyrrole macrocycle oxidatively. A portion of chlorophylls, however, experiences the Type I process in which tetrapyrrole macrocycles are preserved intact with various degree of defunctionalization, hence surviving into sediments. These survived chlorophyll derivatives could be further altered chemically to be fossil porphyrins, red pigments extracted from sedimentary rock as old as the Proterozoic.

Although chlorophyll derivatives in marine sediments have been studied for almost a quarter century, our new HPLC method that excludes analytical artifacts revealed that major chlorophyll derivatives are pyropheophytins, cyclophorbide enols, chlorin carotenyl esters, and chlorin sterile esters. The chemical structures of these derivatives strongly suggest their formations through metabolic processes, not a simple abiotic product. In addition, derivatives of chlorophyll b, chlorophyll d, and bacteriochlorophyll a that should be derived from specific taxa are also commonly identified in marine surface sediments. These chlorophyll derivatives are thus potential proxies of not only photosynthetic productions but also metabolic, biochemical processes in the water column.

We pursue uses of these chlorophyll derivatives as proxies of water column biogeochemical processes by understanding taphonomy of chlorophylls. Here, we particularly focused in cyclophorbide enols which has turned out to be quantitatively most important after our new analytical method. It is thus important to elucidate the environment of its formation and the organisms/processes concerned. We considered various analytical processes for the best method quantifying cyclophorbide enols from both water and sedimentary samples. We also determined physicochemical properties on hemi-synthesized cyclophorbide enols.

Keywords: chlorophyll derivatives, chlorophyll d, fossil porphyrin, cyclophorbide enols, marine photosynthetic production, biogeochemical processes

BPO003-14

Room:201B

Time:May 26 12:00-12:15

Distributions of archaeal membrane lipid and DNA within the modern coastal shallow marine water column

Kazuyoshi MORIYA^{1*}, Michinobu Kuwae², Masanobu Yamamoto³, Tadao Kunihiro⁴, Hidejiro Onishi⁴, Mitsuyo Saito⁴, Hideki Hamaoka⁴, Takuya Sagawa², Junya Shibata⁴, Naoki Fujii⁴, Naoki Yoshie⁴, Koji Omori⁴, Hidetaka Takeoka⁴

¹Dep. Earth Sciences, Waseda Univ., ²Senior Research Fellow Center, Ehime U., ³Div. Earth System Sci. Hokkaido Univ., ⁴CMES, Ehime U.

TEX86 thermometry, based on the number of cyclopentane moieties in the glycerol dialkyl glycerol tetraether (GDGT) lipids of membranes of Crenarchaeota, has been utilized as a paleotemperature proxy for the last ten years (Schouten et al., 2002 and others). This proxy was derived from an empirical relationship between annual mean sea surface temperatures and TEX86 values of core top sediments (Kim et al., 2008). This empirical relationship has been subsequently supported by the results of incubation experiments of marine Crenarchaeota (Wuchter et al., 2004; Schouten et al., 2007). Based on analyses of particulate organic matter from the modern open ocean, the membrane lipids preserved within core top sediments were biosynthesized at about 100 m deep within a water column (Wuchter et al., 2005). The distributions of isoprenoid chain produced within a water column are finally preserved within deep sea sediments (Takano et al., 2010). However, because this proxy has been developed and designed for open ocean settings, applicability of this unique proxy to the shallow coastal ocean, which can be easily affected by the anthropogenic climate change, is still uncertain.

Here we present abundances and distributions of archaeal membrane lipids within the water column at Beppu bay in the Seto Inland Sea. Beppu bay is an archetypal silled basin of 70 m depth, with bottom water that are decoupled from the surface water in summer, producing anoxic conditions in the bottom water. The abundance of GDGTs in the anoxic water mass was considerably higher than those of the oxygen rich water mass. Nonetheless, low particulate organic carbon content in the anoxic water mass indicates that the excess concentration of GDGTs implies in situ biosynthesis rather than accumulation of organic matter settled from the overlying oxygen rich water mass. Calculated TEX86 values from these lipids show a significant positive correlation with the in situ water temperatures observed, confirming the appropriateness of this proxy for shallow coastal ocean paleothermometry. On the other hand, at least two different genotypes were identified from archaeal DNA distributions within the water column. One genotype identified in samples recovered from water depth greater than 30 m was not found in samples recovered from the very shallow water, implying GDGTs in deeper water are synthesized by the different genotype than those found in the surface water.

Kim, J.H., Schouten, S., Hopmans, E.C., Donner, B., Sinninghe Damste, J.S., 2008. *Geochimica et Cosmochimica Acta* 72 (2008) 1154-1173.

Schouten, S., Hopmans, E.C., Schefus, E., Sinninghe Damste, J.S., 2002. *Earth and Planetary Science Letters* 204, 265-274.

Schouten, S., Forster, A., Elda Panoto, F., and Sinninghe Damste, J.S., 2007. *Organic Geochemistry* 38, 1537-1546.

Takano, Y., Chikaraishi, Y., Ogawa, N.O., Nomaki, H., Morono, Y., Inagaki, F., Kitazato, H., Hinrichs, K.U., and Ohkouchi, N., 2010. *Nature Geoscience* 7, 858-861.

Wuchter, C., Schouten, S., Coolen, M.J.L., Sinninghe Damste, J.S., 2004. *Paleoceanography* 19, PA4028.

Wuchter, C., Schouten, S., Wakeham, S.G., Sinninghe Damste, J.S., 2005. *Paleoceanography* 20, PA3013.

Keywords: Archaea, TEX86, DNA, coastal shallow marine

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-15

Room:201B

Time:May 26 12:15-12:30

Effect of surface ocean stratification on the distribution of planktic foraminifers

Ralf Schiebel^{1*}

¹Universite d'Angers

Production and abundance of planktic foraminifer species are correlated to a varying degree to surface ocean temperature (SST), salinity, primary production (PP), i.e., nutrient utilization, availability of prey, and the turbidity of surface waters. The regional and seasonal variability of surface ocean stratification does encompass different environmental parameters. In turn, planktic foraminifers facilitate the reconstruction of past surface ocean stratification on a seasonal and longer term, for example, the modern and Pleistocene surface ocean of the northern Arabian Sea. Changes in stratification are confirmed by the ecological significance of planktic foraminifer species.

Keywords: planktonic foraminifera, stratification

BPO003-16

Room:201B

Time:May 26 12:30-12:45

Stable isotopic composition of polar planktonic foraminifera: Results from sediment trap study in the North Pacific

Azumi Kuroyanagi^{1*}, hodaka kawahata¹, Hiroshi Nishi², Makio C. Honda³

¹Atmosphere and Ocean Research Institute, ²The Tohoku University Museum, ³JAMSTEC

Planktonic foraminifera provide a record of the upper ocean environment through their species assemblage and individual tests. Oceanographic condition at high-latitude plays an important role in the changes in global ocean environment, thus it is significant to assess the detailed past ocean situation in this region. *Neogloboquadrina pachyderma* (Ehrenberg) mainly distribute and dominate in sub-polar and polar region, thus it is an important species for the reconstruction of paleo-oceanography in the high latitude. In the study, we investigated the changes in oxygen isotope of *N. pachyderma* (sin.) using a 3.5-year sediment trap sample (about two-weeks resolution), and inferred 1) vital offset value in the area, 2) size effect, and 3) apparent calcification depth. In this study area, offset values were approximately 1 permil throughout the sampling period for both size except for 2000 (around 0.8 permil offset). Oxygen isotope values of *N. pachyderma* (sin.) exhibit definite seasonal variation throughout the sampling period, 1998 -2001. $\delta^{18}\text{O}$ values of both small (125-180 μm) and large (180-250 μm) shells decreased in autumn (September-October) with a minimum around September-October, and increased in spring with a maximum value around April-May. They ranged from 0.58 to 2.53 permil for smaller shells and 0.52 to 2.27 permil for larger shell throughout the study period. The differences in $\delta^{18}\text{O}$ between small and large shells generally decreased during winter, and increased summer. During winter, water column is well mixed, and differences is small (0.14-0.21 permil), while water column become stratified during summer, and differences became big (0.30-0.51 permil). During stratified water column period, larger shell mostly represented lighter $\delta^{18}\text{O}$ values, up to 0.54-0.79 permil lighter values. The large seasonal change in difference of $\delta^{18}\text{O}$ suggests that the different $\delta^{18}\text{O}$ between shell size would be mainly affected by water column situation rather than individual kinetic/metabolic effect. During stratification period, larger shell and smaller ones would mainly calcify at 24-35m and ~45m water depth, respectively. On the other hand, both size mainly reflect the water environment at 45-55m to their shells during water column mixed.

Keywords: planktonic foraminifer, *Neogloboquadrina pachyderma*, oxygen isotope ratio, sediment trap, northwestern North Pacific

BPO003-P01

Room:Convention Hall

Time:May 26 14:00-16:30

Three sets of temperature proxy revealed from coral *Porites cylindrica*

Alrum Armid¹, Ryuji Asami^{2*}, Tandri Fahmiati³, M. Ali Sheikh¹, Hiroyuki Fujimura¹, Tomihiko Higuchi¹, Eiko Taira⁴, Ryuichi Shinjo¹, Tamotsu Oomori¹

¹University of the Ryukyus, ²University of the Ryukyus, ³University of Haluoleo, Indonesia, ⁴Aqua Culture Okinawa Co., Japan

Numerous studies on scleractinian corals, especially massive *Porites* corals, have been conducted to establish potential tracers (Sr/Ca, Mg/Ca, and U/Ca) of sea surface temperature (SST) (e.g., Beck et al., 1992; Min et al., 1995; Mitsuguchi et al., 1996). In particular, the Sr/Ca of aragonitic coral skeleton is widely used to reconstruct SST variability using modern and fossil corals (e.g., Gagan et al., 1998). It should be noted that most of previously published paleothermometers have been derived from massive *Porites* corals, thus evaluation of other coral species as useful proxy is also significantly required which may provide more detailed investigation of paleoenvironmental reconstruction with high temporal and spatial resolution. Recently some investigations suggested that the coral Mg/Ca ratio significantly reflects not only seawater temperature, but also skeletal growth effects (Inoue et al., 2007). In order to examine such effect, data on culture experiment of coral species having variable growth rate are needed.

With these objectives in mind, we conducted a laboratory culture experiment utilizing scleractinian branching coral *Porites cylindrica* to examine the incorporation of Sr, Mg and U into the skeletons under three temperature settings (22, 26, and 30 degrees C), and subsequently evaluate the reliability of Sr/Ca, Mg/Ca, and U/Ca as potential seawater temperature proxies. The advantage of using *P. cylindrica* is that it has a relatively higher growth rate of up to ~3 cm/year (Custodio III and Yap, 1997) compared to massive *Porites* corals. *P. cylindrica* is a branching stony coral widely distributed in the tropical-to-subtropical Indian and Pacific Oceans and is very common in shallow water near the coast (Veron, 2000). Hence, results of this study will provide new valuable information for reconstructing past SST variability, which will contribute to the development of paleoceanography and paleoclimatology.

Keywords: *Porites cylindrica*, paleothermometer, metal/Ca ratio, culture experiment, distribution coefficient

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-P02

Room:Convention Hall

Time:May 26 14:00-16:30

Long-term preservation of organic carbon originated from the foraminiferal cell under dysoxic/anoxic conditions

Hidetaka Nomaki^{1*}, Nanako O. Ogawa¹, Naohiko Ohkouchi¹, Hiroshi Kitazato¹

¹JAMSTEC

Benthic foraminifera are major components among deep-sea benthic ecosystems and are known as major consumers of phyto-detritus. However, subsequent digestion and mineralization processes of ingested organic matters are still unclear. Here we operated a long-term incubation experiment to reveal how the ingested organic matters were degraded or mineralized by benthic foraminifera under dysoxic to anoxic conditions. Two different organic matters, algae and bacteria, were labeled with ¹³C and fed onto the surface sediments in a closed system of *in situ* feeding core. After 1.5 years, the sediments were recovered and foraminiferal specimens were isolated from the sediments. Although the ¹³C-label was found at extremely low abundances in the bulk sediments and in overlying water as dissolved inorganic carbon, the label was found at high concentration in dead foraminiferal tests particularly in algae-added cores. Benthic foraminiferal feeding affects largely to the distribution and subsequent preservation/mineralization of organic matters produced at both the oceanic surface and seafloor. It further suggests foraminiferal test serves highly isolated space for preserving organic matters from surrounding environments.

Keywords: Sediment-water interface, Benthic foraminifera, organic matter, Carbon mineralization, burial

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-P03

Room:Convention Hall

Time:May 26 14:00-16:30

Perspectives in stable oxygen isotope analysis as paleo-thermometer in biogenic carbonates

Saburo Sakai^{1*}

¹JAMSTEC

In this presentation, I'll summarize the recent progress of the high-resolution analyses of stable oxygen isotopes of biogenic carbonates as paleo-thermometer. Especially, the cold-water coral and fish otolith are treated as materials in order to describe the $\delta^{18}\text{O}$ -thermometer and clumped-isotope thermometer (δ^{47}).

Keywords: biogenic carbonate, oxygen isotope, paleo-thermometer