

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-01

Room:303

Time:May 24 16:30-16:45

River Regime of Ghaggar River in the Mid to Late Holocene, Northern India

Hideaki Maemoku^{1*}, Yorinao Shitaoka², Tsuneto Nagatomo³, Hiroshi Yagi⁴

¹Hiroshima University, ²Kyoto University, ³Nara University of Education, ⁴Yamagata University

It is well known that huge cities of ancient civilization, such as Mohenjo-daro, were abandoned during the post-urban Harappan period. Many reasons for the declination of Harappan culture have been estimated since 19th century, for example huge earthquake, devastative flooding, climate change etc. The Lost Saraswati hypothesis would be one of major estimations for the declination. One of the leading candidates for present remnant of lost Saraswati is the Ghaggar-Hakra River that originates in the Lower and Sub Himalayas of northwestern India. It flows westerly in the Punjab plain and disappears into Cholistan desert as a dried river bed. Naruse(1974) and Yashpal et al.(1980) said the Ghaggar had used to connect with glacial area in the Higher Himalayas through the Sutlej or Yamuna River course, however been disconnected by piracy due to crustal uplift or rejuvenation of adjacent river. We have tried to examine the hypothesis from the view point of chronology of flood plains and sand dunes distributed along the Ghaggar River. We resulted that the Ghaggar was not the mighty Saraswati during mature Harappan period because sand dunes on either side of the Ghaggar had been formed before that.

Acknowledgement

This study was supported by the research project titled Environmental Change and the Indus Civilization (project 3-3) managed by Research Institute for Humanity and Nature, Japan.

Keywords: India, Ghaggar River, Holocene, Indus Civilization, Harappan Culture, OSL dating

HQR023-02

Room:303


Time:May 24 16:45-17:00

Rara lake, the largest lake in Nepal Himalaya, as a pull-apart lake embanked by glacial till

HIROSHI YAGI^{1*}, Hideaki Maemoku², Makoto Okamura³, Hiromi Matsuoka³, Toshiki Osada⁴, Hirofumi Teramura⁴, Yasuhiro KUMAHARA⁵, D.P. Adhikari⁶, V. DANGOL⁶

¹Fac. Edu., Yamagata University, ²Hiroshima University, ³Kochi University, ⁴R.I.H.N., ⁵Gunma University, ⁶Tri-Chandra Campus, Tribvan Iniversity

Bathymetric survey of Rara Lake that is the largest lake in Nepal Himalayas was carried out, using an ultra sonic sounder and GPS. The shoreline of Rara Lake is just adjacent to the piedmont line except its southwest corner, where an alluvial fan develops. The alluvial fan is dislocated by an active fault that shows down-throw of eastern side of the fan surface. The bathymetry of the western half of the Rara Lake is box-shaped with a flat bottom deeper than -160 meters below the lake level though the transverse profile of its eastern half shows V-shaped submerged valley. A western margin of the lake is bounded by a steep submerged cliff deeper than 160 meters. It is a continuation of the active fault. The eastern bank of the lake is fringed by narrow embankment composed of the detritus thicker than 100meters. It is very clear that blockage of the mouse of the rhomboid valley has primarily formed Rara Lake due to detritus embankment. How the box-shaped depression in the western end has been formed? The alluvial fan is cut by the active fault that down-threw its eastern side. The active fault is northwestern extension of the Talphi fault (Nakata,1982) that is one segment of the active fault system with dextral displacement along the Main Central Thrust and that steps its trace to Darma fault in the north of Rara Lake. The trace of the active fault clearly continues northwestward to the submerged steep wall of the western fringe of the lake. That implies western half of the lake is tectonic origin due to pull parting that is liable to occur at a step of active faults. Namely, the box-shaped depression located in the western half of Rara Lake has been formed as a kind of pull-apart basin at the step of a series of the dextral active fault, Darma-Talphi active fault system.



Keywords: Rara Lake, Nepal Himalaya, glacial till, embankment, pull-apart lake

HQR023-03

Room:303

Time:May 24 17:00-17:15

Changes in vegetation and fire regimes since the mid-Holocene around Lake Rara, western Nepal

Nao Miyake^{1*}, Arata Momohara², Atsunori Nakamura³, Yusuke Yokoyama³, Makoto Okamura¹, Hiromi Matsuoka¹, Hideaki Maemoku⁴, Hiroshi Yagi⁵, Vishnu Dangol⁶, Toshiki Osada⁷

¹Faculty of Science, Kochi University, ²Chiba University, ³University of Tokyo, ⁴Hiroshima University, ⁵Yamagata University, ⁶Tribhuvan University, ⁷National Institutes for the Humanities

Vegetation and fire regime changes since the mid-Holocene were investigated using pollen, plant macrofossil and macroscopic charcoal records in a core sediment (Rara09-4) (29.5347°N, 82.0933°E and 3,000 m a.s.l.) from Lake Rara in western Nepal. The pollen record was divided into two local pollen assemblage zones (I and II), even though it was co-dominated by *Quercus* and *Pinus*, with *Abies*, *Picea* and *Betula* pollen throughout the core. In zone I (ca. 6,000 to 3,000 yr B.P.), *Quercus* pollen had high values of more than 50% of total tree pollen. On the other hand, zone II (ca. 3,000 yr B.P. to present) is characterized by the decrease in *Quercus* pollen and increase in *Pinus* pollen. Most of fossil *Quercus* and *Pinus* pollen grains were *Q. semecarpifolia* and *P. wallichiana* types, based on their SEM identification. Fossil leaves of *Q. semecarpifolia* type were founded frequently in the both zones. Macroscopic charcoal influx increased progressively in zone I. Based on these fossil records and spatial patterns of forest vegetation in the present, during the mid to late Holocene, *Q. semecarpifolia* and *P. wallichiana* predominated in the forests mainly on the south-facing slopes, whereas especially on the north-facing slopes evergreen conifer forests consisting of *Abies* and *Picea* with *Betula* were established. The decline of *Q. semecarpifolia* and dominance of *P. wallichiana* in zone I may have been caused by changes in fire regimes associated with combined effects of climate changes and intensified land-use activities around the lake.

Keywords: fire regime, Holocene, Lake Rara, plant remains, pollen, vegetation

HQR023-04

Room:303

Time:May 24 17:15-17:30

Plant fossil assemblages of the Last Glacial Maximum at Nakazato, northern Kanto region

Rika Nishiuchi^{4*}, Arata Momohara², Shigeto Osato³, Kunihiro Endo¹

¹Nihon Univ. Geosystem, ²Chiba University, Horticulture, ³Doshitsu Research Corporation, ⁴Nihon Univ.

Valley-fill deposits were cropped out at Nakazato site, NW of Utsunomiya city, northern Kanto region. Peat and tephra deposits filled the valley, and the peat deposits are overlain by Shichihonzakura and Imaichi tephra of 13~14 ka. Around the middle horizon of the thick peat, Ogawa Scoria is intercalated, which was derived from Nikko volcanoes a little after AT(29 ka). Therefore this peat is inferred to have piled up in LGM. There are very few studies of plant fossil assemblages in LGM not only in the northern Kanto but also whole Kanto region. Therefore, reconstruction of vegetation based on pollen analysis and plant macrofossil research at Nakazato site is very significant.

The plant macrofossil assemblages showed a typical assemblage in the last glacial as *Picea*, *Tsuga*, *Abies* includes *Abies veitchii* and *Betula*. *Picea* and *Betula* fossil pollen were detected abundantly from the peat horizon overlying Ogawa Scoria.

The reconstructed vegetation in LGM at Nakazato site is discussed in comparison with the plant fossil assemblages and paleoenvironments in LGM from other sites in Kanto and the middle part of Honshu.

Geological survey and sampling in Nakazato site were supported by the Research Group of Nakazatohara Outcrop(Yoichi Nakamura).

Keywords: pollen analysis, plant macrofossil, The Last Glacial Maximum, northern Kanto region

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-05

Room:303

Time:May 24 17:30-17:45

Depositional environment change during glacial/interglacial cycle, based on fluvial terraces, around Lake Biwa

Daisuke Ishimura^{1*}

¹Dept. Geophysics, Kyoto Univ.

I clarified landform development around Lake Biwa based on geomorphological and geological survey. Although the lake level is controlled mainly by bedrock exposed along the course of the Seta River and not directly influenced by sea level, older to younger terraces are widely distributed. I interpreted that the formation and preservation of fluvial terraces were results of crustal movement and depositional environment changes during glacial/interglacial cycle.

Keywords: Lake Biwa, fluvial terrace, depositional environment, landform development, tephrochronology

HQR023-06

Room:303

Time:May 24 17:45-18:00

Distribution and chronology of buried terraces in the Arakawa and Menuma Lowland and correlation with surrounding rivers

Takeshi Ishihara^{1*}, Toshihiko Sugai¹, Shoichi Hachinohe²

¹Natural Environmental Studies, ²Cen.Envi.Sci.Saitama

Distribution of basal topography of the latest Pleistocene-Holocene incised valley fill (the Alluvium) was revealed based on analysis of more than three thousand borehole data and boring core samples in the middle and upper part of the Arakawa Lowland and the Menuma Lowland, central Kanto Plain (Ishihara *et al.*, submitted a,b). We will discuss the forming processes of the basal topography of the lowland and their correlation with those in the Tokyo Lowland, and other alluvial lowlands such as Naka, Tama, Obitsu, and Yoro River lowland.

The basal topography beneath the Arakawa lowland can be divided into five buried landform surfaces: buried fluvial terrace surface I (S-I) to IV and incised valley bottom (S-V) which basal gravel (BG) of the Alluvium at the bottom. S-I to S-IV are widely distributed in the Arakawa Lowland, whereas their distribution are less clear in the Menuma Lowland. S-V ranges from the Arakawa Lowland to the Menuma Lowland and extending to the western bank of the current Tone River. S-I, II, III are partly covered by the Kanto loam and As-YP tephra was found in the Kanto loam of S-III.

Six buried landform surfaces are identified in the lower part of the Arakawa Lowland and Tokyo Lowland (Matsuda, 1974; Ando and Watanabe, 1996): buried terrace surface Ar0, T₀, Ar1 (T₁), Ar2 (T₂), T₃, incised valley Ar3 (T₄), in descending order. S-I, II, III, IV, and V are correlated with surface Ar0, Ar1 (T₁), Ar2 (T₂), T₃, and Ar3 (T₄), respectively. Considering above correlation and recognition of As-YP from S-III, S-I can be correlated with the Musashino terrace surface, and S-II to S-IV are correlated with the Tachikawa terrace surface group. In the Arakawa and Menuma Lowland, the buried terrace surfaces can be recognized up to 65 km upstream from the present river mouth and the incised valley can be recognized up to 85 km upstream indicating that the influence of sea-level drop in the Last Glacial extended far inland in the central Kanto Plain, resulting in formation of several fluvial terrace surfaces.

Buried Tachikawa terrace group and incised valleys are also identified in Nakagawa, Tamagawa, Obitsugawa, Yorogawa, and other alluvial lowlands in the Kanto Plain (Kaizuka *et al.*, 1977; Kashima, 1982; Endo *et al.*, 1983). Correlation of basal topography of the Alluvium between in the Arakawa and Menuma Lowland, and in above other alluvial lowland, is discussed.

Ando and Watanabe (1996) the Quaternary Research (Japan),35,281-291. Endo *et al.* (1983) URBAN KUBOTA,21,26-43. Ishihara *et al.* (submitted a) the Quaternary Research (Japan). Ishihara *et al.* (submitted b) Geomorphology. Kaizuka *et al.* (1977) Quaternary Research,8,32-50. Kashima (1982) Geographical Review of Japan,56,679-694. Matsuda (1974) Geographical Reports of Tokyo Metropolitan Univ.,9,1-36.

Keywords: the Arakawa Lowland, the Menuma Lowland, buried terraces, the Last Glacial

HQR023-07

Room:303

Time:May 24 18:00-18:15

Late Holocene environmental change in alluvial lowlands around the Lake Hamana, central Japan

Yoshiki Sato^{1*}, Osamu Fujiwara², Eisuke Ono³, Masatomo Umitsu⁴, Kaoru Kashima⁵

¹Kyushu University, ²AIST, ³Niigata University, ⁴Nara University, ⁵Kyushu University

Lake Hamana is a brackish lake located on the Pacific coast of central Japan, facing the Nankai Trough. It covers ~65.0 km² and is connected to the Pacific via a single channel between the sand bars. The lake has a complicated coast line with many alluvial lowlands, which were originally created as drowned valleys incising the Pleistocene terraces through the Holocene transgression and have progressively buried by sediments from the rivers and resulted in a fresh water marsh. This environmental change is closely related to both of the Holocene sea-level change and crustal movements(Ikeya *et al.* 1990). We carried out coring survey at coastal lowlands of the lake, and reconstructed their environmental changes since the middle Holocene using the depositional facies, diatom assemblages and radiocarbon ages.

Rokken-gawa lowland is located at southeastern coast of the lake. Surface geology of this lowland consists of mud layer, peat layer and sand layer in ascending order. The peat layer is divided into the upper and the lower parts by the mud layer identified between them. Sato *et al.* (2010) has pointed out that environmental changes from fresh water to brackish water condition occurred at ca.5500 calBP and ca. 3400 calBP respectively in this lowland. Here we discuss the environmental change of this lowland with additional data of sediment cores. The mud layer, above and beneath the lower part of the peat layer, shows dominance of brackish-marine species, for example *Cyclotella striata*, *Cocconeis scutellum* and so on. We also recognized at an elevation of -3.0 to -4.5 m in the mud layer, *Staurosira construens*, fresh-brackish water species, increased abruptly to be dominant. On the contrary, in the peat layer, fresh water species are dominant, for example *Tabellaria fenestrata*, *Aulacoseira* spp. and *Pinnularia* spp. These data indicate that water salinity decreased three times since the middle Holocene in this lowland. Radiocarbons ages by Sato *et al.* (2010) suggest that these events have occurred around 3400 calBP, 5500 calBP and before 5500 calBP.

In Shinjo lowland, located at southwestern coast of the lake, surface geology consists of mud layer, peat layer and sand layer in ascending order. The peat layer is subdivided into three parts by mud layer. We obtained new radiocarbon ages from each of these parts: 6420-6640 calBP (at the basis of the lower part), 4970-5300 calBP (in the middle part) and 5640-5900 calBP (at the basis of the upper part). The main components of diatom species in the mud layer are brackish-marine species, for example *Cyclotella striata*, *Cocconeis scutellum* and *Achnanthes submarina*. On the other hand, the peat layer shows abundance of fresh water species and dominance of *Staurosira construens*. These data imply that water salinity decreased three times in this lowland. On the basis of the radiocarbon ages, we infer that these events occurred around 6500 calBP, 5700 calBP and 5100 calBP, respectively.

According to the above data, synchronic occurrence of water salinity decreasing event is recognized between the two lowlands around 6500-5000 calBP. This period is also synchronous with the enclosure of bay by sand bars estimated from foraminiferal assemblages taken from lake bed sediments (Matsubara 2001). This suggests that environmental change at the two lowlands was also caused by the development of the sand bars sheltering the lake from the Pacific to decrease the seawater influx to Lake Hamana. Lake water salinity around 6500-5000 calBP is fluctuated drastically in short period. This indicates the formations and collapses of the sand bars occurred repeatedly.

Reference

Ikeya, N. *et al.* 1985. Geosci. Repts. Shizuoka Univ., 11, 171-179. (in Japanese).

Matsubara A. 2001. The Hiyoshi review of the social sciences, Keio Univ. 11: 20-32. (in Japanese).

Sato Y. *et al.* 2010. JGU, HQR010-P25.(English abstract)

Keywords: environmental change, sand bar, diatom assemblages, Lake Hamana, Holocene

HQR023-08

Room:303

Time:May 24 18:15-18:30

Use of electrical conductivity of Holocene deposits in the Nobi plain to analyze depositional environment

Yuichi Niwa^{1*}, Toshihiko Sugai¹, Ryutaro Naruhashi¹

¹Environmental Studies, KFS, UT

Electrical conductivity (EC) method is simple approach for estimation of depositional environment. Naruhashi et al (2010) indicates the possibility that EC records rapid changes of depositional environment in inner bay floor, on the basis of rapid increasing and decreasing of EC through the faulting event horizons on the Kuwana fault.

In this study, the usefulness of EC of stirred Holocene deltaic sediments from three cores from the Nobi Plain, Japan, was assessed as a proxy for marine transgression and regression. At first, the influence of grain size on EC was evaluated, because for same salinity, finer sediments tend to show higher EC (Yokoyama and Sato 1987). Lack of correlation of EC with mud content for the sediments with >20% mud suggested that permeability was a negligible factor, based on the correlation of mud content to permeability. Then, EC was compared with facies analysis results and the ratios of marine diatom species. Marine deposits showed high EC (>0.9 mS/cm), terrestrial deposits low EC (<0.4 mS/cm), and brackish deposits intermediate. Because it was positively correlated with the percentage of marine diatom species, EC in inner bay deposits primarily reflected salinity. In the YM core (the youngest of the three cores), EC of the inner bay deposits was weakly positively correlated with clay content, which controls pore water content. EC of inner bay clayey deposits was generally higher in YM than in KZN, suggesting that the lower EC in older clayey sediments results from compaction. These findings suggest that the EC values of fine sediments initially record salinity but may decrease gradually over time under the influence of compaction. Thus, to reconstruct the original salinity, the effects of compaction and of grain size distribution and especially clay content should be evaluated.

Keywords: salinity, Holocene, depositional environment, electrical conductivity, Nobi plain

HQR023-09

Room:303

Time:May 25 08:30-08:45

Floodplain evolution in lower reach of the Stung Sen River, central Cambodia

Naoko Nagumo^{1*}, Toshihiko Sugai¹, Sumiko KUBO²

¹The University of Tokyo, ²School of Education, Waseda University

The Stung Sen River which has the largest drainage basin in the Tonle Sap water system flows in an arc at central Cambodia region. At 230 km upstream from the Lake Tonle Sap, the river flows across a 7-km-wide floodplain and water level of the river changes 7 m annually because of two distinct seasons driven by monsoon. This research considers floodplain evolution of the Stung Sen River based on characteristics of the drilled cores in the floodplain and outcrops along the river in dry season, and ¹⁴C datings.

The channel of lower river is about 6-7 m depth and 70-100 m width with a rectangular cross section, which forms prominent meander scrolls along the channel. Although abandoned channels are well developed along the channel, natural levees are poorly formed and ground elevation decreases toward the channel. In addition, the floodplain is surrounded by uplands, so that flood water in monsoon season easily flows into floodplain and becomes remarkably wet especially along the channel. In floodplain, channel deposits of gravel to medium sand and back marsh deposits of silt to clay are accumulated at least 10 m on the basement of floodplain deposits. The channel deposits become thicker as it approaches to the channel, and the particle size becomes larger. ¹⁴C ages of channel deposits are comparatively new along the channel. In meander belt, which is composed of the channel, meander scrolls and abandoned channels, channel shifts and change of the deposits are prominent in particular, and that probably occurs in several decades order. ¹⁴C ages of back marsh deposits suggest to have been accumulated during last 35000 years by the accumulation rate of 0.1 -0.6 mm/yr with occasional channel deposits. This may suggest that the depositional pattern in the edge of floodplain has not drastically changed since the latest Pleistocene. Moreover, the oxidative-reductive reddish patches of the deposits which are highly recognized near the channel show the situation that the groundwater level alternates by monsoon.

Keywords: Stung Sen River, meander, boring core, ¹⁴C age, Cambodia

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-10

Room:303

Time:May 25 08:45-09:00

Relationship between slope collapse and landform evolution of active fold zone

Mamoru Koarai^{1*}, Takaki Okatani¹, Takayuki Nakano¹, Taku Komatsubara², Takahito Kuroki³

¹GSI of Japan, ²AIST, ³Fukuoka Univ. of Education

Large earthquakes occurred in Chuetsu District, such as the 2004 Niigata Prefecture Chuetsu Earthquake and the 2007 Niigata Prefecture Chuetsu-oki Earthquake. There are many active folds in Chuetsu District, and about 10cm uplift of Oginajo Anticline by the Chuetsu-oki Earthquake was detected by InSAR, and several slope collapses occurred in Nishiyama Hills. However, 0.5-1m uplift by the Chuetsu Earthquake was detected by difference between the DEM measured by photogrammetry using aerial photos taken before earthquake and the DEM by airborne laser survey measured after earthquake, and many slope collapses and landslides by earthquake are concentrated in Imokawa-River region. The authors try to survey the relationship between the growth of active fold by earthquake and the concentration of slope collapses by earthquake.

There is no terrace which is older than 15ka along Imokawa-River, because uplifting ratio of Imokawa-River region is large. As the relative elevation between oldest terrace and river bed is 20m in downstream and 30m in upstream, uplift ratio is 1.3-2 mm/year. If amount of uplift of the Chuetsu Earthquake is about 1m, the cycle of one earthquake would be 500-770 years. This research is supported by Grants in Aid for Scientific Research (22500994).

Keywords: active fold, slope collapse, Chuetsu Earthquake, Chuetsu-oki Earthquake, Imokawa River, Nishiyama Hills

HQR023-11

Room:303

Time:May 25 09:00-09:15

Late Quaternary Tephrostratigraphy of Marine Cores off Joetsu, Japan

Yuya Nakamura^{1*}, Toshihiko Sugai¹, Takeshi Ishihara¹, Antonio Freire¹, Ryo Matsumoto²

¹The University of Tokyo, ²The University of Tokyo

The piston cores and gravity cores from off Joetsu in the southeast Japan Sea contain widespread tephtras from distal volcanoes in the Kyushu, Chugoku, Chubu, and Hokkaido area. Identification of these tephtras is based on the mineral composition, and the morphologic feature and chemical composition of volcanic glass shards.

Since the Joetsu offing is located near the center of the Japanese islands, the place is advantageous in a tephtra study geographically. In addition, sedimentation rate in this area is faster than in the other spots of Japan Sea because this area is near the shore. This means that this area has potential to reconstruct paleoenvironment with high resolution. MD179-3312 core with the length of about 32 m is the main target of this study, and the recovery of the core is regarded as 100%.

On the basis of the stratigraphy, petrography, and chemical composition of volcanic glass shards measured by microprobe analyzer (SEM-EDS; JEOL JSM-6390LA), tephtras can be correlated to the As-K, Jo-2, AT, On-Ng, Aso-4, On-Kt, K-Tz, SK, Toya in descending order. As-K, AT, Aso-4, On-Kt, K-Tz, SK have been already found around the Sado Island, and the source of supply volcano has been known (Machida and Arai, 2003). Fernando(2010) finds two marker tephtras of Jo-1 (Joetsu 1 tephtra) and Jo-2, and former of which we compared with As-K. We discovered On-Ng and Toya at around this area for the first time. Furthermore, the tephtra found in the lower of Toya through this project is correlative to Hikage-2 (HK-2: Nagahashi et al., 2007), although SiO₂ of the volcanic glass is slightly lower than that of HK-2. There are some drift pumices (diameters are about 2-4mm) which might be correlative to On-Pm1 between K-Tz and SK based on its stratigraphic position. But it deviates from chemical composition of volcanic glass of On-Pm1.

This study expanded the isopachs of On-Ng and Toya. If the correlation between lower tephtra of Toya and HK-2 is true, it indicates at least one of the eruption ages of Toya (113-114ka: Ganzawa et al., 2007) or HK-2 (105.9ka: Nagahashi et al., 2007) should be revised. In addition, If the drift pumices above can be correlated with On-Pm1, it means the Pm1 pumices were transported by Jinzu River and drifted in the Japan Sea.

Acknowledgements

This study is a part of the GH 2010-11 project supported by METI and JOGMEC.

Keywords: Tephtra, Japan Sea, Chronology, Late Pleistocene, MD179, SEM-EDS

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-12

Room:303

Time:May 25 09:15-09:30

Local marine radiocarbon age offset of Hokkaido region

Yosuke Miyairi^{1*}, Yusuke Yokoyama¹, Hiroyuki Matsuzaki²

¹AORI, The University of Tokyo, ²MALT, The University of Tokyo

The natural radioisotope ^{129}I dating covers an important age range for geological samples. The iodine contains ^{129}I that is the long half-life radionuclide besides stable isotope ^{127}I . ^{129}I is a cosmogenic nuclide generated by cosmic rays in the atmosphere. It is possible to measure it by using $^{129}\text{I}/^{127}\text{I}$ at the iodine age. It is possible to measure iodine dating by using this equilibrium. However, present $^{129}\text{I}/^{127}\text{I}$ ratio is very high. Because a large amount ^{129}I was added by the human act of using nuclear power. Therefore, it is indispensable to know initial $^{129}\text{I}/^{127}\text{I}$ ratio that doesn't contain anthropogenic ^{129}I to measure ^{129}I age.

HQR023-13

Room:303

Time:May 25 09:30-09:45

14C analyses of sedimentation periods for shell middens at Higashimyo site, Saga prefecture

Toshio Nakamura^{1*}

¹Center for Chronological Research, Nagoy

The Higashimyo archeological site is located in huge Saga Plain, the largest in western Japan, at the northern end of Saga city, in Northern part of Kyushu. The Saga plain faces southwards the innermost part of Ariake bay, a bay of 15 km wide and 90 km long with relatively shallow water depths. A huge volume of shell fragments were accumulated at the Higashimyo site, and the most impressive archeological interest is that more than 400 of baskets knitted with twisted ropes made of thinly sliced wood were recovered. The knitted wooden baskets excavated here are the oldest (ca. 7800-7900 cal BP) recorded in the earliest Jomon period in Japan.

Six shell middens were detected at the Higashimyo site, and No.1 and No.2 middens were intensively excavated. For the No.1 midden, thickness of the mound was ca. 1.2m and the elevation of the uppermost layer was -1.1m, and for the No.2, the respective values were 1.5m and -0.5m. The starting and ending layers and several layers in between at No.1 and No.2 middens were 14C dated to estimate the duration period of the shell mound. We have collected shell fragments and terrestrial plant remains from the full layers, from the bottom to the top, of the two middens for 14C dating, to estimate the duration necessary to make up the total volume of the middens. Geographically it was certain that the middens had been formed during the sea-level-rising period a few thousand years after the start of Holocene.

The followings are revealed concerning the chronology of the Higashimyo site.

(1) According to 14C ages for terrestrial nuts for No.1 and No.2 middens, the sedimentation started almost the same time at around 8000 to 7850 cal BP for both middens. The termination time of accumulating shell fragments was, however, different. The No.1 shell middens stopped accumulation at 7950-7850 cal BP, and the No.2 did at 7800-7700 cal BP.

(2) Both terrestrial plants and marine shells were dated for both middens. 14C ages for the terrestrial plants are younger than those for the marine shells, by about 300-350 years. This apparent age difference is from the ocean reservoir effect. However, since the age difference is not established yet firmly (Nakamura et al. 2007), we adopted 0 years as the local reservoir correction value for the ocean reservoir effects, for further analysis.

(3) Our estimates yielded relatively shorter duration periods, i.e., 50-150 years and 150-250 years for the No.1 and No.2 middens, respectively.

Keywords: 14C age, Jomon shell midden, archeological site, sedimentation period, Jomon transgression, ocean reservoir effect

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-14

Room:303

Time:May 25 09:45-10:00

Upper Palaeolithic obsidian exploitation along the Shinano river in central north Japan

Akira Ono^{1*}

¹Meiji University

Present paper focuses on the formulating interpretative models for acquisition and transportation of lithic raw materials, with reference to obsidian and siliceous hard shale, choosing a case of Upper Palaeolithic camp site Mattobara in central-north Japan. The site is located on the left bank of the latest Pleistocene river terrace in the middle course of the Shinano River, dated back ca. 17 ? 14ka from the date of similar assemblages and the dating of tephra in a neighboring areas. The lithic assemblage of this site is rather simple and small Points are significant. Excavations uncovered totally 8,200 lithic tools, flakes, and chips. Dominant raw material is siliceous hard shale, and it takes about 80percent of total lithic materials. Small scale concentrations of this site are representing the activities of small mobile-groups along the Shinano River in the later period of Late Pleistocene. Source identification of obsidian from Mattobara Loc. C by X-ray fluorescence analysis identified that thirty samples are from Wada-Pass, and one is from Kirigamine both located in Nagano Prefecture. On the contrary to the case of Loc. C, 24samples are from north Fukaura (Aomori Pref.) and Oga (Akita Pref.) and only one sample identified as Tadeshina (Nagano Pref.) in the case of Loc. A. These quite different and wide range available areas indicate a complex system of lithic raw materials. Linear linkage of stepping-stone model will be discussed, irrespective of long distant direct procurement /or nearby direct procurement (so called embedded strategy).

Keywords: Uper Paleolithic, Shinano river system, Obsidian, Geologic source identification, Matobara site, Point tool industry

HQR023-15

Room:303

Time:May 25 10:00-10:15

Holocene environmental evolution based on lake deposits in Central Eurasia

Kunihiko Endo^{1*}, Toshihiko Sugai², Tsuyoshi Haraguchi³, Shigeru Suzuki⁴, Takashi Chiba², Haruka Matsuoka⁵, Hidehiro Sohma⁶

¹Dept Geosystem Sciences, Nihon University, ²The University of Tokyo, ³Osaka City University, ⁴Paleolabo, ⁵Tsukuba Gakuin University, ⁶Nara Women's University

To reconstruct environmental evolution in Ili Delta and Balkhash Lake areas, we have continued geological, geomorphological and paleoenvironmental researches under the Ili project, RIHN, since 2007 (Endo et al., 2010). Following the reconstruction of lake level change in the last 2000 years using 2007 core in the western part of Balkhash Lake, we took several cores in 2009 in the easternmost part of the lake, where is the deepest part of the lake.

These 2009 cores, covering almost Holocene, have been analyzed using pollen, diatom, and ostracod, and also geochemistry and magnetic properties. Two cores, 0901 and 0902, provide us continuous environmental records, which are combined with geological and geomorphological events and evidences.

Especially, in mid Holocene, highly lowered lake level stage is recognized from 5500 to 3500 years ago, when dry vegetation like desert was dominant and coniferous forest decreased. It suggests warm and dry climate is dominated.

In the lowest reaches of Heihe, the middle Inner Mongolian, about 20 paleoshore-lines (gravel bars) were dated by AMS 14C method. However, from 6000 to 3500 years BP there are no gravel bars, suggesting very low lake level dominant. After 3500 years BP, alluvial processes were reactivated and lake level attained to the highest (Endo et al., 2006).

Nearly same situation is reported in Mongolian lake, where 5800 to 3100 years BP, warm and dry conditions were dominant using diatom and pollen analyses (Wang et al., 2011). They proposed the existence of mid Holocene drought probably in east and central Asia.

We will compare the paleoenvironmental results in Central Eurasia with those of the eastern part during the middle Holocene, and discuss the extension of warm and dry climate, and the cause of such climate change.

References

Endo et al., (2010) Reconstruction of lake level and paleoenvironmental changes from a core from Balkhash Lake, Kazakhstan. *Reconceptualizing cultural and environmental change in central Asia: an historical perspective on the future, Ili Project*, 93-104.

Endo et al. (2006) Paleoenvironment and Migration of rivers, delta and lakes in the lowest reaches of Heihe River. *Project report on an oasis-region*, 5(2), 161-171.

Wang et al. (2011) A prolonged dry mid-Holocene climate revealed by pollen and diatom records from Lake Ugi Nuur in central Mongolia. *Quaternary International*, 229, 74-83.

Keywords: lake level change, Holocene, hyper arid stage, pollen analysis, ostracod analysis, diatom analysis

HQR023-16

Room:303

Time:May 25 10:15-10:30

Late Holocene fluvial landform development in the middle basin of Ili River, Kazakhstan

Hitoshi Shimizu^{1*}, Toshihiko Sugai¹, Hiroki Montani¹, Akio Sato², Nakayama Yasunori², Kunihiro Endo²

¹Natural Environmental Studies, KFS, UT, ²Department of Geosystem Sciences, College

Ili River runs from Tien-Sian Mountain as a source and flows into Balkhash Lake in the central Kazakhstan, Central Asia. The basin of Ili River belongs in the semi-arid area, and from the middle basin to the downstream basin is large plain called Ili Delta and observed paleo-channel clearly. Near Bakanas, modern channel runs to the west, meanwhile paleo-channel runs to the north. This means avulsion occurred.

About environmental change in Balkhash Lake, water level change past 2000 years is getting reconstructed for diatom and chemical analyses of drilling core (Endo et al., 2010; Chiba et al., 2010; Sugai et al., 2010 etc.). Shimizu and Sugai (2010) tried to reconstruct paleo-discharge of Ili River from the meander wavelength of Ili River course.

Since the 80 percent of influent water of Balkhash Lake is from the water of Ili River, to reveal the process and chronology of fluvial landforms in the middle and downstream part of Ili River will be useful for not only the key of clarification of landform development but also the key of environmental change in Balkhash Lake.

Fluvial landforms of Ili River including its paleo-channels were classified by field survey in August 2010 and interpretation of satellite images by Google Earth and DEM data from SRTM3.

The paleo-channels and fluvial surfaces were dated by AMS-14C method for the materials in the same fluvial sediment layer obtained through trenching survey. Particle analysis and magnetic susceptibility were also examined.

Fluvial landform along the middle part of Ili River was classified into five surfaces from T1 to T5.

T1 is the oldest, probably Pleistocene terrace covered with vegetated dune sand. At least, two cycles of upper-fining fluvial sediment units consist of T1. T2 is distributed along paleo-channel turning to the north at Bakbakhly. Aeolian sand dune partly covers T2. The paleo-channel is buried at least 30cm with fine aeolian sand.

T3 has paleo-channels whose meander length is larger than that of present channel. This implies that paleo discharge of T3 stage was larger than present discharge (Shimizu and Sugai, 2010). Two radiocarbon ages were obtained. One is about 1500 years ago from shell in sorted sand; the other is about 700 years ago from humic soil showing that the channel changed back marsh. That is to say, about 1500 years ago, the main channel of Ili River was forming T3, and after 700 years ago, this channel was abandoned and T3 became terrace.

T4 is distributed along the main stream of modern Ili River and formed after the main channel of Ili River moved to the west. Large flood is likely to flow into paleo-channels on T4.

T5 is the small floodplain of modern channel of Ili River and along the channel.

Fluvial landform development in the middle part of Ili River basin can be summarized as follows.

At first, Ili River ran on T2. Before 1500 years ago, an avulsion occurred in Ili River and T3 started forming. In this period, Ili River had large discharge and sediment. After T3 formed, Ili River channel shifted to west after 700 years ago, and T3 became terrace and T4 started forming. Today, T4 became terrace, and T5 is forming.

Reference

Chiba et al., (2010) Project report on an Oasis-region, 8, 1

Endo et al., (2010) Reconceptualizing Cultural and Environmental Change in Central Asia: An Historical Perspective on the Future.

Shimizu and Sugai (2010) Project report on an Oasis-region, 8, 1

Sugai et al., (2010) abstract Asian 2k symposium

Keywords: avulsion, 14C dating, paleo-channel, land classification

HQR023-17

Room:303

Time:May 25 10:45-11:00

Holocene environment changes in Lake Balkhash reconstructed by high-resolution XRF-core analysis and geomorphic survey

Toshihiko Sugai^{1*}, Montani Hiroki¹, Endo Kunihiko², Haraguchi Tsuyoshi³, Shimizu Hitoshi¹, Chiba Takashi¹, Kondo Reisuke⁴, Jumpei Kubota⁵

¹University of Tokyo, FSKC, ²Nihon University, ³Osaka City University, ⁴Geological Survey of Japan, ⁵Research Inst. for Humanity and Nature

Lake Balkhash is the largest terminal lake in central Asia. The lake sediment provides us to reconstruct climate changes during the Holocene. We obtained several sediment cores with a length of 5-6 m in the eastern part of the lake in 2009 under the Ili project, RIHN. This paper reports Holocene depositional environment changes in and around the eastern part of the lake based on high resolution (0.1 mm) XRF analysis of the longest two cores named as 901 and 902 coupled with the analysis of the Lepsy river deposits. Site 901 is about 20 km distant from the present Lepsy river mouth, while site 902 is 40 km, where is the heart of the eastern part of the lake and its deepest point. The lake bottom topography between proximal site 901 and distal site 902 is almost flat with a water depth of about 20 m. In contrast average slope gradient between 901 site and Lepsy river mouth is about 1/1000, almost equal to the slope of the lower reaches of the Lepsy river bed. For detail near the river mouth, the slope of delta plain is gentler and the slope of the delta front slope is steeper than 1/1000. Under such geomorphic conditions, water-level fall and rise cause quick incision and accumulation of the river channel.

Lepsy river valley had been filled with fine and partly organic sediments with a thickness of several meters from the early Holocene to 5500 years ago. Then, dominant process changed from accumulation to incision. River terraces were formed and well-sorted sandy sediments were transported downstream. At around 2000 years ago, the river turned to accumulation phase again. The river bed and river terrace sediments contain SiO₂ of 40-60% and CaO of 15-20%, while Fe₂O₃, MnO, and Al₂O₃ of less than 10 %.

Both 901 and 902 cores can be divided into three sedimentary units, the top and bottom units are whitish clayey sediments with high Ca content, indicating lots of calcium carbonates which include calcite, aragonite, and dolomite according to XRD analysis (Montani et al, 2011). The middle units deposited from 6000 to 3000 years ago based on AMS -14C dating and showed various kinds of stratified sedimentary structures including sand beds with plant fragments and gastropod fossils, parallel silty lamina (observed only in 901 core) and evaporites (in 902 core). The middle unit contains high SiO₂ indicating siliceous deposits driven by Lepsy river.

Using high resolution cps values (near 30,000 points in total with vertical interval of 0.2 mm for each core) of major elements (Si, Ca, Mg, K, Fe, S, Al, etc) provided by micro-XRF analyzer, we calculated Ca / Si ratio to know the change of relative abundance of chemogenic (and partly biogenic) vs. terrigenous components of the sediments. Trend of Ca / Si ratio curves between the two cores were markedly concordant with each other and the ratio of distal core 902 was always larger than that of proximal core 901. This suggests that the lake-level fluctuations dominantly control the sedimentary environment; water level fall brings the river mouth close to the core sites and causes the river channel incision and terrace formation, resulting that chemogenic sediments are remarkably diluted by increase of terrigenous deposits at the core sites. Water level rise should bring the opposite results. At around 5500 years ago, when Ca / Si ratio showed the minimum (nearly zero), evaporites (gypsum and magnesite) deposited at site 902 and fluvial sand deposited at 901. This implies that the lake level dropped almost 20 m below the present level, and the river mouth reached 901 and playa appeared at the center of the eastern part of the Lake Balkhash.

Keywords: Lake Balkhash, kazakhstan, Holocene, chemical analysis, lake level change, Lepsy river

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-18

Room:303

Time:May 25 11:00-11:15

Reconstruction of past environment by an ice core drilled from Grigoriev ice cap in Kyrgyz Tianshan, Central Asia

Nozomu Takeuchi^{1*}, Koji Fujita², Shuntarou Sera¹, Sachiko Okamoto², Kazuhiro Naoki³, Vladimir Aizen⁴

¹Chiba University, ²Nagoya University, ³JAXA, ⁴University of Idaho

In 2007, ice cores were successfully drilled on Grigoriev Ice Cap located in the Tien Shan Mountains, Kyrgyzstan. The elevation of the drilling site was 4600 m a.s.l. and entire core length was 87 m. We present stable isotope, mass balance, dust concentration, and pollen concentration in the last 200 years of the ice core.

Keywords: ice core, climate change, central asia, stable isotope, pollen

HQR023-19

Room:303

Time:May 25 11:15-11:30

Climate reconstruction using hydrogen and oxygen isotope ratios of tree-ring cellulose in Kyrgyz

Masaki Sano^{1*}, Takeshi Nakatsuka¹, Chenxi Xu¹, Chiyuki Narama², Jumpei Kubota², Kimitaka Kawamura³

¹Grad. Sch. Env., Nagoya Univ., ²Res. Inst. Humanity and Nature, ³Inst. Low Temp. Sci., Hokkaido Univ.

Hydrogen and oxygen isotope chronologies of tree-ring cellulose were established to understand past climate variations in southwestern Kyrgyz. A total of three core samples collected in 1998 from three individual junipers growing in an arid region were subjected to hydrogen and oxygen isotopic analysis. One core sample was analyzed over the last 485 years at pentad (5-year) resolution by simply mixing 5 adjacent rings, while the remaining two cores were individually examined for the last 50 years at annual resolution.

Correlation coefficients between the two trees for the inter-annual variations during the 1948-97 period are 0.60 ($p < 0.001$) and 0.30 ($p < 0.05$), respectively, for hydrogen and oxygen isotopes, suggesting that isotopic ratios are at least partly controlled by common climatic factors. Response analysis with ambient climatic records revealed, however, that the hydrogen and oxygen chronologies did not show significant correlations with monthly temperatures and precipitation at annual resolution. On the other hand, the pentad isotope records compared with the corresponding instrumental data showed significant positive correlations with March-April and March-August temperatures, respectively, for hydrogen and oxygen isotope chronologies. These results indicate that the pentad-based measurements result in the smoothing of short-term and micro-scale environmental noises, such as heterogeneous spatial and temporal distributions of soil waters in the arid land, recorded in annually resolved chronologies.

The 485-year hydrogen isotope chronology, which reflects spring temperatures, shows multidecadal to century-scale fluctuations, with a notable warm period of ca. 1710-1730 and a warming trend over the 20th century. Interestingly, the warm anomalies of the early 1700s, which can be recognized at many other places in the world, correspond to the end of the Maunder Minimum (1645-1715), thus may indicate some causal relationship between them. The oxygen isotope data, which reflect temperatures in spring and summer seasons, show rough similarity in the high-frequency domain with the hydrogen isotope chronology, since oxygen and hydrogen isotopes of precipitation are highly correlated with each other. However, the long-term variations substantially differ from each other, indicating that different factors control isotopic ratios in the low-frequency domain. One important factor is relative humidity, to which oxygen isotopes are known to be sensitive rather than hydrogen isotopes. Though we were not able to compare the isotope records with relative humidity due to the lack of instrumental data, if we consider the effect of relative humidity, it is possible to explain the reason why oxygen isotopes showed the significant correlation with summer temperatures, which were not correlated with hydrogen isotopes, as follows. In the studied region, summer precipitation is much less than spring one, through which only spring temperatures can be recorded in tree-ring hydrogen isotopes. On the other hand, leaf water oxygen isotope enrichment occurs during photosynthesis in summer, controlled by the summer temperature through summer relative humidity. We therefore interpreted that the oxygen isotope chronology capture both of relative humidity in summer and temperatures in spring, and the divergence of long-term variations between the oxygen and hydrogen isotope chronologies originates from relative humidity. When the 485-year oxygen isotope chronology is assumed to be a proxy of relative humidity, it shows an arid condition during the 1750-1900 period, followed by a humid trend continuing up to present.

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-20

Room:303

Time:May 25 11:30-11:45

Fluctuation of glaciers and glacier runoff in Ili Basin during the last millennium

Akiko Sakai^{1*}, Chiyuki Narama², Nozomu Takeuchi³, Koji Fujita¹, Jumpei Kubota²

¹Nagoya University, ²RIHN, ³Chiba University

Ili River Basin is located at the Central Asia across from northwest China to Kazakhstan. A lot of mountain glaciers distributes at the northern Tien Shan Mountains in the Ili Basin. Meltwater from those glaciers flow through the Ili River and pour into the Balkhash Lake. Total glacier area attains only 0.7% of the total area of the basin. But, meltwater from glaciers is significant water resources because precipitation tends to increase with altitude. Furthermore, glacier can supply water during the summer dry season.

Hence, fluctuation of discharge from glaciers should have affect on the human activity in this basin. In this study, we have tried to reconstruct the fluctuations of glaciers and discharge from glaciers during the last millennium using proxy data such as tree ring and ice core record.

We will compare the calculated fluctuation of glacier area with glacier expansion record indicated by moraine dating in this presentation.

Keywords: glacier, discharge, proxy, tree ring, ice core

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-21

Room:303

Time:May 25 11:45-12:00

Water-level changes of Central Asian lakes during the last 1000 years based on historical maps

Chiyyuki Narama^{1*}, Kicengge¹, Jumpei Kubota¹

¹RIHN

To clarify the environmental changes in Central Asia during the past few hundred years, lake levels of three lakes, Aral Sea, Lake Issyk-Kul (Ysyk-kol), and Lake Balkhash in Central Asia were reconstructed by several historical maps and SRTM DEM data. These historical maps from the 17th-19th century were described in English, Russian, Mongolian, Manchu, Tibetan, Turkic, and Chinese languages during the Qing Dynasty for acquiring information about landscape and land-use of local people) in western countries. The maps show that water level of Aral Sea increased ~2m in the mid-19th from 1960s before lake shrinkage. Water level of Lake Issyk-Kul was 14m higher than at the present during the 17th-mid-19th century, because lake water had overflowed to Chu river. Water level of Lake Balkhash increased 2-3m from the present during the 17th-mid-19th century. The maps show three lakes, which have the same water sources in the Pamirs and Tien Shan mountains, expanded in the 17th-19th century on the same timing. As the previous studies, Aral Sea experienced the drastic decline of the lake-level in the 12th-14th century, and the old settlements around Lake Issyk-Kul in the 10th-12th century have sunk under present lake level. According to several proxy data such as summer temperature from tree-rings, snow accumulation from ice-cores, glacier variations, soil development, and historical documents, this lake-level decline occurred in the 12th-14th century under dry period, and the increase of lake-level in the 17th-mid-19th century under the cold and wet conditions of the Little Ice Age. However, drastic decline of the Aral Sea lake-level might be related with water irrigation systems in the 7th-12th century around the Syr Darya and Amu Darya or flow changes of Amu Darya to Uzboi to the Caspian Sea as shown in previous studies. In specially, the water-level decline of Aral Sea was large-scale in the 12-14th century, comparing with the degree of water-level changes in three lakes during the Little Ice Age. These facts highlight the significant environmental changes including human impacts that have occurred in the past millennium in Central Asia.

Keywords: Central Asia, historical map, lake-level change, proxy data

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-22

Room:303

Time:May 25 12:00-12:15

Construction of hydrological model of the Ili River Basin

Takeo Onishi^{1*}, Akiko Sakai², Jumpei Kubota³

¹RBRC, Gifu University, ²Nagoya University, ³Research Inst. Humanity and Nature

The Ili River Basin is stretching over the China and Kazakhstan. It is an internal river that flows into the Balkhash Lake, which is the largest lake in the Central Asia. Its significant discharge comes from the Tian Shan ridge. Thus, elevation and precipitation difference in the basin is very large. Especially, from the middle part of the river to the lower part of the river, precipitation amount is drastically decreased, and there are classified as semi-arid and arid zone. Though traditional living form of this area was combination of nomadic grazing, and agriculture, large-scale developments such as electric power development by a dam construction and agricultural development during the period of the Soviet Union. In addition to this, under the condition of global warming, the amount of water supplied by glacier melting in the Tian Shan Mountain might decrease. Thus, to evaluate these anthropogenic impacts on hydrological cycle of the basin, we constructed a hydrological model of the basin.

Keywords: Ili, Balkhash lake, Kapchagai reservoir, irrigation

HQR023-23

Room:303

Time:May 25 12:15-12:30

Climate change and historical interactions between human activities and the environment in Central Eurasia

Jumpei Kubota^{1*}

¹Research Inst. for Humanity and Nature

In this paper, a multi-disciplinary research project for understanding historical interactions between humans and the natural environment in arid to semi-arid Regions in Central Eurasia will be outlined.

Historically and geographically, Central Eurasia has been a key area of interaction, transit and exchange between East and West. While many Central Eurasian peoples are well recognized in historical records as skilled nomads, merchants and traders, it is more recently acknowledged that these peoples also assimilated the ideas and artifacts passing through their territories into their own cultures, often with material effect on landscapes and livelihoods. At the same time, Central Eurasia is an excellent location for tracing human reactions to both past climate changes and anthropogenic activities. In this climatically sensitive area, which alternates between semi-arid and arid conditions, human influence can be historically traced. The area with extended arid and semiarid deserts has potential agricultural plains along rivers, flowing from high mountains with many glaciers, which were actively cultivated far back in historical time. These border regions could record both natural environmental and anthropogenic changes very sensitively.

Although interactions between environmental changes and human reactions have rarely been studied in Central Eurasia, agricultural development in the Aral Sea basin has caused the severe lake-level regression that started in the 1960's. Recent agricultural development in arid to semi-arid regions, especially in the latter half of the 20th century associated with modern irrigation technology, has contributed to increasing agricultural production. However, considerable environmental issues have resulted. It is important, therefore, to balance resource development and preservation in arid and semi-arid regions.

This project aims to study and clarify the historical interaction between human activities and natural systems in the semi-arid region of Central Eurasia. The project attempts to clarify historical changes, the rise and fall of nomadic groups and countries, their removal, changes in subsistence, the use of natural resources, and climate change through the analysis of historical documents and archaeological investigations as well as various natural proxies such as ice cores, lake sediment samples, tree rings and wind-blown deposits. At the same time, we will investigate the present status of the area and the effects of human activities on the natural environment, with particular emphasis on their social, religious and cultural background.

Keywords: human societies, environment, historical interaction, arid regions, climate change

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-24

Room:303

Time:May 25 12:30-12:45

Diachronic dynamics of human-environment interactions in Eurasia

Jumpei Kubota¹, Kunihiro Endo², Toshihiko Sugai^{3*}

¹Research Inst. for Humanity and Nature, ²Nihon University, ³University of Tokyo

General discussion and prospects of the Diachronic dynamics of human-environment interactions in Eurasia

Keywords: human-environment interactions, Eurasia, Quaternary, paleoclimate, paleoecology, natural environment

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P01

Room:Convention Hall

Time:May 24 14:00-16:30

Holocene sediment of cores from Lake Rara, western Nepal

Makoto Okamura^{1*}, Hiromi Matsuoka¹, Makoto Osaki¹, Hideaki Maemoku², Hiroshi YAGI³, Atsunori Nakamura⁴, Yusuke Yokoyama⁴, Hirofumi Teramura⁵, Toshiki Osada⁵

¹Kochi Univ, ²Hiroshima Univ, ³Yamagata Univ, ⁴University of Tokyo, ⁵Research Institute for Humanity and Nature

Five cores are collected from Lake Rara basin and terrace using piston coring system. These cores are consisted of light gray colored mud and sandy turbidite layers. Sandy turbidites are intercalated periodically among the cores which cover 7500 years by 14C dating. Constant sedimentation rate of cores shows 50cm/kyr among these 7500 years interval. The average occurrence rate of the turbidites have 800 years interval.

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P02

Room:Convention Hall

Time:May 24 14:00-16:30

The Holocene lake level changes at Lake Qarun and Lake Hamura, Egypt

Kaoru Kashima^{1*}, Ryutaro Naruhashi²

¹Faculty of Sciences, Kyushu university, ²University of Tokyo

The joint research project for the inland saline lakes has started with Kyushu university, University of Tokyo and Menoufia University in Egypt. The project is a multidisciplinary research program, focused on the environmental history of the Neolithic and later periods in this region. More in detail, our research is aimed at a high resolution environmental reconstruction in order to make clear the climatic development during the Holocene, and its impact on human societies. We undertook field surveys at inland lakes and Lake Qarun at Fayoum Basin and Lake Hamura at Wadi Natrun in northern Egypt.

The warming and very dry climate could be observed in the data from all our drillings in the early Holocene in Egypt. The three times of fluctuation between arid ? humid environment have occurred after then. The cyclic fluctuation of humidity was also observed in Lake Qarun in Egypt. The lake level of the lake was fell down after the 2200 years BP in the Ptolemaios period , and fluctuated its level with hundreds years intervals.

The above-noted recovery of humidity and its regional variability in the Holocene presumably was one of the major causes for the archaeological events, especially with regard to water availability.

Keywords: Inland Saline Lake, Lake Level Changes, Holocene, Egypt, Diatom, River Nile

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P03

Room:Convention Hall

Time:May 24 14:00-16:30

Interactions between human activity and natural environment in arid and semi-arid regions in Central Eurasia

Mitsuko Watanabe^{1*}

¹RIHN

The purpose of this paper is to propose the new perspective for the studies on the historical changes of the interaction between human and natural environment in the Central Eurasia.

The Central Eurasia belongs to the arid and semi-arid regions. The ecological environment of this region is characterized by the uneven distribution of the water resources and potential land productivity. When discussing about the historical changes of the interaction between human activity and natural environment in the Central Eurasia, we should give careful consideration to those uneven distribution.

During the 20th century, Central Eurasian people experienced drastic change in their surroundings. Agricultural and industrial development in the Soviet Union had a particularly great influence on the way of use of the natural resources, such as water and land. In this paper, we try to clarify the actual situations of the agricultural development during the Soviet era, in the Ili river basin in the Almaty oblast, Republic of Kazakhstan.

The methods employed in this study included document analysis and interviews. At first, through document analysis, the outline of the formation process and historical changes of the kolkhoz and sovkhos organization are described. Second, that outline is given body and substances through the landscape analysis based on the interview.

Finally, we examine to understand the interaction between human activity and natural environment in arid and semi-arid regions in Central Eurasia, from the view point of the 'possibilism'.

Keywords: Central Eurasia, arid and semi-arid regions, human activity, landscape, environmental history

HQR023-P04

Room:Convention Hall

Time:May 24 14:00-16:30

Middle Holocene abrupt water-level drop of Lake Balkhash revealed by mineralogical analysis of the lake sediments

Hiroki Montani^{1*}, Toshihiko Sugai¹, Tsuyoshi Haraguchi², Kunihiko Endo³

¹Natural environment, Univ. of Tokyo., ²Geosci.,Osaka City Univ., ³Geosystem Sci.,Nihon Univ.

Lake Balkhash located in Kazakhstan in the dry area is a terminal lake of the maximum waters area in Central Asia. It is important to evaluate climatic change and the human activity to restore the environment of old Lake Balkhash.

The paleoenvironment of Lake Balkhash of about 10000 years is being restored because through core analysis and acoustic profiling analysis in the Lake Balkhash (Endo et al.,2010).

However, there is only limited study related to mineralogy. In the restoration of the paleoenvironment around the lake, it is important to understand the mineral deposit in Lake Balkhash. In addition, it will be able to supplement the place without microfossils such as pollens and diatoms in mineral analysis.

Powdery diffractometry was used in the present study. Up to now, the mineral analysis has been done to 80 samples (0901 core 40 samples and 0902 core 40 samples) in total for the crushed sample.

Present result showed the presence of quartz, feldspars, calcites, aragonites, dolomites, micas, and chlorites were contained in 0901 cores. It was also confirmed that 0902 cores contained hydro-magnesite, magnesite, and gypsum present in addition to all minerals of 0901 cores. In the present study, attention is paid to quartz, feldspars in 0901 cores, and magnesite and gypsums in 0902 cores.

Magnesite is generated from the environment with an extremely high Mg/Ca ratio in water (Last,1992). That is, the maximum peak of the magnesite confirmed by 0902 cores suggests the decrease under the evaporation environment in the lake level. In addition, the gypsum confirmed by the same depth is one of the typical evaporites generated with the evaporation environment. It provides an evidence to support the drawdown. 0901 cores also have the coarse-grained sediment. It was confirmed that a lot of quartz and feldspars existed from the mineral analysis in the same depth. In a word, it is suggested that this coarse-grained sediment be a land source material. Moreover, there are white silt in upper and lower 0901 cores and 0902 cores. When this sedimentary facies is compared, the gypsum horizon in 0902 cores is corresponding to the coarse-grained sedimentary facies of 0901 cores. Concerning that the AMS-14C ages of the shell and plant material within the coarse sedimentary unit of 0901 show around 5.5 ka (Sugai et al., 2010), We conclude that the water level of Lake Balkhash markedly decreased in the middle Holocene.

Keywords: Lake Balkhash, X-ray Diffraction, sediments mineralogy, Holocene, lake level fluctuation

HQR023-P05

Room:Convention Hall

Time:May 24 14:00-16:30

Sand dunes formation and development in Ili-River delta and Lake Balkhash area, Kazakhstan

Akio Sato^{1*}, Kunihiro Endo², Reisque Kondo³, Toshihiko Sugai⁴, Hitoshi Shimizu⁴, Yasunori Nakayama²

¹Graduate School of IBS,Nihon University, ²Dept Geosystem Sciences,Nihon University, ³Inst of Geology and Geoinformation,AIST, ⁴GSFS, University of Tokyo

1. Research background

The Subjects of this study is about the relationship between environmental change and sand dune formations in Saryyesik Atyrau Desert, South-East Kazakhstan. Environmental change of the desert is closely related to those in Lake Balkhash and Ili River delta. The 'ILI-PROJECT' members have been studying environmental changes in Lake Balkhash and Ili River delta during Pleistocene to Holocene (Kubota,2010;Endo et al,2010a;Endo et al,2010b). Some researches, acoustic survey and boring in Lake Balkhash (Haraguchi et al, 2010), analysis of lake sediment (Sugai et al, 2010), diatom (Chiba et al, 2010), river terrace sediments and the ¹⁴C dating at Lepsy River (Sugai et al, 2010), analyses for meander wavelength of Ili River and the paleochannel (Shimizu et al, 2010), and ¹⁴C dating about the sediments (Shimizu et al, 2011) have been conducted. Dune formations and patterns show important factors about paleoclimate and paleoenvironment in arid area (e.g. Lancaster, 1995). In case of this study, research for sand dune patterns, stratigraphical relation of sand dunes and fluvial sediments, and age determination of those, are investigated. On the basis of the results we will discuss the lake level change, arid condition about paleo ili River delta, paleoenvironment and paleoclimate in central Asia.

2. Classify of Ili River delta and Sand dune pattern

In this study area, the annual precipitation is about 150mm. Ili River delta are classified into alluvial plain and some paleo-delta plain(T1: high terrace covered with sand dunes, T2 and T3: paleochannels running to north) (Shimizu et al, 2011). In T1 delta, main sand dune pattern are running NW-SE and also NE-SW direction (crossed pattern). These are composed of Draa(mega-linear dune: length:30km over, height:50m over, wavelength:1km over) are developed in the south desert. At the estuary of Aksu River, the southern coast of Lake Balkhash, there are NE-SW linear dunes partially submerged. Sand dunes in T2 paleo-delta covered paleochannel. Small vegetated Sand dunes in T3 paleo-delta before 13th C are developed (Shimizu et al, 2011).

3. Prevailing wind directions at present

Surface wind direction (influenced by the prevailing wind) is principal factors of the sand dunes pattern. The observatory data in wind direction indicate mainly NNE-NE(55%) and SW-SSW(12%) at Balkhash city (Northern coast of Lake Balkhash), NNE-NE-ENE(71%) at Bakanas(center of Ili River delta). In winter season, the temperature and wind direction shows strong cold air supplied from NE-NNE, and the origin of the wind is considered to be 'Siberian high'. We need more researches for understanding about the relationship between present wind direction and sand dunes pattern.

4. Paleoenvironment changes from sand dune studies

The Crossed Draa patterns covering T1 paleo-delta, suggests shift of paleo-wind directions after T1 paleo-delta. The OSL dating result, from the upper part of a linear dune partially submerged at the estuary of Aksu River, shows about AD 200 (Endo et al, 2010a), matching to the reduced phase of Lake Balkhash from studies of the lake sediments. In the lake shrinking phase, dry-up lake bottom might have provided aeolian sand to form the linear dunes. The small dunes in T3 paleo-delta, show that the condition in dune formation has continued after 13th C. But most of sand dunes are 'inactive' or 'vegetated' at present. It suggests that those inactive or vegetated dunes had been formed under the past conditions of higher aridity and stronger wind. The presentation will discuss using more climate data and analysis of geomorphology.

Reference..Chiba et al (2010):Abs.JAQR.40.16-17..Endo et al(2010a):Abs.JGUM. HQR010-12.. Endo et al (2010b):Ili Project, 93-104.. Haraguchi et al (2010):Abs.JGUM.HQR010-10.. Kubota et al(2010): Abs.JGUM.HQR010-09.. N.Lancaster (1995):228-254.. Shimizu et al(2010):Abs.JAQR.40.14-15..Shimizu et al(2011):Abs.AJG.100240.. Sugai et al (2010): Abs.JAQR.40.82-83..

Keywords: sand dunes, Draa, wind direction, Ili-River delta, Lake Balkhash

HQR023-P06

Room:Convention Hall

Time:May 24 14:00-16:30

Optically stimulated luminescence dating of sediments from the Lake Balkhash area, Kazakhstan

Reisuke Kondo^{1*}, Kenichi Uchida², Toshihiko Sugai³, Kunihiko Endo², Takashi Suzuki², Sumiko Tsukamoto⁴, Tatsuhiko Sakamoto⁵

¹Geological Survey of Japan, AIST, ²Nihon University, ³GSFS, University of Tokyo, ⁴Leibnitz Institute, ⁵JAMSTEC

The lake level of Lake Balkhash of Kazakhstan has been fluctuated by the changes in the hydrological and/or climatic environments through the Late Quaternary. It is thought that some landforms have been formed in response to the lake level changes of the Balkhash. However, only a few numerical ages have been obtained from the landforms and sediments from the area, and the lack of the age constraint has prevented studies of precise geomorphic development and terrestrial paleoenvironmental reconstruction. This study applied the optically stimulated luminescence (OSL) dating to fluvial terraces, sand dunes, and gravel bars around Lake Balkhash in order to establish their chronology. Twenty samples for OSL dating were taken from nine localities. The emerged gravel bars are especially important landforms, because they are the direct indicators of higher lake levels than the present Lake Balkhash. Along the lower reach of the River Repsy which flows into southeast shore of Lake Balkhash, fluvial terraces are classified into at least three surfaces (lower, middle, higher terraces). The middle surface occupies the broadest area among these terraces.

Fine grain quartz was extracted from all samples. The OSL measurements of these quartz samples were conducted by using the single aliquot regenerative-dose (SAR) protocol to calculate equivalent dose (D_e) values. The dose rates of the samples were calculated from U, Th and K concentrations which were obtained by neutron activated analysis.

The results of the OSL dating indicate that one of the gravel bars from the lakeshore was formed at 24.5 ka, and the formation of the fluvial middle terrace had occurred at 9.1 ka. Three OSL ages obtained from the lower and the middle fluvial terrace deposits were in agreement with calibrated AMS¹⁴C ages from nearby the OSL samples, which supports that the OSL ages in this study are reliable. In addition, the age of emerged gravel bar (24.5 ka) indicates that the lake level of the Balkhash had been higher during the Last Glacial Maximum. All these chronological data will contribute significantly to the reconstruction of the paleoenvironment in the Lake Balkhash area.

Keywords: Lake Balkhash, gravel bars, fluvial terraces, sand dunes, OSL dating

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P07

Room:Convention Hall

Time:May 24 14:00-16:30

A trial of grasping development of fluvial terraces and hills using OSL dating in Hanno area, Saitama Prefecture

Takashi Suzuki^{1*}, Reisuke Kondo², Kunihiro Endo¹, Tatsuhiko Sakamoto³

¹Nihon University, ²Geological Survey of Japan, AIST, ³JAMSTEC

In Hanno area of Saitama in eastern Japan, fluvial terraces and hills distribute broadly, but detailed dating data and information corresponded with climatic change are only a few. The aim of this study is classification of the river terraces, and establishment of the stratigraphy to elucidate the development of terraces of Late Quaternary in Hanno area. In this area, aeolian deposit known as Kanto Loam, covered terraces and hills thickly. Therefore we applied the OSL dating to aeolian deposit to establish the chronology of Kanto Loam and fluvial landforms. Recently, OSL dating has been expected and applied to elucidate the age of various sediments. We tried to clarify the development of terrace and hill topography using OSL dating in Hanno area, Saitama Prefecture, Japan.

As a result of the trench survey, upper formation of the Tachikawa Loam and fluvial sandy deposits were cropped out. Near the top horizon, Upper Grassy Ash (UG, 12ka) was found out. The result of OSL dating shows linear trend and parallel to an age-depth line determined tephrochronologically. However the difference between two lines is about 8ka. We need to examine the effect of the source and emplacement process of fine-grained quartz, for example volcanic origin, reworking process, etc.

Keywords: Iruma river, fluvial terraces, hills, aeolian deposit, OSL dating, tephra

HQR023-P08

Room:Convention Hall

Time:May 24 14:00-16:30

Exposure age of cirque wall and cirque bottom of Nogaïke Cirque and Komakainoike Cirque: estimates from cosmogenic Be-10

Yasuhide EZURE^{1*}, Yuki Matsushi², Hiroyuki Matsuzaki³, Toshihiko Sugai¹

¹NENV, Univ. of Tokyo, ²DPRI, Univ. of Kyoto, ³NEM, Univ. of Tokyo

This study revealed exposure ages of blocks on cirque wall and cirque bottom of Nogaïke Cirque and Komakainoike Cirque near Mt. Kiso-komagatake in Central Japanese Alps, using TCNs (Terrestrial Cosmogenic Nuclides). TCNs enable us to know how long rock surfaces have been exposed on the ground. Formation age of cirque has been estimated from TCN exposure ages of glacial deposits, but disappearance of glacier is not clear. We measured the concentration of ¹⁰Be and ²⁶Al in rocks, using Micro Analysis Laboratory, Tandem Accelerator (MALT), The Univ. of Tokyo. As a result, the concentration of ¹⁰Be ranged from 10⁴ to 10⁵ atoms g⁻¹ (¹⁰Be) and ²⁶Al ranged from 10⁵ to 10⁶ atoms g⁻¹. Calculated results showed that the upper part of Nogaïke Cirque (NOG-7B and NOG-8B) exposed 4.1-4.7 ka, the lower part (NOG-2B, NOG-3B and NOG-4B) exposed 2.3-6.0 ka, the lower bedrock of Komakainoike cirque (KC-1) exposed 11.8-12.9 ka, the upper bedrock (KC-2) exposed 6.0-7.6 ka, considering with the production rate of ¹⁰Be is 31.6-34.4 atoms g⁻¹ yr⁻¹ and ²⁶Al is 194.2-210.0 atoms g⁻¹ yr⁻¹). These results imply that the cirque glacier had also existed in both cirques between Neoglaciation and Younger Dryas Period, not only during Last Glacial Period.

Keywords: cirque, Terrestrial Cosmogenic Nuclides (TCN), exposure age, Accelerator Mass Spectrometry (AMS)

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P09

Room:Convention Hall

Time:May 24 14:00-16:30

Sensitivity change of TL signal of quartz extracted from the Lake Biwa1400 m Core sample, Japan.

Masashi Takada^{1*}, Aiko Shimada², Shin Toyoda³, Keiji Takemura⁴, Hidehiro Sohma¹

¹Nara Women's University, ²Japan Electron Optics Laboratory Co. Ltd, ³Okayama University of Science, ⁴Kyoto University

Quartz is one of the most common minerals on the Earth and thermoluminescence (TL) signal of quartz is widely used for Quaternary dating in Earth science and Archeology. TL signal in quartz are also used for purposes other than dating. For example, variety of sensitivity change of TL signal of quartz is recorded for quartz crystals of different origins (Takada, 2010), suggesting the possibility for rough estimates of the provenance of sediments. In this study we analyzed TL sensitivity change of quartz grains from the Lake Biwa 1400 meter Core sample, to discuss their chronological background.

(Reference)

Takada M. (2010): Characteristics of 110 degrees Celsius TL signal in quartz from a variety of rocks and sediments: a clue to sediment provenance. *Studies in Geography and Regional Environment Research*, Nara Women's University, VII, 105-112.

Keywords: the Lake Biwa1400 m Core sample, quartz, thermoluminescence signal, sensitivity change

HQR023-P10

Room:Convention Hall

Time:May 24 14:00-16:30

DNA analysis of a single *Pinus* pollen grain in Belukha glacier, Altai Mountains, Russia

Fumio Nakazawa^{1*}, Jun Uetake¹, Yoshihisa Suyama², Ryou Kaneko³, Nozomu Takeuchi⁴, Koji Fujita⁵, Hideaki Motoyama¹, Hiroshi Kanda¹

¹National Institute of Polar Research, ²Tohoku University, ³University of Tokyo, ⁴Chiba University, ⁵Nagoya University

Pollen is a regular component in mid- and low- latitude glaciers since most of them are located within a few ten km from vegetation. In addition, many of the pollen grains contain protoplasm, this suggests that DNA information of the pollen preserved for long periods can be obtained. Modern pollen analysis is used to classify pollen taxon from pollen morphology. Thus, the classification of related species based on the morphology is difficult and limited primarily to plant genus or family. Obtaining DNA information from each single pollen grain in glaciers should allow classification below genus level. It is therefore extremely useful for reconstructions of past vegetation, climate and environment in ice core study. However, no studies have ever tried the detailed classification by obtaining DNA sequence from pollen in glaciers. This study attempted to analyze DNA of a single *Pinus* pollen grain extracted from surface snow taken in Belukha glacier of Russia's Altai Mountains in July 2003. A 149 bp *rpoB* fragment from the chloroplast genome in each *Pinus* pollen grain was amplified using polymerase chain reaction and the DNA products were sequenced in order to classify them at section level. *Pinus* is recognized with approximately 111 species in two subgenera, four sections, and 17 subsections. From the obtained sequences, the pollen grains were classified as sect. *Quinquefoliae* and sect. *Pinus* within four sections namely *Quinquifoliae*, *Parrya*, *Trifoliae* and *Pinus*. *Pinus sibirica* in sect. *Quinquefoliae* and *P. sylvestris* in sect. *Pinus* are currently distributed on the surrounding of the glacier. The consistent results for the section may suggest that the source of the pollen in the glacier is *Pinus* trees found on its surroundings.

Keywords: glacier, pollen analysis, DNA, ice core

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P11

Room:Convention Hall

Time:May 24 14:00-16:30

Last glacial vegetation in Kamishiro, north Nagano, Japan

Arata Momohara^{1*}, Bai Chaobudu¹, Nao MIYAKE², Yoshihiko KARIYA³, Susumu Okitsu¹

¹Chiba University, ²Kochi University, ³Sensyu University

Plant fossil assemblages in last glacial stage between 35,240yBP and 29,630yBP from Kamishiro, north Nagano, central Japan were studied. Plant macrofossil assemblages from 5 horizons include *Larix kaempferi*, *Picea jezoensis*, *Picea* sect. *Picea*, *Tsuga*, *Abies veitchii*, *Pinus* subgen. *Haploxylon*, *Betula ermanii*, *B. platyphylla*, and *Alnus hirsuta* with herbaceous plants as *Carex*, *Chrysosplenium kamtschaticum*, and *Stellaria alsine* var. *undulata*. In pollen assemblages, *Pinus* subgen. *Haploxylon*, *Abies*, *Picea*, *Tsuga diversifolia* type and *Betula* were dominant. The composition represents typical composition in LGM in central and northeast Honshu. Occurrence of seeds of *Chrysosplenium kamtschaticum* with pollen of *Abies homolepis* type indicates that the refugia of temperate plants had been distributed in mesic places in subarctic coniferous forest under cold and dry climate in last glacial maximum inland Honshu.

Keywords: plant macrofossil, last glacial, refugia, pollen analysis

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P12

Room:Convention Hall

Time:May 24 14:00-16:30

Environment and paleobiology based on diatom analysis in Kanto plain during Jomon Transgression

Marie Noguchi^{1*}, Kaoru Kashima², Kunihiro Endo³

¹Nihon University, ²Kyushu University, ³Geosystem Sci., Nihon Univ.

In the Kanto plain, sea water entered into the area of Tokyo Lowland about 10,000 years ago. Around the maximum stage of the Jomon Transgression the coastline was 60km inner than the present. On the basis of diatom analysis, environmental change was investigated mainly using cores in Misato city. This area around Misato at about 9,000 cal.y.BP were tidal to shallow inner bay, then changing to wide inner bay, Oku-Tokyo bay, corresponds to Jomon transgression. About 8,500 cal.y.BP marine plankton species of *Paralia sulcata* was dominant to show from outer bay to inner bay. From 8,500 to 7,500 cal.y.BP, *Crassostrea gigas* of Oyster reef formed, showing wider inner bay. From 7,500 to 5,500 y.BP, Oku-Tokyo Bay was in the maximum stage, and from 5,300 y.BP the regression stage started (Endo *et al.*,1983;Kosugi,1992). Around 4,000 ~ 3,500 cal.y.BP, the coastline situated near Misato, and tidal flat distributed.

Keywords: Jomon Transgression, Diatom, Oku-Tokyo bay

HQR023-P13

Room:Convention Hall

Time:May 24 14:00-16:30

Holocene relative Sea-level Changes in Sekiyado lowland, Japan

Takashi Chiba^{1*}, Shigeo SUGIHARA², Yoshiaki Matsushima³, Kazuo Masubuchi⁴, Yusuke Arai²

¹Grad. School of Frontier Sci., The Univ., ²Meiji Univ., ³Kanagawa Pre. Museum of Natural History, ⁴Kawasakishi board of education

Paleo Kinu bay has formed on lowlands located along the middle to lower reaches Tone river during the Holocene transgression (called the Jomon transgression) period (Endo et al., 1983). Moreover, in Holocene, many shell mounds were made by Jomon and Yayoi people in coastal areas of this bay. So far Kashima lowland (Kikuchi, 1969; 1988), Takagami lowland (Ota et al., 1985; Kashima et al., 1985), Kinosaki lowland (Sugihara et al., 1997; 2000) and lake Kasumigaura (Saito et al., 1990) were surveyed for paleo environmental changes by the Jomon transgression. However, timings and details of paleo sea level changes by the Jomon transgression is not emerged in paleo Kinu bay during Holocene.

In order to reveal the timings and details of sea level changes in paleo Kinu bay during Holocene, we have basically analyzed 4 cores, and drawn Age-depth and sea level curve. The results are as follows;

Stage 1 (about 12000-7000 cal yrBP) : the Jomon transgression occurred and the sea level rose from -25m to +2.5m (Masubuchi 2010). In this period, however, two small regressions (about 8500 cal yrBP and 7500 cal yr BP) are recognized.

Stage 2 (7000-4000 cal yrBP) : sea level fell from +3m to 0m gradually.

Stage 3 (4000-2500 cal yrBP) : sea level fell and reached its the lowest stand (about -2m) during 3000-2500 cal yr BP.

Stage 4 (2500 cal yrBP-) : sea level rose from -2m to 0m

These three regression events (about 8500, 7500, and 3000-2500 cal yrBP) consistent with the colder stage in 8200 cal yrBP, the colder stage in 7400cal yr BP, and the Yayoi regression, respectively (Koizumi and Sakamoto, 2010). In addition, the tide range in this bay is estimated at least a range of plus/minus 4m from diatom analysis during Holocene.

Keywords: Relative sea level change, Holocene, Paleo kinu bay, Sekiyado lowland, Diatom analysis

HQR023-P14

Room:Convention Hall

Time:May 24 14:00-16:30

Variation in magnetic susceptibility of latest Pleistocene to Holocene succession at Kiso river delta, central Japan

Takashi Ogami^{1*}, Toshihiko Sugai²

¹Chuo University, ²University of Tokyo

Applicability of magnetic susceptibility for shallow marine and fluvial sediments is examined based on analysis of latest Pleistocene to Holocene succession. We measured thousands of magnetic susceptibility and grain size distributions for seven drilling cores obtained at Kiso river delta, central Japan.

The cores are drilled at fluvial and delta plains. Deltaic succession contained in the cores is classified into following sedimentary facies, in ascending order: A, braided river; B, flood plain, tidal river and tidal flat; C, bay floor and prodelta; D, delta-front slope and delta-front platform; E, delta plain.

Magnetic susceptibility is measured at 5 cm depth interval using portable magnetic susceptibility meter (SM30, ZH Instruments). Grain size of facies B, C, D and E was measured using laser diffraction particle size analyzer (SALD3000S, Shimadzu Corp.). Detailed age-depth curves of these cores are reconstructed based on densely measured radiocarbon dating.

Profile of magnetic susceptibility change indicates that magnetic susceptibility roughly changes according to sedimentary facies though magnitude of the susceptibility differs among cores. The susceptibility is high at facies B and D, moderate at facies E and low at facies C. Such change appears to be controlled by grain size distributions. It is also recognized that there are lags in timing between change of grain size and magnetic susceptibility.

To evaluate effect of grain size control on magnetic susceptibility, we classified sediments into several clusters by grain size distribution. Ward's method is chosen for clustering operation. Distribution of magnetic susceptibility of each cluster was examined and compared. The median values of magnetic susceptibility increase as median values of grain size increase in case median values of grain size smaller than 0.25 mm (2 phi). In case median values of grain size bigger than 0.25 mm, the values of susceptibility decrease as that of grain size increase.

Variation in magnetic susceptibility among cores and core depth remain in each grain size cluster. These extracted differences should be results of sedimentary processes and backland settings.

Keywords: magnetic susceptibility, grain size, Holocene, Kiso river delta

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P15

Room:Convention Hall

Time:May 24 14:00-16:30

Landform development in Tedori river basin and Togatani limestone cave

Yuki Sakamoto^{1*}

¹Faculty of Letters,Kanazawa University

*

Keywords: Tedori river, Limestone cave, Landform development, Deepening

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P16

Room:Convention Hall

Time:May 24 14:00-16:30

Marine terrace distribution and tectonic geomorphology around the source region of the Rumoi Nanbu earthquake in 2004

Toshinori Sasaki^{1*}, Keiichi Ueta¹, Yasuhira Aoyagi¹, Daiei Inoue¹, Takenobu Tanaka², Masaru Sato², Makoto Yanagida², Norihisa Goto², Hideki Amano³

¹CRIEPI, ²Hanshinconsultants Co. Ltd., ³Hokudensogosekai Co.

To reveal the relation between active fault and earthquake source fault, we have investigated geomorphological and geological survey around the source region of the Rumoi Nanbu earthquake in 2004. In this presentation, we provide the results of high resolution topographic investigation by LiDAR DEM and air photo, and we conclude activity on active fault of this region by boring survey. As a result of topographic survey, we found consecutive tectonic topography which is reverse scarplets on the marine terraces along the coastal line in the 25 km long in N-S direction. The boring survey reveal that the active fault which is formed the reverse scarplets displaces the wetland deposit in the last glacial age.

Keywords: 2004 Rumoi-nanbu earthquake, marine terrace, tectonic geomorphology, DEM investigation, aerial photo investigation, active fault