

SMP046-01

Room:201B

Time:May 27 08:30-08:45

## The CPO and deformation processes of K-feldspar in the Kawai mylonites from the Ryoke metamorphic belt, SW Japan

Kazuhiko Ishii<sup>1\*</sup>, Tatsuya Sakabe<sup>1</sup>, Kyuichi Kanagawa<sup>2</sup>

<sup>1</sup>Osaka Prefecture Univ., <sup>2</sup>Chiba Univ.

The temperature of brittle-plastic transition for K-feldspar has been considered to be 500-550 degrees. Under higher temperature condition, K-feldspar is deformed by dislocation creep with development of crystallographic preferred orientation (CPO). Some granitic rocks deformed under lower temperature, upper-greenschist to lower-amphibolite facies condition, include bands of K-feldspar fine-grained aggregates. These fine-grained aggregates are considered to be deformed by diffusion or dissolution-precipitation creep based on their microstructures and random crystallographic orientation. However, some granitic mylonites deformed under these condition include fine-grained K-feldspar bands that show clear CPO, requiring to understand more detailed processes. We discuss the formation and deformation processes of fine-grained K-feldspar aggregates in granitic mylonites based on their microstructures and CPO.

Kawai mylonite zone is sinistral mylonite zone developed in the Cretaceous Ryoke metamorphic belt, SW Japan. The studied samples are mylonites deformed under upper-greenschist to lower-amphibolite facies condition and are mostly composed of fine-grained quartz bands and porphyroclasts of plagioclase and K-feldspar.

In moderately deformed mylonites including porphyroclasts more than 30 % of whole volume, fine-grained K-feldspar aggregates develop between and around porphyroclasts. The fine-grained aggregates can be divided into two types based on their microstructures. Most of type 1 aggregates develop between porphyroclasts and are characterized by straight grain boundaries and elongated grain shape. In most case, their long axes are sub-parallel to relative displacement direction of porphyroclasts. Type 2 aggregates show undulose extinction and are characterized by less elongated grain shape with irregular grain boundaries.

Highly deformed mylonites include fine-grained K-feldspar bands that can be divided into type A and B based on the geometrical relationship with K-feldspar porphyroclasts. Type A aggregates are fine-grained bands that include several porphyroclasts and type B aggregates are thin and long tails on porphyroclasts. Type A aggregates show CPOs with the same orientation as the porphyroclasts and their [100] and [010] axes tend to be sub-parallel to XZ plane. The CPOs for most of type B aggregates gradually rotate from the same orientation as the porphyroclast as away from the porphyroclast and their [100] axes tend to be high angle to XZ plane. The axes of the rotation are sub-parallel to Y-axis of mylonites but the sense of the rotation is variable without particular relation with the sinistral shear sense of mylonitic deformation.

The features described above may indicate that fine-grained K-feldspar aggregates were formed by recrystallization along microcracks and kink boundaries within porphyroclasts, inheriting crystallographic orientation of the host grain, and then were deformed by granular flow associated with dissolution-precipitation processes. Further, present samples indicate that clear CPO originated from host grain can survive after large deformation through granular flow.

**Keywords:** Granitic ultramylonite, K-feldspar, CPO, EBSD, fine-grained aggregates

SMP046-02

Room:201B

Time:May 27 08:45-09:00

## Deformation of granitoids in brittle-ductile transition: insights from the Asuke shear zone

Junichi Fukuda<sup>1\*</sup>

<sup>1</sup>Dept. Earth & Space Sci., Osaka Univ.

Under brittle-ductile transition, various kinds of fault rocks such as mylonite, cataclasite and pseudotachylyte are formed. It is important to study these rocks to comprehensively understand the processes of inland earthquake generation. Especially, quartz in fault rocks shows various types of recrystallization processes sensitively in response to experienced temperature, pressure and stress; bulging, subgrain rotation, grain boundary migration, which has been investigated experimentally (Hirth and Tullis, 1992). In natural systems, extrapolations of experimental results often give good results to estimate the above parameters (Jerabek et al., 2007; Stipp and Tullis, 2008). Therefore, investigations of quartz texture would give the processes which are experienced during shear deformation. In addition, the slip systems of quartz, induced by plastic deformation, are indicative of the conditions of shear deformation, and they have been studied from experiments and analyses of natural samples (Reviews in Passchier and Trouw, 1996; Toy et al., 2008). However, a lot of reports for recrystallization processes and slip systems of quartz so far focus on regions where plastic deformation of rocks is dominated. Therefore, studies for mineral textures in brittle-ductile transitions are limited.

Based on the above questions, this study focuses on one of the representative areas of brittle-ductile transition in Japan; the Asuke shear zone, Aichi prefecture. The studied area consists of Inagawa granite as host rock (Kanaori et al., 1991; Sakamaki et al., 2006), and the major mineral constituents in studied rocks are quartz, plagioclase, K-feldspar, amphibole, and biotite. The fault rocks in the Asuke shear zone are basically cataclasite, and partly mylonite on outcrop scale. Under optical microscopy, quartz fine grains less than 1  $\mu\text{m}$  are recognized around large quartz grains up to 3 mm. Plagioclase and K-feldspar are fragmented by brittle deformation, and fine grains in micrometer scales can be recognized. It has been discussed that these grain-size-reduction processes are induced by above dynamic recrystallization (bulging or subgrain rotation in this case) or introductions of clacks (Vernooij et al., 2006a). Moreover, Vernooij et al. (2006b) suggested that dissolution-precipitation process partly contributes to grain size reduction. Also, it has been argued that dauphine twinning may induce dynamic recrystallization of quartz (Lloyd, 2004; Stipp et al., 2008) or not (Neumann, 2000). These grain-size-reduction processes as well as slip systems can be inferred from analyses of crystallographic preferred orientations (CPOs) and misorientation axes. In this study therefore, we use electron backscatter diffraction (EBSD) and measure relationship of crystallographic orientations between quartz fine grains and host large grains. Then, I will discuss the grain-size-reduction processes of quartz in brittle-ductile transition. Also, surfaces of quartz grains are observed by using scanning electron microprobe (SEM). Grain-size-reduction processes and contributions of fluid will be discussed from their morphologies, together with the EBSD results. In addition to analyses for quartz, deformation mechanisms of feldspars are investigated: The compositions of feldspars between fine grains in micrometer scales and porphyroclasts are determined. EBSD analyses and observations of grain morphologies by SEM for feldspar fine grains are performed. Then, I will discuss deformation mechanisms of feldspar fine grains (in this case, grain boundary sliding or possibility of solution-precipitation creep).

Keywords: quartz recrystallization, grain morphology, feldspar deformation, composition, electron backscatter diffraction

SMP046-03

Room:201B

Time:May 27 09:00-09:15

## Elastic anisotropy of deformed rocks

Tohru Watanabe<sup>1\*</sup>, Yuto Shirasugi<sup>1</sup>, Katsuyoshi Michibayashi<sup>2</sup>

<sup>1</sup>University of Toyama, <sup>2</sup>Shizuoka University

The seismic anisotropy will provide us the information about deformation in the Earth's interior. In order to interpret observed anisotropy, we must understand the relationship between deformed textures and elastic properties. Plastic deformation gives rise to the lattice preferred orientation (LPO) of mineral grains, which leads to elastic anisotropy in deformed rocks. The development of SEM-EBSD has enabled us to measure the orientation of individual mineral grains in a deformed rock. Using the measured orientations and elastic constants of single crystals, we can calculate elastic properties of the deformed rock using Voigt or Reuss averaging schemes. No information about the shape or arrangement of grains is used in these averaging schemes. The Voigt average gives an upper bound, and the Reuss average a lower bound to elastic stiffness. The stiffness of the rock sample should be found within these bounds. When component minerals have similar elastic properties and weak anisotropy, Voigt and Reuss averages are nearly equal. These values can provide a good prediction of elastic properties. However, when component minerals have strong anisotropy, Voigt and Reuss values are far apart (Mainprice and Humbert, 1994). Additional information like the shape and arrangement of grains should be taken into account for a better prediction of elastic properties.

As the arrangement of grains, we focus on a layered structure seen in deformed rocks. For simplicity, we consider an alternation of two compositionally different layers. The two layers are composed of different mineral grains, which are well aligned in each layer. Two layers have different effective elastic constants. The elastic properties are assumed to have the symmetry of orthorhombic systems in both layers. Two layers are assumed to have the same principal axes of elastic tensors. The calculated effective elastic constants show Reuss values in relation to deformation perpendicular to the layers. The calculation gives Voigt values in association in relation to deformation parallel to the layers. In this presentation, we will also compare calculated and measured velocities for serpentinite mylonites.

Keywords: elasticity, anisotropy, deformed rock, LPO, layered structure

SMP046-04

Room:201B

Time:May 27 09:15-09:30

## metasomatic instability and constitutional oversaturation

mitsuhiro toriumi<sup>1\*</sup>, fukuyama mayuko<sup>2</sup>

<sup>1</sup>JAMSTEC, <sup>2</sup>Insti.Earth Sciences, Acad.Sinica

Migrating interface derived from interdiffusion of ionic species in permeable solution in the plate boundary rocks occurs as wavy boundary between mono- and bi-mineralic bands. Simple examples are found in the banded basic schists of the Sambagawa metamorphic belt. Albite - quartz - calcite bands having wavy interface are commonly sandwiched by thin chlorite bands and sometimes by thin epidote bands. The trails of hematite grains are continuously pervaded into albite - quartz bands from chlorite band, indicating the advancement of interface into chlorite bands.

The wavelength of the interface increases firstly and then reaches the constant level with width of albite - quartz band. It suggests that the preferable wavelength of the interface develops with time, considering the increasing width of albite - quartz band with time. However, the ratios between width of band and wavelength of the interface varies in different rock specimens.

The modeling of the wavy interface development should be constructed in the system of chemical equilibrium of solution with albite, chlorite, quartz and calcite. At the interface chemical equilibrium attains but being apart from the interface, the diffusion of ionic species relevant with chlorite and other minerals should make change in equilibrium concentrations. As the concentrations gradients of ionic species in grainboundary solution makes constitutional oversaturation (1) of one side mineralogy, the interfacial geometry becomes unstable for small waveform perturbation. The selective wavelength in this case is governed by the ratio of oversaturation degree and interfacial energy. In this study, the authors can propose the capillarity effect constitutional oversaturation instability of the metasomatic banding interface. This is called as metasomatic instability which controlled by velocity of interface advance.

### Reference

(1) W.Kurz and D.J.Fisher, Fundamentals of solidification, Trans Tech Pub., 1986.

Keywords: metasomatic instability, constitutional oversaturation, metamorphic banding

SMP046-05

Room:201B

Time:May 27 09:30-09:45

## Implication of the residual pressure recorded in quartz inclusions in garnet from high T metamorphic terranes

Tadao Nishiyama<sup>1\*</sup>, Kyoko Aikawa<sup>1</sup>

<sup>1</sup>Kumamoto University

Quartz inclusions in garnet from metamorphic terranes reveal residual pressure depending on the metamorphic pressure and temperature of their entrainment (Enami et al, 2007). The residual pressure occurs due to the difference in elastic properties of host garnet and quartz inclusions and also due to the pressure and temperature change according to the exhumation path. The residual pressure can be measured by Raman micro-spectroscopy, and the results are consistent with those estimated by a simple elastic model of the sphere-in-hole problem (Enami et al, 2004). This is an important progress in metamorphic petrology, because it represents a new development of metamorphic barometry independent of thermodynamic methods.

However, several problems have remained to be solved in the application of this method. One problem is the applicability of this method to high temperature metamorphic terranes in which low - high transition can occur in quartz. This study aims to solve this problem and examined quartz inclusions in garnet from the Higo Metamorphic Rocks, one of the high T metamorphic terrane of Cretaceous in age in Central Kyushu.

The studied area is Kosa district, Kumamoto Prefecture, which is the same area studied by Maki et al (2004). Here we adopt the metamorphic zonal mapping after Maki et al (2004): biotite zone, garnet - cordierite zone, and orthopyroxene zone from north to south. This study newly defined the muscovite - out isograd and the tourmaline - out isograd in the studied area. The peak metamorphic temperatures are about 600 °C at the muscovite - out isograd and about 800 °C at the orthopyroxene isograd, which are estimated by the garnet - biotite thermometer and the petrogenetic grid (Maki et al, 2004).

We examined eight samples collected from the area between the muscovite - out isograd and the orthopyroxene isograd by Raman micro-spectroscopy. According to Enami et al (2004), we made plots of  $Dw_1$  vs  $Dw_2$ . In the plots we found two kinds of data: one plotted in the first quadrant (both  $Dw_1$  and  $Dw_2$  are positive) and the other plotted in the third quadrant (both  $Dw_1$  and  $Dw_2$  are negative). The two kinds of data are taken from the same garnet grain. The latter data has not been reported by Enami et al (2004) who studied mostly high pressure metamorphic rocks. We interpret that the data plotted in the first quadrant show compressive stress and the data plotted in the third quadrant do tensile stress as suggested by Enami et al (2004).

The peak metamorphic pressure and temperature condition of this area is likely to have reached the stability field of high quartz (Obata et al, 1994, Osanai et al, 1998, Yoshimura, 2004, Maki et al, 2004 and Miyazaki, 2004). Therefore quartz inclusions showing tensile stress may be originally high quartz when they are entrained in the host garnet. The high quartz will transform into low quartz when the host rock crosses the transition curve during the exhumation, and the volume decrease by about 1 per cent associated with the transition may cause tensile stress in the garnet. Quartz inclusions showing compressive stress from the same garnet grain may have been entrained in garnet when the rock crosses the transition curve from high temperature side in the retrograde path. Therefore we assume that quartz inclusions with compressive stress have been entrained at the low- high transition temperature and evaluated the entrainment pressure by combining the residual stress measured by Raman micro-spectroscopy and the elastic model. The result shows 0.35 - 0.5 GPa, independent of sample location in the area. The estimated pressure is higher by 0.1 - 0.3 GPa than the peak metamorphic pressure estimated by Maki et al (2004) and also by Miyazaki (2004), and lower than the pressure by 0.3 - 0.5 GPa recorded in a calc-silicate granulite in the same area reported by Maki et al (2009).

Keywords: garnet, quartz, residual pressure, high T metamorphic terrane, Raman micro-spectroscopy, elastic interaction

SMP046-06

Room:201B

Time:May 27 09:45-10:00

## Distinct difference of metamorphic texture between high P/T and low P/T metapelites

Kazuhiro Miyazaki<sup>1\*</sup>

<sup>1</sup>GSJ AIST

Localization of deformation and reaction will affect macroscopic evolution of metamorphic belt in the crust. To evaluate localization of deformation and reaction, metamorphic textures of metapelites from high P/T metamorphic belt (Sanbagawa Metamorphic Rocks) and low P/T metamorphic belt (Ryoke Metamorphic Rocks) are examined.

Distinct difference of metamorphic textures between high P/T and low P/T metapelites is observed with EPMA mappings of thin sections as follows. Metamorphic differentiation and coarsening of metamorphic minerals progressed well in high P/T metapelites. Quartz-rich layers or lenses are formed in high P/T metapelites. On the other hand, mosaic textures of quartz-rich and Al-bearing mineral domains are developed in low P/T metapelites. Textures of migmatite of low P/T metamorphic rocks are exceptional and complex. Coarsening of metamorphic plagioclase is clear in high P/T metamorphic rocks. Average size of metamorphic plagioclase in higher-grade high P/T metapelites is about 10 times larger than those of lower-grade high P/T metapelites. At the same metamorphic temperature (around 500 °C), average size of metamorphic plagioclase of high P/T metapelites is about 10 times larger than those of low P/T metapelites. Exceptionally, average size of metamorphic plagioclase in migmatites is much larger than those in the non-migmatitic low P/T metapelites.

It is suggested that quartz-rich layers or lenses in high P/T metapelites were formed by deformation, because these layers or lenses are parallel or subparallel to schistosity. To examine the deformation-associated metamorphic differentiation, deformation of immiscible two-phase viscous fluid is simulated. The results of deformation of the two-phase fluids without viscosity contrast show that coarsening of the each mosaic domain takes place. And multi-layers or elongated mosaic structures are formed by deformation. On the other hand, the results of deformation of the two-phase fluids with viscosity contrast show that branching layered structures or lenses of lower viscosity fluid are formed. These types of structures are developed with strain localizing in low viscosity fluid.

The numerical simulations suggest that formation of quartz-rich layers or lenses in high P/T metapelites is caused by strain localization in quartz-rich domain. On the other hand, coalescence and precipitation-dissolution are important for coarsening of metamorphic minerals with deformation. Because dihedral angle between quartz and water becomes less than 60° at high P/T conditions, fluid network will be formed within quartz-rich layers or lenses, which will enhance reaction-diffusion in high P/T metapelites. The above-mentioned mechanism has a positive feedback, therefore, it is expected that textural evolution due to deformation and reaction in high P/T metapelites will accelerate progress of deformation and reaction.

Keywords: metamorphic rock, metamorphic texture



SMP046-07

Room:201B

Time:May 27 10:00-10:15

## Heterogeneous nucleation, Ostwald step rule and Potts model: Implications for precipitation of silica polymorphs

Atsushi Okamoto<sup>1\*</sup>

<sup>1</sup>Tohoku University

The rock-forming minerals include several polymorphic minerals such as aluminosilicates, serpentine, carbonates, carbon, and silica. These minerals have been used as indicators of P-T condition and/or fluid compositions, when they formed. It is also well known that metastable phases commonly appear especially during fluid-rock interaction, including aragonite in carbonate sinters, and opals in silica sinters. Okamoto et al. (2010) found that the dominant silica minerals precipitated from aqueous solutions in order of amorphous silica, cristobalite to quartz, and that quartz is dominant in the solutions including minor Al and Na. The formation of less stable phase prior to the most stable phase is called as the Ostwald step rule. The phenomena following the Ostwald step rule is explained by relative differences in growth rate among the polymorphic minerals in the presence of seeds of all phases, but nucleation of these seeds should be considered. In this study, heterogeneous nucleation and the effects of impurity are investigated by a simple microscopic model, Potts model, in Monte Carlo simulation.

In the Potts model, each spin (molecule) in a lattice has one of  $q$  states (phases). Each lattice site  $i$  has a spin  $s_i$  taking values in the range of 1 to  $q$ , and the energy of a configuration  $s$  is given by the Hamiltonian,  $H = - [J]_{s_i, s_j} - [h_a M_a]_a$  (eq. 1), where  $M_a$  is the number of spins of the spin type  $a$ . The first term is the sum of nearest-neighbor pairs of spins with the symmetric interaction energy (surface energy), and the second term describes the effect of external fields  $h_a$  acting on spin type  $a$  (bulk free energy). To evolve the system, we employed discrete-time Metropolis dynamics. Here, we consider spins of four states (1-4) in a  $30 \times 30$  square lattice in two dimensions and periodic boundary conditions. We set  $T = 0.80$ , and impose the external fields as  $h_a = 0.1 \times (a - 1)$ , indicating that state 4 is the most stable. When we set the diagonal elements of the interaction matrix to unity and the other elements to zero, any metastable phase does not appear. The sequential nucleation of metastable phases following the Ostwald step rule is realized when we introduce a non-diagonal interaction,  $J_{a, a+/-1} = F1 > 0$  (attractive), between neighboring phases, and all other non-diagonal interactions have  $J_{a, b} = F2 < 0$  (repulsive) (Sanders et al. 2007). The successive nucleation of metastable phases in order of 1, 2, 3 and 4 appears with  $F1 = 0.1$  and  $F2 = -1.0$ . We also find that phase  $i$  nucleated in phase  $i-1$ . This result is qualitatively same to the occurrences of the silica minerals in the pure Si solution of Okamoto et al. (2010): homogeneous nucleation of opal-A from the solution, heterogeneous nucleation of opal-C upon opal-A and that of quartz on opal-C. A new phase preferentially nucleates on the next-less-stable phases, as they are probably more structurally similar to new phase than are other phases.

The effect of impurities was introduced as  $-[J_{pa}]$ , that is the interaction between the impurity,  $p$ , and spin  $a$  into equation 1 following Sear (2005). To consider the case of silica, we propose that an interaction between phase 4 and an impurity equals 1, and that interactions between other states and an impurity are zero. The simulation results reveal that the evolution of the system varies with increasing number of impurities. When the number of impurities is small, nucleation of the most stable phases occurs via the formation of metastable phases. In contrast, with a larger amount of impurity, phase 4 is able to nucleate directly. The direct nucleation of quartz in the solution with small amounts of impurities (Okamoto et al. 2010) indicates the strong interaction between quartz and these cations.

### References:

- Okamoto A. et al. (2010) *Geochim Cosmochim Acta*, 74, 3692-3706.
- Sanders DP. (2007) *Phys Rev B*, 75, 132101.
- Sear PS (2005) *J. Phys. Condens. Matter* 17, 3997-4004.

Keywords: polymorphic minerals, Ostwald step rule, heterogeneous nucleation, Potts model

SMP046-08

Room:201B

Time:May 27 10:15-10:30

## Zircon behavior in the upper amphibolite facies metamorphic rocks from the Aoyama area, Ryoke metamorphic belt, SW Japan

Isao Yamaguchi<sup>1</sup>, Tetsuo Kawakami<sup>1\*</sup>, Yokoyama D. Takaomi<sup>1</sup>, Tomoyuki Shibata<sup>1</sup>, Takafumi Hirata<sup>1</sup>

<sup>1</sup>Kyoto University

Recent researches on zircon behavior showed that zircon can crystallize at various stages of metamorphism. In order to interpret the results of the U-Pb zircon dating, it is essential to understand how zircon behaves and responds to the metamorphic processes (Harley et al. 2007). However, zircon-forming, zircon-consuming and zircon-modifying reactions are controlled by the whole-rock composition, pressure-temperature conditions, and fluid-melt-rock interactions. This complexity leaves zircon behavior in metamorphic rocks still unraveled. This research aims to describe the mode of occurrence of zircon in the upper amphibolite facies metamorphic rocks exposed in the western part of the Aoyama area (west of the Kachiji fault), Ryoke metamorphic belt, SW Japan (Yamaguchi and Kawakami, 2008), and to understand the behavior of zircon during the high-temperature metamorphism.

In the Aoyama area, upper amphibolite facies pelitic schists and pelitic-psammitic migmatites are widely exposed and the metamorphic grade increases from the north to the south (Takahashi & Nishioka, 1994; Kawakami, 2001). The metamorphic conditions of the study area estimated by the geothermobarometry were about 610 °C, 3 kbar in the north and about 715 °C, 6 kbar in the south. Kawakami and Suzuki (2008) reported the CHIME monazite age of the Ao granite that is intruded to the south of the Aoyama area to be 79.8±3.9 Ma.

Eighteen pelitic-psammitic metamorphic rock samples from the study area were examined under SEM-EDS and about 250 grains of zircon were described in detail. As a result, at the north of the study area, zircon grains larger than 20 µm in diameter were abundant than in the south.

In order to confirm whether this trend is controlled by the whole-rock Zr composition or not, the remaining half of the chips that was used for the thin section preparation were powdered and analyzed by the XRF analysis. As a result, most of the whole-rock Zr was resided in the zircon grains larger than 20 µm in the north, whereas zircon grains larger than 20 µm occupied only 20-30 % of the whole-rock Zr in the south.

Based on the result above, a hypothesis is made that most of the coarse-grained zircon in the north larger than 20 µm would be detrital. In contrast, at the south, zircon is likely dissolved and may have newly nucleated during the metamorphism, so that Zr may be resided in zircon smaller than 20 µm and in other minerals.

In order to check this hypothesis, zircon U-Pb dating by LA-ICPMS was carried out. As a result, zircons (> 20 µm) in the north showed the Proterozoic to the Jurassic discordant ages both in the core and the rim. These ages would represent detrital ones. On the other hand, zircon grains in the south showed the Cretaceous discordant ages. This may represent the partial resetting of the old, detrital zircons during the Cretaceous or the Tertiary time. Therefore, it is likely that in the study area, detrital zircon grains including coarse ones dissolved and partially reset during the Ryoke metamorphism and/or during the contact metamorphism by the Ao granite and the Kabuto granodiorite intrusions postdating the Ryoke metamorphism. To confirm that new zircon grains nucleated in the south of the study area during the Ryoke metamorphism or not, the dating of tiny grains and thin overgrowths of zircon and check of the REE patterns of them are important.

**Keywords:** high-temperature metamorphism, zircon, U-Pb dating, laser ablation ICPMS, zirconium, Ryoke metamorphic belt



SMP046-09

Room:201B

Time:May 27 10:45-11:00

## Heterogeneous distribution of garnet in the Sanbagawa metamorphic rocks in Kanto Mountains

Mutsuko Inui<sup>1\*</sup>, Kosuke Hatori<sup>1</sup>, Youji Hanagaki<sup>1</sup>, Ryota Hadachi<sup>1</sup>

<sup>1</sup>Kokushikan University

Occurrence of garnet in the Sanbagawa metamorphic rocks from Nagatoro area, Kanto Mountains, east Japan, was investigated in detail.

Garnet have been known to occur in the Nagatoro-area pelitic rocks, which contributed to the definition of the metamorphic zonation in the area. The occurrence, or texture, of the garnet grains have, however, not been described in detail. Textural observation of garnet is becoming more and more important, since garnet grains with different texture has now been reported from the Sanbagawa metamorphic rocks in Shikoku.

In this study, pelitic samples were taken from the high-grade zone area in the Sanbagawa belt exposed in the Kanto Mountains. Most garnet grains found in the pelitic schists were smaller than 0.1 mm in diameter, and were either included in the albite porphyroblasts (referred as albite-spot, hereafter) or among muscovite grains. The heterogeneous distribution is similar to the type-B garnet found in Shikoku (Inui, 2010). The average size of the albite-spot seemed to be larger in samples with garnet than in those without garnet. The shape of the garnet grains included in the albite-spots were mostly euhedral, whereas about half of the garnet grains within muscovite layers had round shape. Many of the rounded grains had aspect ratios larger than 2. Such grains often accompanied chlorite "tails" at their either end, which suggests that the grains have been resorbed after their euhedral growth. The overall texture suggests that the euhedral garnet grains in albite-spots are perfectly preserved, on the other hand, the long and round grains among muscovites are partly preserved. It is likely that garnet grains formed in the other parts of the rocks are mostly resorbed and are lost. It infers mechanism that resulted in the heterogeneous distribution of the garnet in the schists. The correlation of the size of garnet grains and the distance to its nearest neighbor suggested that the initial garnet growth was controlled by the velocity of material transfer in the rocks.

The heterogeneously distributed garnet grains in the Nagatoro area were compared to the similar garnet in the Asemigawa River area and the origin of the garnet will be discussed.

Inui, M. (2010) Two types of garnet in Sanbagawa pelitic schists along Asemigawa River, central Shikoku *Journal of Mineralogical and Petrological Sciences*, 105, 274-279.

Keywords: garnet, Kanto Mountains, Nagatoro, grain size distribution, heterogeneous distribution

SMP046-10

Room:201B

Time:May 27 11:00-11:15

## High-Mg cores of the garnets in the Sambagawa pelitic schists from the Besshi district, central Shikoku, Japan

Kabir Md Fazle<sup>1\*</sup>, Takasu Akira<sup>1</sup>

<sup>1</sup>Shimane University

Pelitic schists adjacent to the Seba metagabbro consist mainly of garnet, phengite, epidote and quartz with small amounts of amphiboles (Na-Ca and Ca-amphibole), albite, chlorite, biotite and carbonaceous matter. Rutile, titanite, ilmenite, calcite, paragonite, hematite and zircon are occasionally present as accessory minerals. A schistosity is defined by preferred orientation of coarse-grained phengite (3.5 mm).

Garnets in the pelitic schists occur as euhedral to subhedral grains up to 3 mm across. The garnets are optically zoned, from pale red-colored cores to colorless rims. The garnets have almandine-rich composition ( $X_{Alm}=0.50-0.60$ ) with variable amounts of the grossular ( $X_{Grs}=0.18-0.31$ ) and pyrope ( $X_{Prp}=0.08-0.24$ ) components. Two zones (almandine-pyrope rich core and almandine rich rim) were identified based on chemical composition. The cores are abundance of rich in pyrope, decreasing toward the core-rim boundary ( $X_{Prp}=0.24-0.18$ ) and they show antithetic zoning of  $X_{Alm}$  (0.50-0.56). The boundary between the core and the rim shows a sharp chemical discontinuity. In the rim,  $X_{Alm}$  (0.56-0.58) and  $X_{Prp}$  (0.09-0.13) increase,  $X_{Sps}$  (0.02-0.01) decreases, and  $X_{Grs}$  (0.29-0.31-0.28) increases and then decreases slightly. The garnets contain inclusions of epidote, Ca-amphibole (Mg-hornblende), phengite ( $Si=6.53-7.27$  pfu), paragonite, albite (An 0-3), chlorite, calcite, ilmenite, rutile, titanite and zircon. The garnets also contain polyphase inclusions of Mg-hornblende+paragonite+quartz; epidote+paragonite; epidote+paragonite+quartz and epidote+albite. Garnets are partly replaced by aggregates of amphibole (Mg-hornblende), phengite ( $Si=6.47-6.52$  pfu), epidote and albite (An 0-3) and symplectites of Ca-amphibole (Mg-hornblende) and albite (An 0-5) with rare quartz. The margins of the garnets are occasionally replaced by chlorite, phengite and biotite, or by biotite and calcite aggregates.

Amphiboles occur in the matrix as subhedral to anhedral prismatic grains up to 2 mm long. Some amphiboles in the matrix are zoned from winchite, barroisite cores to Mg-hornblende rims.

According to the petrography and chemical compositions of the minerals the pelitic schists are adjacent to the Seba metagabbro mass probably suffered high-pressure metamorphic condition. The chemical discontinuity between the core and the rim of the garnets is formed in the different metamorphic conditions for the core and the rim. Similar zoning of garnets have been reported by Higashino and Takasu (1982), Takasu (1984) and Nomizo (1992) from the Besshi district. Takasu (1984) reported similar garnet from the Seba metagabbro mass, the core of garnet undergone an eclogite facies metamorphism at higher temperature conditions (720-750 C, 12-24 kbar) and the rim together with the surrounding Seba eclogitic basic schists (610-650 C, 7-24 kbar).

### Reference:

Higashino, T and Takasu, A., 1982, Journal of Japan Association of Mineralogy, Petrology and Economic Geology. 77, 362-367. Takasu, A, 1984, Journal of Petrology. 25, 619-643. Nomizo, A, 1992, Journal of Geological Society of Japan. 98, 49-52.

\*Corresponding author: fazlekabir@gmail.com

Keywords: pelitic schist, high-Mg garnet, Sebadani, Sambagawa

SMP046-11

Room:201B

Time:May 27 11:15-11:30

## Metamorphism and thermal structure of subduction zone: a case study on the Sanbagawa pelitic rock

Soichi Omori<sup>1\*</sup>, Kazumasa Aoki<sup>2</sup>

<sup>1</sup>Earth & Planetary Sciences, Tokyo Tech., <sup>2</sup>The University of Tokyo

The thermal structure of the present subduction zones has been a focus of geophysical studies regarding the origin of arc magma, transportation of volatile component such as H<sub>2</sub>O and CO<sub>2</sub> into the deep mantle, and origin of the subduction zone earthquakes. Many studies employing numerical modeling have been done for the aim estimating the thermal structure of the subduction zone. The results contributed semi-quantitative understanding of the thermal structure of the subduction zone, however, diversity of the results among the models is not negligible to be applied to the natural system.

It has also been one of the motivations of metamorphic geology that analysis of prograde metamorphism of a suite of metamorphic rocks in a metamorphic belt yields information of the subduction geotherm. Recent advance in decoding metamorphic P-T condition enabled revealing true peak P-T condition and prograde P-T path excluding extensive retrograde hydration. Therefore, now, we can discuss the subduction geotherm of past subduction zone in a context of metamorphic geology.

In this presentation, we review nature of the thermal structure in the subduction zone, and prograde- and progressive-metamorphism, at first. Then a case study on the Sanbagawa pelitic rock is demonstrated with implications to the mantle-wedge dynamics and the material transportation.

Keywords: subduction zone, thermal structure, metamorphic rock, exhumation

SMP046-12

Room:201B

Time:May 27 11:30-11:45

## Wedge extrusion followed by major out of sequence thrusting accompanying duplexing, the high P/T Sambagawa blue schist


Soichi Osozawa<sup>1\*</sup>

<sup>1</sup>Tohoku University

I am seeking for a sponsor, like museum, city or town office, who kindly publishes the colored geological map, very wide, my studied area. Map file is pc based, and only 200,000 Yen for 500 copies is needed.

The Sambagawa high P/T zone of Besshi area is regionally mapped, and characteristic major structures performed the exhumation of high grade metamorphic rock is clarified. As noted by Osozawa and Pavlis (2007), the most fundamental structure is a D2 extrusional wedge. The southeastward wedge consists of a series of normal faults domain at hanging wall of the Asemigawa detachment, and a series of thrust faults domain of the footwall. These faults bound every metamorphic mineral zone, and the thermal culmination of the Asemigawa section is oligoclase-biotite zone, hanging wall of the detachment fault. To the NW, we now confirm that the culmination is an eclogite body, constituting an axis of the extrusional wedge. To the SE, these extrusional wedge and a series of normal and thrust faults are linearly traceable on regional map, but suddenly not continuous to the chlorite zone and disrupted. The disruption is due to the Hamegano out-of-sequence thrust, newly found in this paper, and the Besshi unit is divided into two units. The OST divides the chlorite zone the upper L tectonite consisting of varicolored mafic schist, and the other lower pelitic schist. D2 folds at the hanging wall is disjunctive and broken by the movement of brittle OST. The OST is clearly traceable to the NW from the Asemigawa area. To the NE, a syncline is observed as mapped by previous studies, but it exists only on hanging wall, and the structure is discordance with not-folded or obscured footwall chlorite zone. Its SE limb, hanging wall of the OST, is extrusional wedge, but the NW limb consists of duplexing metamorphic mineral zones, and the syncline is lithologically asymmetric. Further to the NE, OST and hanging wall duplex is involved in NW plunging anticline. We named the duplex stack formed near antichinal axis of OST, the Tomisato duplex. OST and duplex moved SW, as a final expression of metamorphic exhumation, following the wedge extrusion. The Median Tectonic line might have played an important role for exhumation as a root, but most of the Besshi unit, including eclogite body, is consequently rootless. Interestingly, the chlorite zone rocks hanging wall of the OST include a psammitic schist, only at the NW limb of anticline. The schist is lithologically similar to that of the Oboke unit, the structurally lowest unit. The Oboke unit and thrust is also concordantly folded with the OST and duplex, for the antichinal axis. Another anticline is to the SW, Nakashichiban area, where the psammitic schist of Oboke unit is once again exposed.

Metamorphosed ultramafic rocks, including eclogite, are distributed in every metamorphic zone. The metamorphic minerals constitute D1 foliation overprinted by D2 deformation. Therefore, ultramafic rocks were amalgamated with surrounding rocks before D1. Two possibilities of mixing mechanism are expected; sedimentary or tectonic melange process, and we prefer in this case tectonic slicing during D0 at deeper subduction zone, by considering eclogite metamorphism.



Keywords: Sambagawa high P/T zone, extrusional wedge, out-of-sequence thrust, duplex, exhumation, D2

SMP046-13

Room:201B

Time:May 27 11:45-12:00

## Tectonics of the Kamuikotan metamorphic rocks distributed in the western part of Asahikawa-city, Hokkaido, Japan

Ayumi Okamoto<sup>1\*</sup>, Toru Takeshita<sup>1</sup>

<sup>1</sup>hokkaido University

We have studied the tectonics of the Kamuikotan metamorphic rocks, central Hokkaido, which were developed in the Sorati-Yezo Belt convergent boundary between the Eurasian continent and the subducting oceanic plate in the Cretaceous. The Kamuikotan metamorphic rocks are known as typical high-P/T type metamorphic rocks, where P is pressure, T is temperature, respectively. In this study, we investigated the western part of the Kamuikotan gorge area, the western part of Asahikawa-city. The rocks in this area were characterized by a peculiar P-T path and mineral assemblages different from any other metamorphic rocks in the Kamuikotan metamorphic belt. First, we analyzed the tectonics of the Kamuikotan metamorphic rocks from geological and petrological approach (i.e., field work, chemical composition analyses of both minerals and whole rocks). Based on field work, it has been found that the protolith stratigraphy in this area is composed of a typical accretionary stratigraphy (basaltic rocks, limestone, chert and mudstone in ascending order), which was repeated by thrust. Further, three deformation stages (D1, D2 and D3) have been revealed by the overlapping relationships among folds. D1 is characterized by main schistosity (S1), D2 is characterized by east-vergent close folding (F2), and D3 is characterized by crenulation cleavages (S3). Next, we analyzed the P-T path from blueschist to greenschist facies, and further to the surface of the earth for a mafic schist that includes Na-amphibole surrounded by actinolite. We constrained the range of P and T by the pseudosection method. As a result, it has been found that the P-T range lies on the reaction line between pumpellyite and epidote on condition that pumpellyite, epidote and albite coexist. (P is 3.6-4.5 kbar, T is 280-290°C or P is 3.4-4.7 kbar, T is 275-300°C. This difference in P and T is from amphibole model.) Further, we estimated the formation temperature of epidote in equilibrium with pumpellyite from the pistacite component (0.27-0.31) (Nakajima et al., 1976), which yielded the temperature range of 296-310°C. In the same way, we estimated the formation temperature of chlorite from the chemical composition (Inoue et al., 2009), which yielded the temperature range of 121-240°C if all Fe is Fe<sup>2+</sup>, and that of 109-220°C if some Fe is Fe<sup>3+</sup> (Vidal et al., 2005). In addition, we estimated isochore from homogenization temperatures of fluid inclusions in quartz (101-130°C) that coexisted with chlorite used for the estimate of formation temperature. Integrating all these results, we analyzed P-T path. First, from blueschist facies, temperature increased and pressure decreased to reach or pass through the reaction line between pumpellyite and epidote. Next, pressure decreased toward the isochore of fluid inclusions at constant temperature of c. 300 °C inferred from deformation microstructures in quartz indicating brittle-ductile transition. Furthermore, temperature and pressure decreased toward the formation temperature of chlorite. Finally, we have speculated that the increase of temperature from blueschist to greenschist facies could have been caused by fluid infiltration in the rocks.

SMP046-14

Room:201B

Time:May 27 12:00-12:15

## A clockwise P-T path deduced from metapelites and aluminosilicates-bearing veins from the Tseel terrane, SW Mongolia

burenjargal ulziiburen<sup>1\*</sup>, Atsushi Okamoto<sup>1</sup>, Noriyoshi Tsuchiya<sup>1</sup>

<sup>1</sup>Tohoku University

The Tseel terrane of the Central Asian Orogenic Belt, SW Mongolia, contains a record of amphibolite-facies (locally granulite-facies) metamorphism related to several igneous activities. In the central part of the Tseel area, the andalusite (And) + sillimanite (Sil) + kyanite (Ky)-bearing quartz veins occur, whereas only sillimanite occurs in host pelitic gneisses, that contain garnet + biotite + plagioclase + quartz. Textural relations indicate that aluminosilicate polymorphs formed in the order of And-Ky-Sil. Garnet in a sample of gneiss collected from near an aluminosilicates-bearing quartz vein shows compositional zoning, characterized by decreases in Ca and Mn from core to rim, and increases in Fe and Mg, along with minor retrograde zoning at the outermost rim.

We calculate P-T conditions by garnet-biotite geothermometry and garnet-biotite-plagioclase geobarometry based on compositional zoning in garnet, assuming constant compositions for biotite and plagioclase, to roughly constrain the P-T path during garnet growth. This approach is based on the following assumptions: (1) biotite, plagioclase and quartz coexisted with garnet; (2) the compositional ranges of biotite and plagioclase during garnet growth were retained in the thin section; and (3) intracrystalline diffusion was negligible within the crystals. For individual samples, we selected the biotite compositions with highest and lowest Mg/Fe<sup>2+</sup> ratio and plagioclase compositions with anorthite content,  $X_{An}$ . The P-T estimates along the garnet zoning is carried out for four cases as follows: (1) highest  $X_{An}$ , highest Mg/Fe<sup>2+</sup> Bt; (2) highest  $X_{An}$ , lowest Mg/Fe<sup>2+</sup> Bt; (3) lowest  $X_{An}$ , highest Mg/Fe<sup>2+</sup> Bt; and (4) lowest  $X_{An}$ , lowest Mg/Fe<sup>2+</sup> Bt.

We obtained the decompression P-T path from the kyanite stability field (530-570 C and 6.0-9.6 kbar) to the sillimanite stability field (590-620 C and 2-6 kbar), with slight increase in temperature. Although garnet does not record the P-T conditions at the burial stage, the occurrence of aluminosilicates indicates the change from the andalusite stability field to kyanite stability field. These observations suggest that the metamorphic rocks in the Tseel terrane experienced a clockwise P-T path, although the peak pressure remains unknown.

Microthermometry was based on analyses of fluid inclusions in quartz in an aluminosilicates-bearing quartz vein collected from the locality of sample 0701c. Heating and cooling experiments were performed for the primary inclusions within quartz to measure the homogenization (Th) and ice melting (Tim) temperatures. The values of Th are scattered over the range 110-240 C, with most between 160 and 200 C (mean value, 171 ± 28 (1 $\sigma$ ) C). The values of Tim range from 24.0 to 29.6 C, with a mean value of 26.8 ± 1.3 (1 $\sigma$ ) C, corresponding to salinity of 10.2 ± 1.6 wt.% NaCl equivalent. Microthermometric analyses of fluid inclusions reveal that the aluminosilicates-bearing quartz veins formed in the kyanite stability field (530-600 C and 6.0-8.5 kbar). Abundant fluid supply along fractures would have enhanced the formation of coarse-grained kyanite in quartz veins.

The P-T path during the exhumation in the Tseel area cannot be explained by subduction of old slab, but is well consistent with the geothermal gradient along the interface between the slab and the arc crust just after the ridge subduction (after c. 1 Myr). The intrusion of granitoids and mafic dikes and high temperature metamorphism of the Tseel terrane would be caused by the subduction of young oceanic lithosphere during the evolution of CAOB in the Devonian ages.

**Keywords:** Tseel terrane, aluminosilicates, garnet, clockwise P-T path, ridge subduction



SMP046-15

Room:201B

Time:May 27 12:15-12:30

## Multi-events metamorphism of the eclogites in the Lake Zone, southwest Mongolia

Javkhlan Otgonkhuu<sup>1\*</sup>, Akira Takasu<sup>1</sup>, Dash Bat-ulzii<sup>2</sup>, Gerel Ochir<sup>2</sup>

<sup>1</sup>Shimane Univ., <sup>2</sup>Mongolian Univ. of Sci. and Tech.

Early Cambrian eclogites have been described from the Chandman district in the Lake Zone, southwest Mongolia (Hanzl and Aichler, 2006; Takasu et al., 2008; Stipska et al., 2010). The Alag Khadny metamorphic complex consisting of metamorphic rocks and ultramafic bodies is exposed for about 10 km across and 4 km wide, and it occurs between the ophiolite complex and migmatized metamorphic rocks in the Lake Zone. Several lenticular-shaped bodies of eclogites and amphibolites (max. 2 km x 0.8 km) occur in the matrix of orthogneisses and minor pelitic gneisses.

Eclogites consist mainly of garnet, omphacite ( $Jd < 46\%$ ), and amphibole with subordinate amounts of epidote, phengite, paragonite, plagioclase, biotite, K-feldspar, rutile, titanite, quartz, calcite, hematite, ilmenite and zircon. The eclogites experienced three metamorphic events i.e. the precursor metamorphic event (M1) of HT-amphibolite facies; HP metamorphism (M2) of the eclogite facies; and HP-metamorphism (M3) of the epidote-amphibolite facies.

Garnets occur as porphyroblast and they show a prograde zoning. The core of the garnets contain polyphase and single grain inclusions of high  $TiO_2$  (up to 1.32%) taramite, taramite+quartz, Fe-pargasite, tschermakite, plagioclase ( $An < 19$ )+biotite+epidote. Those inclusions indicate relatively high-temperature metamorphism of amphibolites facies conditions (M1).

The prograde stage of the first HP metamorphic event (M2) is characterized by polyphase and single grain inclusions in the garnets such as barroisite+taramite+epidote+quartz, barroisite+Mg/Fe-hornblende+quartz, plagioclase ( $An = 3-5$ )+epidote, chlorite, calcite, and rutile. The peak eclogite facies conditions of 560-680 C and 22-25 kbar (Ravna, 2000; Ravna and Terry, 2004) estimated by the compositions of garnet+omphacite ( $Jd < 46$ )+phengite ( $Si = 6.58-7.11$ ) assemblage. The retrograde stage of eclogite facies is characterized by symplectite of sodic plagioclase ( $An = 1-11$ )+amphibole and/or Na-poor clinopyroxene ( $Jd = 2-25$ ). These mineral assemblages give 450-560 C and 4-11 kbar (Holland and Blundy, 1994; Holland, 1983).

The second HP metamorphism of the epidote amphibolites facies (M3) is characterized by prograde zoned amphiboles with winchite, actinolite, tremolite core and barroisite rim. They contain inclusions of garnet, omphacite and symplectite of clinopyroxene+sodic plagioclase suggesting that the amphiboles crystallized after the M2 metamorphism. The cores of the amphiboles indicate 300-400 C and 3-8 kbar, whereas the rims indicate >400-600 C and 3-12 kbar (Otsuki and Banno, 1990). Taramite/tschermakite outermost rim is occasionally developed.

### Reference:

- Hanzl, P. and Aichler, J., (2007) Pracha, Czech Geological Survey  
Holland, T., 1983 Contributions to Mineralogy and Petrology, 82, 214-220;  
Holland, T.J.B and Blundy, J.D., 1994. Contributions to Mineralogy and Petrology, 116, 433-447.  
Otsuki, M. and Banno, S., 1990 Journal of Metamorphic Geology., 8, 425-439  
Ravna, E.J.K., 2000. Journal of Metamorphic Geology, 18, 211-219  
Ravna E.J.K and Terry, M.P., 2004. Journal of Metamorphic Geology, 22, 579-592  
Stipska, P. et al., 2010. Journal of Metamorphic Geology, 1-19.  
Takasu, A. et al., 2008 Abstract of the 115th Annual Meeting of the Geological Society of Japan. 246p. (in Japanese)

Keywords: eclogite, Alag Khadny metamorphic complex, Lake Zone, Mongolia

SMP046-16

Room:201B

Time:May 27 12:30-12:45

## Deformation of Lake Shorelines and Mid Crustal Flow in Tibet

Simon Wallis<sup>1\*</sup>, Kazuhiro Ozawa<sup>1</sup>, Hiroshi Mori<sup>1</sup>, Takeshi Sagiya<sup>2</sup>, Toshio Nakamura<sup>3</sup>

<sup>1</sup>Earth & Planetary Sciences, Nagoya Uni., <sup>2</sup>Geodynamics Group, Nagoya Uni., <sup>3</sup>Center Chronological Research, Nagoya U.

The mid crust beneath Tibet is generally thought to be highly mobile low viscosity material. A low viscosity layer of mid crustal material can account for the relatively flat nature of the high plateau?high elevation but low relief?and injection of mid crust into low lying regions around the Tibetan Plateau is thought to be one of the main processes involved in the expansion of the Plateau. Mechanical modeling shows that many of the first order features of the Tibetan Topography can be explained by the presence of a mid crust with a viscosity of  $10^{19}$  Pa s or less. However, there has been no independent quantitative estimate of the effective viscosity of the mid crust. Lake shorelines offer a way to achieve this.

Despite its low rainfall, Tibet contains a large number of lakes due to the lack of water outlets from the central plateau. Many of these lakes are surrounded by well-preserved paleo shorelines. The presence of these shorelines shows that the lakes were once much larger than they are now. One of the largest lakes in Tibet is Lake Nam Co, which lies 150 km to the north of Lhasa. This lake shows good development of plaeo-shorelines and is of a suitable size to investigate properties of the mid crust. When there is a drop in the water level of a lake, it reduces the weight on the underlying crust and resulting in a buoyancy force that tends to uplift the substrate. The maximum uplift possible is determined by the ratio of the densities of water to rock: approximately 1 m of uplift for every 3 m decrease in the water level. The reason for the uplift is the inflow of mobile rock at depth. Re-equilibration will not be instantaneous?it will take time for crustal flow to occur. The time scale for this crustal flow depends mainly the geometry of the lake basin and the viscosity of the crust. The geometry is well-known and viscosity can then be estimated from measurements of the amount of uplift and the time that it took for the uplift to occur. Shorelines are palaeo-horizontal markers and, therefore, any uplift can be recognized by careful measurement that reveals present day deviations from horizontal. Preliminary results of age dating and surveys of shorelines using kinematic GPS show the potential of this methodology for obtaining good first order estimates of the mid crustal viscosity.

Keywords: Tibet, Lake shorelines, Crustal Flow, Mid crustal viscosity

SMP046-P01

Room:Convention Hall

Time:May 27 14:00-16:30

## Deformation experiment of Serpentine with preferred orientation

Yoshito Urata<sup>1\*</sup>, Jun-ichi Ando<sup>1</sup>, yusuke soda<sup>2</sup>

<sup>1</sup>Graduate School of Hiroshima University, <sup>2</sup>Waseda University

We conducted constant strain rate experiment of antigorite serpentinite, in order to understand the effect of microstructural anisotropy on deformation behavior. The sample is naturally deformed foliated antigorite serpentinite which is characterized by preferential arrangement of (001) of antigorite parallel to the foliation. We prepared two types of oriented starting samples, whose foliations were set at 30 degree and 90 degree with respect to the axial stress. Experimental conditions were 500 C and 550 C at ca. 1 GPa confining pressure with 600 um/h displacement rate of piston. The experimental data indicate that the maximum and yield strengths of the 90 degree are ca. 40 % larger than those of 30 degree and that those of 500 C are 60 to 70 % larger than those of 550 C. The stress drop occurred at the final stage of all experiments. The microstructural observations with optical and scanning electron microscopes suggest that the yielding and stress drop in 30 degree experiments were due to kinking, and the deformation concentration on the axis of kink, respectively. On the contrary, those of 90 degree were due to plastic deformation of antigorite itself and extensional breakage of antigorite grains, respectively. All these behaviors are thought to be caused by two typical characters of foliated antigorite serpentinite; 1) easy to make open crack parallel to foliation if compression stress applies in the direction parallel to the foliation, and 2) weakness of (001) for extensional stress.

Keywords: Antigorite, Serpentine, Solid medium deformation experiment, SEM

SMP046-P02

Room:Convention Hall

Time:May 27 14:00-16:30

## Subduction related Antigorite CPO patterns from forearc mantle in the Sanbagawa belt, southwest Japan

Aya Nishii<sup>1\*</sup>, Simon Wallis<sup>1</sup>, Tomoyuki Mizukami<sup>2</sup>, Katsuyoshi Michibayashi<sup>3</sup>

<sup>1</sup>Earth & Planetary Sci., Nagoya Univ., <sup>2</sup>Nat. Sci. Tech., Kanazawa Univ., <sup>3</sup>Instit. Geosci. Shizuoka Univ.

Antigorite (Atg) is stable throughout large parts of the wedge mantle of most subduction zones. Atg shows very strong acoustic anisotropy, and recent studies have emphasized that the crystallographic preferred orientation (CPO) of Atg should be considered as a possible cause of seismic anisotropy in convergent margins.

Only a few Atg CPO patterns have been published (Bezacier et al., 2010; Hirauchi et al., 2010; Moortele et al., 2010; Soda & Takagi, 2010). From these limited data, two main types of Atg CPO pattern can be defined: one with an a-axis parallel to the stretching direction (A-type) and the other with the b-axis parallel to the stretching direction (B-type). In this study, we report antigorite CPO patterns from the Higashiakaishi (HA) body, a sliver of forearc mantle preserved in the Sanbagawa belt of southwest Japan. These CPO patterns are a further example of the B-type antigorite patterns.

Because Atg has a low plastic yield strength, it is possible that original orientations of the Atg crystals may be affected by mechanical damage caused by production of the thin sections used for measurement. However, statistical analysis using the eigen vector method of Atg CPO in two thin sections from two distinct directions in the same sample (YZ-section perpendicular to foliation and lineation and XZ-section perpendicular to foliation and parallel to lineation) shows no significant differences. Atg CPO developed during the same phase of deformation was also stronger in the sample with a greater proportion of Atg: the opposite to that expected if Atg CPO is disturbed by sample preparation. We conclude that sample preparation by standard polishing techniques has no significant affect on the resulting CPO.

Seismic anisotropy associated with the Atg-bearing HA peridotite calculated using the combined Olivine and Atg CPO patterns requires thicknesses of 1.47-4.6 km for a time delay of 0.1 s and 5.31-11.56 km for a time delay of 1 s. The large range of possible thicknesses represents the difference between Reuss and Voigt averages.

[References] Bezacier, L. et al. 2010, EPSL; Hirauchi, K. et al. 2010, EPSL 299, 196-206; Moortele, B. et al. 2010, J. Microscopy 239, 245-248; Soda, Y. and Takagi, H. 2010, J. Structural Geology 32, 792-802.

Keywords: Antigorite, CPO, seismic anisotropy, Higashi Akaishi body

SMP046-P03

Room:Convention Hall

Time:May 27 14:00-16:30

## The deepest peridotites in ocean floor: Tonga trench peridotites revealing forearc extension

Yuri Shinkai<sup>1\*</sup>, Katsuyoshi Michibayashi<sup>1</sup>, Shigeki Uehara<sup>1</sup>, Teruaki Ishii<sup>2</sup>

<sup>1</sup>Instit. Geosci. Shizuoka Univ, <sup>2</sup>Fukada Geological Institute

The Tonga trench is one of the deepest trenches in the world. We used peridotite samples collected from dredge hauls by Boomerang Leg 8 Cruise aboard R/V Melville in 1996 at the deep landward trench slope (19°15.19S, 172°56.29W; depth 8,194-9,371m; Bloomer et al., 1996, Fall Meeting, Abstract, OG32B-01). Most of samples are remarkably fresh, indicating that tectonic erosion is active in the Tonga trench. The samples are harzburgites and show some variations in microstructure consisting of dominantly coarse (>5mm) granular texture to minor fine-grained (~0.5mm) parts. They contain high-Cr# spinels in a range between 0.5 and 0.8 with very low Ti contents, suggesting that these peridotites were derived from the Tonga forearc. Equilibrium temperatures estimated by Ca in orthopyroxene geothermometer are approximately 900-1250. Olivine fabrics are characterized by intense [100]-fiber pattern, which could be developed by transtension type of strain (Tommasi et al., 1999, EPSL, 168, 173-186). These indicate that the Tonga trench peridotites have probably been derived from the lithospheric mantle due to the forearc extension during slab rollback (Smith et al., 2001, Science, 292, 713-716).

Keywords: Tonga Trench, peridotite, crystal-preferred orientation, forearc, slab rollback

SMP046-P04

Room:Convention Hall

Time:May 27 14:00-16:30

## Elastic Wave Velocities of Antigorite-Bearing Serpentine Mylonites

Yuto Shirasugi<sup>1\*</sup>, Tohru Watanabe<sup>1</sup>, Katsuyoshi Michibayashi<sup>2</sup>

<sup>1</sup>University of Toyama, <sup>2</sup>Shizuoka University

The relationships between elastic wave velocities and petrofabrics were studied in antigorite-bearing serpentinite mylonites. Rock samples with antigorite content from 40 to 80 vol.% were collected from the Happo ultramafic complex, Central Japan. Compressional and shear wave velocities were measured by the pulse transmission technique at room temperature and confining pressures of up to 180 MPa. Petrofabrics were examined by optical microscopy and SEM-EBSD. Olivine a- and c-axes are weakly oriented perpendicular to the foliation and parallel to the lineation, respectively. Antigorite b- and c- axes are distinctly oriented parallel to the lineation and perpendicular to the foliation, respectively. Samples show strong anisotropy of velocity. The compressional wave velocity is fastest in the direction parallel to the lineation, and slowest in the direction perpendicular to the foliation. The shear wave oscillating parallel to the foliation has higher velocity than that oscillating perpendicular to the foliation. As the antigorite content increases, the mean velocity decreases but both azimuthal and polarization anisotropies are enhanced. Measured velocities were compared with velocities calculated from petrofabric data by using Voigt, Reuss and VRH averaging schemes. All averaging schemes show velocity anisotropy qualitatively similar to measurements. There are large velocity differences between Voigt and Reuss averages (0.7~1.0 km/s), reflecting the strong elastic anisotropy of antigorite. Measured velocities are found between Reuss and VRH averages. We think that the relatively low velocity is due to the platy shape of antigorite grains, the well developed shape fabric and their strong elastic anisotropy. Measured velocities will be compared with calculation considering layered structures in serpentinite mylonites.



SMP046-P05

Room:Convention Hall

Time:May 27 14:00-16:30

## Elastic wave velocity and petrofabric of amphibolites and eclogite from the Sanbagawa metamorphic belt

Tohru Watanabe<sup>1</sup>, Kenya Ono<sup>1\*</sup>, Motoki Kitano<sup>1</sup>, Akiyoshi Higuchi<sup>1</sup>, Yuto Shirasugi<sup>1</sup>, Katsuyoshi Michibayashi<sup>2</sup>

<sup>1</sup>University of Toyama, <sup>2</sup>Shizuoka University

Hydration and dehydration processes in the oceanic crust materials play important roles in subduction zone processes as water transportation and slab earthquakes. Seismological constraints to constituent materials and their water content in the slab crust should lead to further understanding of subduction zone processes. It is thus essential to understand elastic properties of slab crust materials. We have made velocity measurements and petrofabric observations on amphibolites and an eclogite from the Sanbagawa metamorphic belt, which might have been subducted oceanic crust.

Rock samples are amphibolite schist, garnet amphibolite (Iratsu amphibolite body, Shikoku-chuou, Ehime Pref.), and eclogite (Higashi-Akaishiyama peridotite body, Shikoku-chuou, Ehime Pref.). The densities are 3120, 3250, 3460 kg/m<sup>3</sup>, respectively at room conditions. A rectangular parallelepiped (the edge length ~ 30 mm) was cut from each rock sample for velocity measurements. Two faces are parallel to the foliation plane, two faces perpendicular to the elongation direction, and the remaining two faces perpendicular to the foliation plane and parallel to the elongation direction. Preliminary velocity measurements were made at room conditions by the pulse transmission technique using Pb(Zr, Ti)O<sub>3</sub> transducers with the resonant frequency of 2 MHz. One compressional wave velocity and two shear wave velocities were measured in each of three orthogonal directions. Two shear waves propagating in one direction oscillate in mutually orthogonal directions. Arithmetic means of V<sub>p</sub> and V<sub>s</sub> are 5.70 km/s and 3.60 km/s for amphibolite schist, 3.92 km/s and 2.69 km/s for garnet amphibolite and 5.50 km/s and 3.60 km/s for eclogite. The azimuthal anisotropy of V<sub>p</sub> is around 10% for amphibolite schist and eclogite, whereas it is 36% for garnet amphibolite. However, these velocity values cannot be compared with petrofabrics, because they must be affected by pores in rock samples. We are now conducting velocity measurements under the confining pressures of up to 180 MPa to remove the influence of pores. The relationship between velocity under the confining pressure and petrofabrics will be presented in our poster.

Keywords: amphibolite, eclogite, elastic wave velocity, petrofabric

SMP046-P06

Room:Convention Hall

Time:May 27 14:00-16:30

## Application of Micro-XRF analysis for estimation of igneous mineral compositions from subduction zone meta-peridotites

Tatsuya Hattori<sup>1\*</sup>, Tomoyuki Mizukami<sup>1</sup>

<sup>1</sup>Natural Science, Kanazawa university

Ultramafic rocks in the metamorphic belt have been generally re-crystallized into fine-grained aggregates and the elemental components in the rocks have been re-distributed into newly formed minerals including hydrous phases. In order to discuss the igneous processes related to the origins, we have to estimate such chemical and modal modification. For examples, Ni, Mg and Fe are important indicators of fractional crystallization and partial melting but they are expelled out from olivine and pyroxene into magnetite (Mag), antigorite (Atg) and sulfides (pentlandite (Pn) etc.). The precise modal compositions of the metamorphic constituents are necessary for reproduction of the igneous mineral compositions.

We test an elemental mapping using a Micro-XRF Analyzer (XGT-5000, HORIBA) to determine modal compositions of metamorphic minerals. In this system, an X-ray beam is focused on a polished thin section (30 micron thick) in a diameter of 0.1 mm and fluorescent X-ray maps (512 x 256 pixels for a region of 2 x 1 cm<sup>2</sup>, for example) of elements with higher atomic number than Ca are gained using an EDS detector.

Analytical samples are ultramafic rocks in Higashi-akaishi peridotite body (HA) in Sanbagawa metamorphic belt, southwest Japan. They have been a part of olivine (Ol)-clinopyroxene (Cpx) cumulate composed of dunite and Cpx-bearing rocks. The present constituent minerals are Ol, Cpx, Atg, Cr-Spl, ferricht Chr, Mag, Pn. The modal compositions of Cpx, Mag, Cr-Spl/ferricht Chr and Pn are determined using Ca map, Fe map, Cr map and Ni map, respectively. Proportions between Ol and Atg are determined by point counting. As a result, modal compositions are determined to one place of decimal, for example: Ol (4.2%), Cpx (32.1%), Spl (0.5%), Mag (2.6%), Pn (0.1%) and Atg (60.5%).

Primary compositions are calculated from the present mode and mineral chemistry assuming a primary assemblage of Ol + Cpx + Cr-Spl, a semi-closed system with additions of H<sub>2</sub>O and SiO<sub>2</sub> and  $KD(Ol-Cpx) = 0.8$ . Resultant Mg# and NiO content in Ol show significant effect of metamorphic changes from 0.862 to 0.822 and from 0.17 to 0.11, respectively. Our calculation indicates that a modal composition of Pn should be determined with a precision to two places of decimals. Measuring sizes and numbers of sulfide grains under a reflecting microscope will be the best way to make reliable estimation of primary NiO in Ol instead of a XRF mapping.

Keywords: Micro-XRF analysis, modal composition, pentlandite, subduction zone, meta-peridotites

SMP046-P07

Room:Convention Hall

Time:May 27 14:00-16:30

## Mode of occurrence of chlorine-rich biotite and zircon in the pelitic gneiss from Sor Rondane Mountains, East Antarctica

Fumiko Higashino<sup>1\*</sup>, Tetsuo Kawakami<sup>1</sup>, M. Satish-Kumar<sup>2</sup>, Noriyoshi Tsuchiya<sup>3</sup>, Masahiro Ishikawa<sup>4</sup>, Geoff Grantham<sup>5</sup>

<sup>1</sup>Kyoto University, <sup>2</sup>Shizuoka University, <sup>3</sup>Tohoku University, <sup>4</sup>Yokohama National University, <sup>5</sup>Council for Geoscience, South Africa

The Bt-Grt-Sil gneiss from northern Balchenfjella, Sor Rondane Mountains, East Antarctica contains Grt porphyroblast (5-10 mm in diameter) and fine-grained Grt (100-200 μm in diameter). The Grt porphyroblast has characteristic zoning in P. The core is P-rich with oscillatory zoning in P, and the rim is P-poor without oscillatory zoning. In this way, the core-rim boundary is defined by the discontinuous P zoning. This discontinuity suggests that the Grt porphyroblast has experienced resorption and recrystallization (e.g., Kawakami & Hokada, 2010). The Grt core is homogeneous in composition while Fe and Mn increase and Mg and Ca decrease toward the rim. This trend is significant where the Grt is in contact with matrix Bt. The fine-grained Grt has similar composition with the Grt porphyroblast rim, suggesting that the rim and the fine-grained Grt crystallized simultaneously.

Chlorine-rich Bt (0.08-1.08 wt%) and coarse Zrn (100 μm in diameter) are included exclusively in the rim of the Grt porphyroblast. Bt in the matrix and that included in the core of the Grt porphyroblast are Cl-poor (< 0.06 wt%). Coarse-grained Zrn is present in the matrix and also included in the fine-grained Grt. Only one grain of tiny Zrn (20 μm in diameter) is found included in the core of the Grt porphyroblast so far.

Almost all of the Cl-poor Bt grains in the matrix give lower  $X_{Mg}$  [=  $Mg/(Mg+Fe_{total})$ ] ( $X_{Mg} = 0.56 \pm 0.03$ ) than the Cl-rich Bt included in the P-poor rim of the Grt ( $X_{Mg} = 0.60 \pm 0.04$ ). Detailed examination of retrograde Fe-Mg exchange between the inclusion Bt and the host Grt revealed that the Cl-rich Bt was originally as Mg-rich as the matrix Bt and changed its composition to the Mg-rich one through the retrograde Fe-Mg exchange reaction between Grt. There is a possibility, therefore, that matrix Bt was once Cl-rich and lost Cl via interaction with the Cl-poor fluid that subsequently infiltrated into the matrix.

The presence of resorption texture at the core-rim boundary of the Grt porphyroblast, and the intimate coexistence of Cl-rich Bt and coarse-grained Zrn in the rim of the Grt porphyroblast implies the genetic relationship between them. The Cl-rich Bt may be formed through the effect of Cl-rich fluids during metamorphism (e.g., Sisson, 1987) or through magmatic-hydrothermal evolution (e.g., Coulson, 2001). Therefore, it is likely that the Cl-rich Bt included in the garnet rim is the evidence for the infiltration of Cl-rich fluid, and coarse-grained Zrn (at least the rim part) and the Cl-rich Bt were formed almost simultaneously, during or after the Grt porphyroblast resorption. Future dating of included Zrn may help constrain the timing of Cl-rich fluid infiltration during the metamorphism in the Sor Rondane Mountains.

Keywords: chlorine, zircon, resorption, metamorphic fluid

SMP046-P08

Room:Convention Hall

Time:May 27 14:00-16:30

## Two modes of occurrence of "arrested charnockite" in Sri Lanka

Yukiko Yamasaki<sup>1\*</sup>, Takeshi Ikeda<sup>1</sup>, Yoichi Motoyoshi<sup>2</sup>, Yoshikuni Hiroi<sup>3</sup>, Bernard Prame<sup>4</sup>

<sup>1</sup>Earth and Planet. Sci., Kyushu Univ., <sup>2</sup>National Institute of Polar Research, <sup>3</sup>Earth Sci., Chiba Univ., <sup>4</sup>Geological Survey of Sri Lanka

Felsic to intermediate lithologies charnockitized incompletely have been called as "arrested charnockite". These types of high-grade metamorphic rocks occur in Sri Lanka, as well as Antarctica and South India that formerly constituted Gondwana-land. We described the mode of occurrence and chemical composition of the constituent minerals of the arrested charnockite from two outcrops in Sri Lanka. In the first outcrop near Kurunegala, charnockite occurs as a number of patches with lenticular to ovoidal shapes. The long axes of the patches are not parallel to the foliation defined in the surrounding felsic gneiss. The boundary between charnockite patch and surrounding gneiss is obscure. Compositional layerings composed of Hbl-rich domain and Hbl-poor domain disappear gradually into charnockite. In the second outcrop near Kandy, charnockite is developed as a vein along a fracture of Grt-Bt felsic gneiss. The boundary is obscure. The compositional layerings and mineral preferred orientation recognized in the host gneiss are preserved in the charnockite domain.

In the charnockite from the first outcrop, orthopyroxene occurs commonly. The orthopyroxene grains are locally replaced by cummingtonite along rim and cleavage. Orthopyroxene appears at inner portions of the boundary of charnockite patches recognized by naked eye. The modal abundance of orthopyroxene increases with increasing distance from the boundary. In contrast, vein-like charnockite from the second outcrop, many symplectites occur while orthopyroxene is not found. The surrounding gneiss also has symplectites locally. The symplectites are composed of various mineral assemblages, such as Amp+Pl+Ilm+Mag+Cpx, Amp+Pl+Ilm+Mag and Pl+Ilm+Mag.

Amphibole can be divided into two based on the chemical composition. One has higher Ca and Al than another. The former occurs in the host gneiss and also in the charnockite that defines gneissosity. The latter amphibole (cummingtonite) is recognized only in the charnockite and either replaces orthopyroxene in the patchy charnockite or consists of symplectite in the vein-like charnockite. The cummingtonite in the symplectite showing similar composition to that replacing orthopyroxene suggests that the symplectites in the vein-like charnockite were originally orthopyroxene which has broken down during retrograde metamorphism.

Keywords: charnockite, Sri Lanka

SMP046-P09

Room:Convention Hall

Time:May 27 14:00-16:30

## Finding of brown hornblende bearing basic rock from Bungo-Ohno, Oita prefecture

Shunji Tojo<sup>1\*</sup>, DOHI, Shintaro<sup>1</sup>

<sup>1</sup>Fukuoka University of Education

The Asaji metamorphic rock have been studied by many authors (e.g. Ohshima et al., 1971; Fujii et al., 2008). As the result of field surveying, several basic rock samples which contain brown hornblende, size of up to 0.8 mm were collected. The sample localities are concordant to the area of basic rock belt, and belonged to the metamorphic zone B described by Ohshima et. al. (1971). However, the brown hornblende is characteristic mineral of the metamorphic zone C, and uniquely found in xenolith or roof pendant of the plutonic rocks. There have been no described plutonic rock near around the localities of these basic rocks we found. It is possibly suggested that the potentially underlaying of plutonic rocks at shallow depth.

Keywords: Asaji metamorphic rock, brown hornblende, contact metamorphism

SMP046-P10

Room:Convention Hall

Time:May 27 14:00-16:30

## Metamorphic and granitic tectonic blocks of the Atokura Nappe in the Yorii-Ogawa district, central Japan

Akira Ono<sup>1\*</sup>

<sup>1</sup>None

The Atokura Nappe in the northeastern Kanto Mountains is composed of Yorii pyroclastic rocks, Yorii Formation, Atokura Formation, Kinshozan Quartzdiorite, Greenstone Merange and various small tectonic blocks (Figure 1). The geological bodies are usually in contact with each other by high-angle faults. Metamorphic and granitic rocks often occur as small tectonic blocks. The representative ones are (a) mid-Cretaceous metamorphic and granitic rocks, (b) early Paleogene Kiroko metamorphic rocks within the Greenstone merange (Ono, JpGU Meeting 2008, G122- P002), (c) late Cretaceous Yorii Granitoids in the Mure.

### mid-Cretaceous metamorphic rocks

A small tectonic block of mid-Cretaceous metamorphic and granitic rocks is exposed between the early Paleogene Yorii Formation and the late-Permian Kinshozan Quartzdiorite near Mt. Kinshozan (Figure 1). Coarse-grained garnet-bearing gneisses are found in the southern part of the tectonic block. Fine-grained chlorite-muscovite schists and chlorite-amphibole schists are exposed in the northern part. Calcareous rocks of approximately 20m thick are exposed in the easternmost part of the block. The calcareous rocks show various rock textures. A few pelitic and tuffaceous thin layers are intercalated. Fusulinacean fossils are found in the calcareous rocks and pelitic rocks. Hence, the low-grade metamorphic rocks are metamorphosed Paleozoic formations. This fact suggests that highly metamorphosed calcareous rocks of the Atokura Nappe are also metamorphosed Paleozoic rocks. The mid-Cretaceous metamorphic rocks have properties similar with those of the Hitachi metamorphic rocks in the Abukuma belt.

### early Paleogene Kiroko metamorphic rocks


The Greenstone Merange which is exposed in the southernmost part of the Atokura Nappe (Figure 1) is mainly composed of the Kiroko metamorphic rocks, serpentinite, actinolite-rocks and various metamorphic and granitic rocks. Serpentinites are poor and granitic rocks are common in the eastern part of the Greenstone Melange. The Kiroko metamorphic rocks mainly consist of mafic rocks, pelitic rocks and psammitic rocks. They are well-recrystallized low-grade metamorphic rocks. There is no evidence for recrystallization and alteration after the main phase of the regional metamorphism.

The Greenstone Melange is in contact with the Atokura Formation by high-angle faults. Serpentinite is common adjacent to the Atokura Formation although serpentinite is rare in the eastern part of the Greenstone Melange. The common occurrence of serpentinite near the high-angle faults suggests that serpentinites played important roles in the formation of the high-angle faults in the root zone of the Atokura Nappe.

### late Cretaceous Yorii Granitoids

The Yorii Granitoids consist of aplite and biotite tonalite which are exposed in the Mure. Biotite tonalite is massive and medium in grain sizes. Biotite is partly altered to chlorite. Magnetite is not observed. The biotite tonalite is in contact with a small tectonic block of chert, slate and sandstone by a high-angle fault. The biotite tonalite may be one of the late Cretaceous granitoids of the Southwest Japan judging from the lack of similar granitic rocks in the Atokura Nappe.





Keywords: Atokura Nappe, metamorphic rocks, serpentinite, granitoid, fusulinid