

BPT002-01

会場:201B

時間:5月25日 08:30-08:45

琉球列島産現生腕足動物 (*Basiliola lucida*) 殻の炭素・酸素同位体組成の個体差 Intraspecific Variations in Carbon and Oxygen Isotope Compositions of a Modern Brachiopod Collected off Okinawa-jima

高柳 栄子^{1*}, 浅海 竜司², 大竹 二雄³, 阿部 理¹, 北川 浩之¹, 井龍 康文¹

Hideko Takayanagi^{1*}, Ryuji Asami², Tsuguo Otake³, Osamu Abe¹, Hiroyuki Kitagawa¹, Yasufumi Iryu¹

¹ 名古屋大学・院・環境, ² 琉球大学・超域, ³ 東京大学・大気海洋研

¹Nagoya University, ²University of the Ryukyus, ³University of Tokyo

Carbon ($d^{13}C$) and oxygen ($d^{18}O$) isotope compositions in rhynchonelliform brachiopod shell calcite have been widely used as proxies of $d^{13}C$ of dissolved inorganic carbon (DIC) and $d^{18}O$ of ancient seawater, respectively. The use of brachiopods as a paleoenvironmental proxy is based on the presupposition that the shell calcite is precipitated in isotopic equilibrium with ambient seawater. Recently, it was shown that the modern brachiopod calcite display variable carbon and oxygen isotope offsets from the range of equilibrium calcite (= calcite precipitated in isotopic equilibrium with ambient seawater) even within a single shell and degree of the disequilibrium varies from species to species [e.g., Auclair *et al.*, 2003, *Chem. Geol.*, 202, 59-78; Yamamoto *et al.*, 2010a, *Palaeo-3.*, 291, 348-359; Yamamoto *et al.*, 2010b, *G-cubed*, 11, Q10009]. However, a single specimen was examined for each species in the previous studies that dealt with within-shell variations. Therefore, further geochemical investigations are needed to reveal intraspecific variations in isotopic compositions to establish the isotopic compositions of brachiopod shells as a much more reliable paleoenvironmental proxy.

This study presents intraspecific variations in carbon and oxygen isotope profiles along the growth axis of modern brachiopod shells of *Basiliola lucida* collected from shelf to shelf slope environments (180-320 m water depth) off Okinawa-jima and Amami-o-shima, southwestern Japan. The $d^{13}C$ and $d^{18}O$ values of each shell are rather constant with no significant variations. The $d^{13}C$ values are greater than those of equilibrium calcite with two outlier shells. The $d^{18}O$ profiles show enrichment in ^{18}O relative to equilibrium calcite throughout the growth axis, which underestimates seawater temperature, although the differences in the $d^{18}O$ values among the samples correspond to those of seawater temperatures among the brachiopod growth sites. Significant offsets of $d^{13}C$ and $d^{18}O$ values from the range of equilibrium calcite indicate that the isotopic compositions in the shells of *B. lucida* are influenced by vital effects that are generally explained by kinetic fractionation and metabolic effects. However, the degrees of the effects on $d^{13}C$ and $d^{18}O$ values are comparatively constant in each individuals. In conclusion, the use of *B. lucida* as a proxy of ancient seawater temperature is not very recommend. However, if water depths of the brachiopod-yielding horizons are determined independently by other fossils (e.g., benthic foraminifers) in a given geologic section, $d^{18}O$ value of *B. lucida* can be used as a proxy to determine depth gradient of seawater temperature for the past.

キーワード: 腕足動物, 炭素同位体組成, 酸素同位体組成, 沖縄沖, 奄美沖

Keywords: brachiopod, carbon isotope composition, oxygen isotope composition, off Okinawa-jima, off Amami-o-shima

BPT002-02

会場:201B

時間:5月25日 08:45-09:00

ガンジス川およびブラマプトラ川堆積物の地球化学的研究 Geochemistry of sediments from the Ganges and Brahmaputra Rivers

Hossain H. M. Zakir^{1*}, Barry P. Roser², M. Sultan-UI-Islam³

H. M. Zakir Hossain^{1*}, Barry P. Roser², M. Sultan-UI-Islam³

¹Jessore Science and Technology Univ., ² 島根大学, ³Rajshahi University

¹Jessore Science and Technology Univ., ²Shimane University, ³Rajshahi University

World-scale big rivers play an important role in the transportation of river water and particles to the ocean. The Ganges and Brahmaputra rivers are especially important for reconstruction of Himalayan monsoon climate. Chemical compositions of the Ganges and Brahmaputra River sediments have been determined to investigate their provenance and source area weathering conditions.

Major and trace element concentrations normalized to upper continental crust (UCC) show the sediments in both rivers are depleted in CaO, K₂O, Na₂O, Sr, MgO, Al₂O₃ - Sc and Fe₂O₃ and enriched in Zr, Th, Ce, SiO₂, Y, TiO₂ and Cr. Silica enrichment is caused by quartz concentration, mineral sorting and/or enhanced compositional maturity during fluvial transportation. Chemical Index of Alteration (CIA) ratios in the river sediments (Ganges, ~64 and Brahmaputra, ~60) are low compared to those of Miocene Sylhet sediments in Bangladesh (mostly ~70-87). CIA ratios are low in the river samples because they mostly contain fresh sandy material within the active channels, while highly weathered fine-grained detritus has been transported by the river waters into the ocean. Major element and trace element ratio plots indicate the sediments in both rivers were derived from felsic continental crust sources. A - CNK - FM relationships further suggest minimal weathering of the modern sediments, major contribution from felsic materials, and minor ferromagnesian mineral content. Chemical similarity of the river sediments, the Sylhet and the equivalent Siwalik sequence sedimentary rocks in Nepal represents homogenization of material derived from the Himalayan source, and no major compositional changes have occurred in the source during the last 20 million years.

Keywords: river, sediments, climate, weathering

BPT002-03

会場:201B

時間:5月25日 09:00-09:15

放射性炭素年代測定を用いた、沖縄県久米島下地原鍾乳洞石筍の年代モデル構築 Age models of stalagmite from Shimochibaru cave in Kume Island, using radiocarbon dating

関有沙^{1*}, 横山 祐典¹, 鈴木 淳², 宮入 陽介¹, 菅 浩伸³, 石村 豊穂², 松崎 浩之¹

Arisa Seki^{1*}, Yusuke Yokoyama¹, Atsushi Suzuki², Yosuke Miyairi¹, Hironobu Kan³, Toyoho Ishimura², Matsuzaki Hiroyuki¹

¹ 東京大学, ² 産業技術総合研究所, ³ 岡山大学

¹The University of Tokyo, ²Geological Survey of Japan, AIST, ³Okayama University

古環境復元のプロキシとしての石筍には、分布地域が限られておらず、世界中の石灰岩地域に石筍が存在すること、千年から10万年スケールで連続的に成長するため、長期記録を得ることができること、そしてウラン系列の年代測定を用いて、60万年前までの正確な年代測定ができることなどの利点があり、近年多くの研究例が報告されている(たとえば Wang et al., 2005)。一般に、放射性同位体を用いた石筍の年代決定は、ウラン系列核種を用いた方法が用いられているが、本研究では、約5万年前までの年代測定が可能であり、少量のサンプルで測定可能な放射性炭素年代測定を用いて石筍の年代モデル構築の可能性について検証を行った。石筍の場合、ambient atmospheric carbon (大気中の同位体比を持つ炭素)のみでなく、母岩である石灰岩からの dead carbon (固定されてから時間が経過し、¹⁴Cに枯渇した炭素)の混入があるため、サンプル中の¹⁴C濃度から直接年代を計算することはできない。しかし、成長期間を通じて dead carbon の混入割合を一定と仮定してその効果を補正することで、成長年代を得ることができる可能性が示唆されている (Beck et al., 2001)。

今回得られた久米島の下地原鍾乳洞から採取した石筍は、X線粉末回折を用いた組成分析によって、純粋な Calcite であることが確認された。久米島下地原鍾乳洞から採取した石筍のうち全長約24cm(石筍A)と約173cm(石筍B)のものを、成長軸に沿ってサンプリングを行い、加速器質量分析装置を用いて放射性炭素年代測定を行った。

放射性炭素年代測定で得られた結果より、Lambert & Aharon (2011)に従って dead carbon の混入割合を計算した。下地原鍾乳洞で採取した3つの石筍の先端部の計算結果から、採取時まで成長していた石筍の先端部でも、約20%の dead carbon の混入があることがわかった。この結果を元に、石筍が成長していた期間、dead carbon の混入割合は常に一定であったと仮定して年代値の補正を行った。

その結果、石筍Aでは、成長方向と年代の逆転が全く見られない年代モデルが得られた。最上部から10cmより深部では、成長速度一定で連続的に成長していたことがわかり、ハイエイタスの存在が示唆される部分については、サンプルの縞の観察と年代モデルが一致した。連続的に成長していた部分の成長速度は0.25m/yrであり、先行研究(例えば Wang et al., 2005)や琉球石灰岩地帯の石筍の成長速度(神谷, 2007)と整合的なものであった。

一方、石筍Bは先端部付近の多少のばらつきはみられたものの、ほぼ一定の成長速度で成長していたことがわかった。成長速度は約1mm/yrであり、Bと比較して4倍の速い値を示した。

今回の結果から、ウラン系列による年代測定を行わなくても、dead carbon の混入割合を適切に補正することにより、放射性炭素年代測定から簡便に石筍の年代モデルを導くことができる可能性が示された。また、年代モデルから成長速度を計算することで、環境復元に適したサンプルであるか否かが判断できることが示唆された。今後はウラン系列各種を測定し、これらの結果の妥当性についても検証していく予定である。

キーワード: 石筍, 放射性炭素年代, 年代測定, 久米島

Keywords: stalagmite, radiocarbon age, dating, Kume Island

BPT002-04

会場:201B

時間:5月25日 09:15-09:30

完新世中期の西太平洋熱帯域表層における海水温および海水の酸素同位体比の復元 Reconstruction of temperature and oxygen isotopes in the surface water of the Mid-Holocene tropical western Pacific

井上 麻夕里^{1*}, Terrence M. Quinn², Frederick W. Taylor², 鈴木 淳³, 石村 豊穂³, 川幡 穂高¹, Hai Cheng⁴, R. Lawrence Edwards⁴

Mayuri Inoue^{1*}, Terrence M. Quinn², Frederick W. Taylor², Atsushi Suzuki³, Toyoho Ishimura³, hodaka kawahata¹, Hai Cheng⁴, R. Lawrence Edwards⁴

¹ 東大・大気海洋研, ² テキサス大学地球物理研究所, ³ 産総研・地質情報研究部門, ⁴ ミネソタ大学地質地球物理学科
¹AORI, the Univ. Tokyo, ²UTIG, the Univ. Texas, ³GSI, AIST, ⁴Dep. Geol & Geophys, Univ Minnesota

Oxygen isotope ratios ($d^{18}O$) in foraminifera from deep sea cores have been used successfully to reconstruct continental ice volume, sea level and deep-ocean temperatures throughout the last glacial-interglacial cycle. Yet there is still considerable debate regarding the relative contributions of changes in the oxygen isotopic composition of seawater in the tropical surface ocean. Obtaining accurate estimate of the past $d^{18}O$ distribution in tropical surface waters is crucial to establishing the role of the tropical oceans in global climate change. In this study, Sr/Ca ratios and $d^{18}O$ values in a fossil coral collected from Vanuatu with the age of 5.2 ka were measured to reconstruct past sea surface temperature (SST) and $d^{18}O$ in seawater. XRD analysis and SEM observation revealed that skeletal material was composed of only aragonite and the age determination was precisely conducted using U-Th dating method. Then Sr/Ca and $d^{18}O$ were analyzed with an approximate time resolution of 2 month. As a result, slightly higher mean SST and about 0.4 per mil enrichment of $d^{18}O$ relative to modern seawater have shown from 5.2 ka coral. The result is corresponding to the previous study from Great Barrier Reef which has suggested that the temperature increase enhanced the evaporative enrichment of ^{18}O in seawater.

Keywords: western Pacific, coral Sr/Ca, oxygen isotope, Holocene

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPT002-05

会場:201B

時間:5月25日 09:30-09:45

中期-後期完新世のスリランカ海水準変動からもたらされる南極氷床変動の情報 Detecting Antarctic melting during the Holocene using sea-level information from Sri Lanka

横山 祐典^{1*}, 宮入 陽介¹, 奥野 淳一², 後藤 和久³, 原口 強⁴, 松崎浩之¹

Yusuke Yokoyama^{1*}, Yosuke Miyairi¹, Jun'ichi Okuno², Kazuhisa Goto³, Tsuyoshi Haraguchi⁴, Hiroyuki Matsuzaki¹

¹ 東京大学, ² 国立極地研究所, ³ 千葉工業大学, ⁴ 大阪市立大

¹University of Tokyo, ²National Institute of Polar Research, ³Chiba Institute of Technology, ⁴Osaka City University

Mid to Late Holocene sea-level change can be used for evaluating long-term stability of the Antarctic ice sheet since the most of the Northern hemisphere major ice sheets disappeared by approximately 8,000 years ago. Ongoing global warming may trigger disintegration of this ice sheet, with collapse of the West Antarctic Ice Sheet alone potentially producing a more than 3 to 4 m global sea-level rise. Relative sea level records from sites far away from former ice sheet regions (far-field) provide information on total volume of the ocean mass change, which can be interpreted as global ice volume change. Here we report Holocene sea-level records from Sri Lanka compared with glacio-hydro-isostatic modeling to better understand the melting history of Antarctic ice sheet during the Holocene.

キーワード: 海水準, 完新世, 南極, アイソスタシー, 堆積物コア, 放射性炭素年代測定

Keywords: Sea Level, Holocene, Antarctica, Isostasy, sediment core, Radiocarbon dating

BPT002-06

会場:201B

時間:5月25日 09:45-10:00

海洋酸性化に対する底生大型有孔虫の石灰化応答 Effects of ocean acidification on calcification of symbiont-bearing reef foraminifers

氷上 愛^{1*}, 藤田和彦², 入江貴博³, 牛江裕行¹, 黒柳あずみ¹, 野尻幸宏⁴, 鈴木 淳⁵, 川幡穂高¹

Mana Hikami^{1*}, Kazuhiro FUJITA², Takahiro IRIE³, Hiroyuki USHIE¹, Azumi KUROYANAGI¹, Yukihiro NOJIRI⁴, Atsushi SUZUKI⁵, Hodaka KAWAHATA¹

¹ 東大・大気海洋研, ² 琉球大学・理, ³ アムステルダム大学, ⁴ 環境研・地球環境研究センター, ⁵ 産総研・地質情報
¹AORI, The Univ. of Tokyo, ²Science, Univ. of the Ryukyus, ³BED, Univ. of Amsterdam, ⁴CGER, NIES, ⁵GSJ, AIST

Ocean acidification (decreases in carbonate ion concentration and pH) in response to rising atmospheric pCO₂ is generally expected to reduce rates of calcification by reef calcifying organisms, with potentially severe implications for coral reef ecosystems. Large, algal symbiont-bearing benthic foraminifers, which are important primary and carbonate producers in coral reefs, produce high-Mg calcite shells, whose solubility can exceed that of aragonite produced by corals, making them the "first responder" in coral reefs to the decreasing carbonate saturation state of seawater. Here we report results of culture experiments performed to assess the effects of ongoing ocean acidification on the calcification of symbiont-bearing reef foraminifers using a high-precision pCO₂ control system. Living clone individuals of three foraminiferal species (*Baculogypsina sphaerulata*, *Calcarina gaudichaudii*, and *Amphisorus hemprichii*) were subjected to seawater at five pCO₂ levels from 260 to 970 ppm. Cultured individuals were maintained for about 12 weeks in an indoor flow-through system under constant water temperature, light intensity, and photoperiod. After the experiments, the shell diameter and weight of each cultured specimen were measured. Net calcification of *Baculogypsina* and *Calcarina*, which secrete a hyaline shell and host diatom symbionts, increased under intermediate levels of pCO₂ (580 and/or 770 ppm) and decreased at a higher pCO₂ level (970 ppm). Net calcification of *Amphisorus*, which secretes a porcelaneous shell and hosts dinoflagellate symbionts, tended to decrease at elevated pCO₂. These different responses among the three species are possibly due to differences in calcification mechanisms (in particular, the specific carbonate species used for calcification) between hyaline and porcelaneous taxa, and to links between calcification by the foraminiferal hosts and photosynthesis by the algal endosymbionts. Our findings suggest that ongoing ocean acidification might favor symbiont-bearing reef foraminifers with hyaline shells at intermediate pCO₂ levels (580 to 770 ppm) but be unfavorable to those with either hyaline or porcelaneous shells at higher pCO₂ levels (near 1000 ppm).

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPT002-07

会場:201B

時間:5月25日 10:00-10:15

Speed of environmental change and ocean acidification Speed of environmental change and ocean acidification

hodaka kawahata^{1*}

hodaka kawahata^{1*}

¹Atmosphere and Ocean Research Institute,

¹Atmosphere and Ocean Research Institute,

Warm tropical ocean is analogy of those in Cretaceous and the Paleocene/Eocene transition. But both showed different features. In the late Albian, the deep sea temperature is estimated at about 15 degree C in the equatorial Pacific and atmospheric CO₂ concentrations is suggested to vary between about 2 and 5 times more than that at present atmospheric pCO₂. Larger oceanic alkalinity content can also be argued from the perspective of the production of calcifying organisms at the surface. Recently Zondervan et al. (2001) pointed out that an acidification of the ocean slows or prevents growth of calcifying primary producers. So the fact that we see some carbonate preservation during the Albian suggests that the ocean surface was sufficiently basic, despite higher atmospheric pCO₂, to allow growth of calcifying producers. In contrast, The P/En is characterized by large excursion in the oxygen and carbon isotope records due to the disintegration of methane hydrate and the most dramatic extinction of 35-50% of cosmopolitan benthic foraminifera in the past 100 m.y.. ODP Leg 199 Site 1220 provides a continuous sedimentary section across the P/E boundary in the carbonate-bearing sediments on 56-57 Ma oceanic crust. The methane is expected to be rapidly oxidized to carbon dioxide. Because of an acidic gas, CO₂ will lead to a reduction in deep-sea pH. A pH decrease was very likely responsible for the emergence of agglutinated foraminiferal fauna as calcareous fauna was eliminated by acidification at the P/E transition at Site 1220.

キーワード: Speed, nvironmental change, ocean acidification

Keywords: Speed, nvironmental change, ocean acidification

BPT002-08

会場:201B

時間:5月25日 10:45-11:15

Subtropical coral records of Northern Hemisphere climate variability and abrupt surface ocean salinity changes

Subtropical coral records of Northern Hemisphere climate variability and abrupt surface ocean salinity changes

Thomas Felis^{1*}, Atsushi Suzuki², Norel Rimbu³, Martin Koelling¹, Henning Kuhnert¹, Mihai Dima³, Gerrit Lohmann³, hodaka kawahata⁴

Thomas Felis^{1*}, Atsushi Suzuki², Norel Rimbu³, Martin Koelling¹, Henning Kuhnert¹, Mihai Dima³, Gerrit Lohmann³, hodaka kawahata⁴

¹MARUM, University of Bremen, Germany, ²GSJ, AIST, Tsukuba, Japan, ³AWI, Bremerhaven, Germany, ⁴AORI, The University of Tokyo, Japan

¹MARUM, University of Bremen, Germany, ²GSJ, AIST, Tsukuba, Japan, ³AWI, Bremerhaven, Germany, ⁴AORI, The University of Tokyo, Japan

Subtropical corals provide an ultra-high resolution archive of past temperature and salinity variations at the sea surface. In contrast to tropical corals, which primarily document interannual variability associated with the El Niño-Southern Oscillation, subtropical coral records document mid- to high-latitude atmosphere-ocean variability. Due to their sub-seasonal resolution, these coral records allow to reconstruct climate variability for individual seasons such as boreal winter, which is still underrepresented in high-resolution proxy reconstructions of the last millennium. The northern Red Sea and the western North Pacific Ocean represent unique locations at the western and eastern margins of the Asian continent, where ocean currents transport warm tropical waters to higher latitudes, enabling coral reef growth at subtropical latitudes. Coral oxygen isotope records in the northern Red Sea (28-29 deg N) were shown to reflect a combined signal of aridity and temperature variations and document atmospheric variability of the Arctic Oscillation/North Atlantic Oscillation (AO/NAO). Coral Sr/Ca and U/Ca records in the western subtropical North Pacific Ocean (27 deg N) were shown to reflect temperature variations and document the oceanic variability of the Pacific (inter)Decadal Oscillation (PDO). Combined analysis of oxygen isotope and Sr/Ca (U/Ca) ratios in corals enables to reconstruct past changes in salinity at annual or higher resolution. Surface ocean salinity is a major component in climate dynamics. However, continuous salinity observations in the surface ocean are scarce prior to 1970, and the magnitude of salinity changes during the last centuries is largely unknown. A coral record from the western subtropical North Pacific indicates that an abrupt regime shift toward fresher surface ocean conditions in this region occurred during the early 20th century, between 1905 and 1910 A.D.. This abrupt freshening resulted from a combination of atmospheric and oceanic advection processes, including a weakening of the westerlies that transport dry continental air from Asia to the North Pacific. New coral Sr/Ca data from the northern Red Sea, in combination with oxygen isotopes, reveal that an abrupt regime shift toward fresher surface ocean conditions occurred in this region at the end of the Little Ice Age, between about 1850 and 1855 A.D.. Possible mechanisms for this abrupt freshening in the northern Red Sea, such as the re-organization of the Northern Hemisphere atmospheric circulation at the end of the Little Ice Age, will be discussed.

キーワード: coral paleoclimatology, oxygen isotopes, trace elements, Northern Hemisphere, Last centuries, Porites corals
Keywords: coral paleoclimatology, oxygen isotopes, trace elements, Northern Hemisphere, Last centuries, Porites corals

BPT002-09

会場:201B

時間:5月25日 11:15-11:30

タヒチサンゴ化石から推定される南太平洋熱帯域の海洋環境 Tropical South Pacific Climate Variability Inferred from Tahiti Fossil Corals: IODP Expedition 310 -Tahiti Sea Level-

浅海 竜司^{1*}, Thomas Felis², Pierre Deschamps³, 花輪 公雄⁴, 井龍 康文⁵, Edouard Bard³, Nicolas Durand³, 村山 雅史⁶
Ryuji Asami^{1*}, Thomas Felis², Pierre Deschamps³, Kimio Hanawa⁴, Yasufumi Iryu⁵, Edouard Bard³, Nicolas Durand³, Masafumi MURAYAMA⁶

¹ 琉球大学・超域, ²University of Bremen, Germany, ³CEREGE, France, ⁴ 東北大学・院理, ⁵ 名古屋大学・院環境, ⁶ 高知大学・理・海洋コア

¹University of the Ryukyus, Japan, ²University of Bremen, Germany, ³CEREGE, France, ⁴Tohoku University, Japan, ⁵Nagoya University, Japan, ⁶Kochi University, Japan

Massive Porites corals, living in shallow waters of the tropical to subtropical oceans, precipitate annually-banded aragonite skeletons. These colonies provide robust chronological control and allow sampling at subseasonal resolution. Ages of fossil corals can be determined accurately by radiocarbon and uranium/thorium dating. Coral strontium/calcium (Sr/Ca) ratio has been widely used as a robust paleothermometer (e.g., Beck et al., 1992). Coral oxygen isotopes (d18O) reflect variations in sea surface temperature and seawater d18O with the latter being closely related to salinity reflecting the precipitation/evaporation balance at the sea surface and changes in water mass transport (e.g., Tudhope et al., 1995). Therefore, coupled determinations of Sr/Ca and d18O in a coral enable the construction of proxy records for both sea surface temperature and salinity (e.g., Gagan et al., 1998, Felis et al., 2009). Fossil corals were shown to provide subseasonally-resolved proxy records of sea surface temperature and salinity for time windows of the Holocene, the last glacial period and older interglacials. However, most fossil reefs of glacial age are located today at water depths of >100 m, as a result of the lower sea level caused by ice sheet build-up during glacial periods (e.g., Bard et al., 1990). Therefore, it has been difficult to collect corals of glacial and deglacial age for paleoclimatic studies.

In 2005, the Integrated Ocean Drilling Program (IODP) Expedition 310 (Tahiti Sea Level), conducted by the European Consortium for Ocean Research Drilling-Science Operator using the mission-specific platform (DP Hunter), drilled the coral reef system off Tahiti (French Polynesia), an island located in the central tropical South Pacific (Camoin et al., 2007). During the expedition, massive fossil coral colonies, mostly of Porites, were recovered in-situ at depths of about 40 to 150 m below modern sea level. We present monthly resolved Sr/Ca and d18O records from well-preserved fossil corals recovered during the expedition (Asami et al., 2009). Our coral-based estimates of sea surface temperature and seawater d18O document thermal and hydrologic variations around Tahiti for selected time windows during the Quaternary, furthering our understanding of tropical South Pacific climate change that accompanied the sea-level change.

Keywords: coral skeleton, Tahiti, oxygen isotope composition, paleothermometer, tropical South Pacific, IODP

BPT002-10

会場:201B

時間:5月25日 11:30-11:45

北西太平洋4島のサンゴが記録する、PDOやEAWM(東アジア冬季モンスーン)に関連した10年規模気候変動 PDO- and East Asia Winter Monsoon-related variability detected in coral records from 4 islands in the North Pacific Gyre

福嶋 彩香^{1*}, 川幡 穂高², 鈴木 淳³, 小島 啓太郎⁴, 岡井 貴司³, 石村 豊穂³, Fernando P. Siringan⁵

Ayaka Fukushima^{1*}, hodaka kawahata², Atsushi Suzuki³, Keitaro Kojima⁴, Takashi Okai³, Toyoho Ishimura³, Fernando P. Siringan⁵

¹ 東大院・理・地惑, ² 東大・大気海洋研, ³ 産総研・地質情報, ⁴ 東大院・工・社会基盤, ⁵ フィリピン大・海洋研
¹EPS, Univ. Tokyo, ²AORI, Univ. Tokyo, ³AIST, ⁴GSE, Univ. Tokyo, ⁵MRI, Univ. Philippines

Decadal variations in the East Asian Monsoon, Pacific Decadal Oscillation (PDO), as well as El Niño Southern Oscillation (ENSO), are considered to play an important role for the regional climate of the NW Pacific. Their interactions may evolve under the progress of the global warming. However, our knowledge on the future climate is limited mostly due to the lack of long-term and precise records. Here we compared the centennial-long $d^{18}O$ and Sr/Ca records in modern *Porites* corals from four sites along the North Pacific Gyre in order to examine past changes in the relationship among the major climate variability. By applying a possible regime shift detection method, it was revealed that evident PDO pattern dominated in the records from the mid-ocean sites, Ogasawara (27°6'N, 142°11'E) and Guam (13°35'N, 144°50'E). In contrast, corals from Ishigaki (12°10'N, 124°20'E) and Bicol (13°03'N, 124°01'E), proximal sites to the continent, indicated strong influence of the East Asian Winter Monsoon, with an evident decreasing trend in its decadal component. Interestingly, both Ishigaki and Bicol records during the 19th century showed significant covariation linked to the PDO, suggesting the PDO as the ruling variability all over the NW Pacific in the past. Weakening of the decadal component of winter monsoon intensity and PDO influence in the marginal sites of the NW Pacific may be associated to impaired Siberian High development during winter while the Aleutian Low remains unchanged under the global warming trend in the 20th century.

キーワード: 太平洋十年規模振動, 東アジア冬季モンスーン, サンゴ骨格気候学, 北西太平洋, レジームシフト

Keywords: Pacific Decadal Oscillation, East Asian Winter Monsoon, coral skeletal climatology, North Pacific Gyre, climatic regime shift

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPT002-11

会場:201B

時間:5月25日 11:45-12:00

喜界島サンゴ骨格を用いた過去432年間の海洋環境復

A 432-year-long paleoceanographic record in Porites coral in Kikai Island, Southern Japan

川久保 友太^{1*}, 横山 祐典¹, 鈴木 淳², シャンテル アルバート³, スティーブ イギンス³
Yuta Kawakubo^{1*}, Yusuke Yokoyama¹, Atsushi Suzuki², Chantal Alibert³, Steve Eggins³

¹ 東京大学大気海洋研究所, ² 独立行政法人産業技術総合研究所, ³ オーストラリア国立大学地球科学研究所

¹AORI, The University of Tokyo, ²Geological Survey of Japan, AIST, ³RSES, The Australian National University

In June 2009, we obtained a long modern core from a coral reef in Kikai Island, Japan. The island is located on the eastern boundary of the East China Sea in the northwestern Pacific. The coral core is approximately 440 cm long and dates back to 432 years (1578-2009 A.D.) including the Little Ice Age (LIA).

The LIA was a time when Europe and other regions neighboring the North Atlantic experienced colder conditions between the 16th to mid-19th centuries. It is reported that this was characterized by the most extensive period of mountain glacier expansion in the recent past. However, owing to the sparseness of the available proxy data during the LIA, there is still no consensus concerning its spatial pattern, timing and cause. In particular, there is a lack of records in the Pacific during the LIA. Thus our 432-year-long coral record from Kikai Island is a useful tool to understand the global-scale picture of climate change during the LIA.

Here we show a continuous 432-year record of sea surface temperature and other parameters based on coral paleo-climate proxies including trace elements, such as Sr, U, and Ba. We analyzed the skeletal elements using laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS). This is a very powerful tool to handle long-term records since it requires a relatively brief experimental time compared with analysis using isotope dilution or thermal ionization ICP-MS. Based on these results, we discuss the paleoceanographic conditions in the northwestern Pacific during the LIA.

キーワード: サンゴ, レーザーアブレーション ICP-MS, 小氷期, 東アジアモンスーン, 太平洋十年規模振動

Keywords: Coral, Laser Ablation ICP-MS, Little Ice Age, East Asian Monsoon, Pacific Decadal Oscillation

BPT002-12

会場:201B

時間:5月25日 12:00-12:15

ハマサンゴ骨格における Mg 同位体分別：飼育実験による低緯度域の新規古気候プロキシの評価

Mg isotope fractionations in *Porites* coral skeletons: evaluation of a new climate proxy from culture experiments

吉村 寿紘^{1*}, 川幡 穂高², 鈴木 淳³, 井上 麻夕里², 谷水 雅治⁴, 古賀 奏子⁵, 酒井 一彦⁵

Toshihiro Yoshimura^{1*}, Hodaka Kawahata², Atsushi Suzuki³, Mayuri Inoue², Masaharu Tanimizu⁴, Kanako Koga⁵, Kazuhiko Sakai⁵

¹ 東京大学・新領域, 大気海洋研究所, ² 東京大学・大気海洋研究所, ³ 産業技術総合研究所・地質情報研究部門, ⁴ 海洋研究開発機構・高知コア, ⁵ 琉球大学・熱帯生物圏研究センター

¹GSFS&AORI, the University of Tokyo, ²AORI, the University of Tokyo, ³IGG, AIST, ⁴KCC, JAMSTEC, ⁵TBRCSS, University of the Ryukyu

Recent developments in inorganic mass spectrometry have allowed new stable isotope systems to be explored by paleoceanographers. In the field of low-latitude climate reconstruction, proxy developments in massive hermatypic corals are still highly desirable for more accurate temperature reconstructions. Stable isotope compositions of the major and minor constituents, such as Ca and Sr, of coral aragonite are potentially new proxies for environmental studies (Bohm et al., 2006; Fietzke and Eisenhauer, 2006). These studies have focused on the temperature dependency of isotope fractionation during the precipitation of biogenic carbonates, because water temperature is the most fundamental parameter controlling the earth's surface environment. Although Sr/Ca is regarded as useful temperature proxy, Mg/Ca is thought to mainly reflect the growth rate (e.g., Inoue et al., 2007). Evaluation of Mg isotope fractionation process in hermatypic coral aragonite must be addressed in order to understand a coral biomineralization and develop a new palaeoceanographic proxy.

Many researchers have analyzed Ca isotopes in biogenic and inorganic calcium carbonates, with the first objective being to evaluate them as a potential new paleothermometer. They have found a small but resolvable temperature dependence of Ca isotope fractionation (e.g., Gussone et al., 2009). Strontium is important minor element in CaCO₃, and in inorganic aragonite and cultured hermatypic corals, it displays similar isotope fractionation mechanisms to those of Ca (Fietzke and Eisenhauer, 2006). Previous studies have reported some steeper temperature dependence (slopes) in Ca and Sr isotope fractionation of biogenic CaCO₃ and the differences in temperature slopes are interpreted as a large kinetic isotopic effects depending on how fast the crystal growing rate and relative mass difference (rmd) among isotopes. Growth effects on Mg isotope are expected to be highly concerned since Mg/Ca is known to clearly dependent on growth rates (Inoue et al., 2007).

In this study, we performed high-precision Mg isotope measurements in *Porites* coral cultured in thermostated tanks. We here evaluate the plausible factors controlling Mg isotope fractionation in coral CaCO₃ skeletons and present the relationships between Mg isotope values and e.g., water temperature, minor elemental concentrations, growth rates.

キーワード: 造礁サンゴ, マグネシウム同位体, MC-ICP-MS, 飼育実験

Keywords: hermatypic coral, magnesium isotope, MC-ICP-MS, culture experiment

BPT002-13

会場:201B

時間:5月25日 12:15-12:30

飼育実験試料を用いたサンゴ骨格気候指標に対する成長速度の影響の評価 Growth-rate influences on coral climate proxies tested by multiple clone culture experiment

林 恵里香², 鈴木 淳^{1*}, 石村豊穂¹, 中村 崇³, 井口 亮⁵, 酒井一彦⁴, 岩瀬晃啓⁴, 川幡 穂高²

Erika Hayashi², Atsushi Suzuki^{1*}, Toyoho Ishimura¹, Takashi Nakamura³, Akira Iguchi⁵, Kazuhiko Sakai⁴, Akihiro Iwase⁴, Hodaka Kawahata²

¹産業技術総合研究所, ²東京大学大学院新領域創成科学研究科, ³琉球大学理学部海洋自然科学科, ⁴琉球大学熱帯生物圏研究センター, ⁵東京大学大気海洋研究所

¹Geological Survey of Japan, AIST, ²The University of Tokyo, ³Faculty of Science, University of the Ry, ⁴Tropical Biosphere Research Center, Univ, ⁵AORI, The University of Tokyo

While coral-based climate reconstruction has been increasingly reported from many tropical sites, potential ambiguity of oxygen isotope thermometer intrinsic in biomineralization process attracts much attention, including so-called "vital effects", "growth-rate-related kinetic effect", "[CO₃²⁻] effect" and so on. Here we study growth-rate influences on skeletal oxygen and carbon isotope ratios based on a long-term culture experiment using *Porites australiensis* clone colonies. The oxygen isotope ratios of colonies showed negligible influence of growth rate over a large variation (2 - 10 mm yr⁻¹) based on the comparison of the seasonal minimum oxygen isotope values during summer. Intercolony variation in the winter extremes can be attributed to growth slowdown / cessation during the cold months. Positive shift in carbon isotope ratios for slower-growing corals was found, and it can be attributed to a kinetic behavior of calcification reaction. Seasonal fluctuation pattern in carbon isotope ratios did not correspond to light intensity. These lines warrant the signal recording ability of coral skeletal oxygen isotope ratios from a long-lived colony of clonal growth as paleo-climate archives, and pose implication for the proper intercomparison of carbon isotope ratio records.

Keywords: coral, oxygen isotope ratio, Sr/Ca ratio, skeleton

BPT002-14

会場:201B

時間:5月25日 12:30-12:45

造礁性サンゴ骨格の窒素同位体比：熱帯・亜熱帯海洋における栄養塩窒素の新指標 Nitrogen isotope of coral skeletons: A new proxy for nitrogenous nutrients in subtropical and tropical ocean

山崎 敦子^{1*}, 渡邊 剛¹, 角皆 潤¹

Atsuko Yamazaki^{1*}, Tsuyoshi Watanabe¹, Urumu Tsunogai¹

¹ 北海道大・理

¹Hokkaido University

Coral skeleton could be a high resolution recorder of past nutrient dynamics in tropical and subtropical ocean with decades to millennia time scales. Nitrogen isotope of organic matter in the coral skeleton could vary with that of nitrogenous sources and be a potential to capture nitrogen isotope changes in marine nitrate. However, nitrogen isotope of coral skeletons had problems to be a paleoenvironmental proxy. Reef corals uptake various nitrogen compounds such as (1) dissolved inorganic nitrogen (DIN) assimilated by zooxanthellae, (2) coral consumption of particle and dissolved organic nitrogen (PON, DON), and (3) nitrogen fixation of symbiotic cyanobacteria. In this study, we developed a new method for coral nitrogen isotope analysis and demonstrated which nitrogen sources controlled nitrogen isotope in coral skeletons. For nitrogen isotope analysis of coral skeletons, each 28 mg powder samples were decalcified. Then, all of organic matter in coral skeleton resolved to nitrate and converted into nitrous oxide to introduce into continuous-flow isotope ratio mass spectrometry. The standard deviation of sample measurements was less than 0.2 permil (1sigma). We compared distribution of coral nitrogen isotope with that of nitrate nitrogen isotope in Shiraho coral reef, Ishigaki Island. We sampled the coral cores from five coral colonies distributed from the mouth of Todoroki River that flow into the coral reef to the reef edge. Seawater samples for nitrate were taken at 50 m intervals along the parallel line of coral sampling. Distribution of coral nitrogen isotope was decreasing from +8.6 permil to +3.0 permil toward off-shore. Nitrogen isotope of nitrate was also decrease from +9.0 permil at river mouth to +2.3 permil at reef edge, which showed similar trends with coral nitrogen isotope. Nitrogen isotope of nitrate near the river mouth was originated from nitrate in river water (nitrogen isotope: +8.2 permil). This result suggested that nitrogen source for reef corals was mainly DIN in reef water, and coral skeleton recorded nitrogen isotope of nitrate. Time series of coral nitrogen isotope has a possibility to reconstruct seasonal to interannual variations of nitrate sources to coral reef such as land-derived nitrogen, nitrogen fixation, and upwelled nitrate. In this presentation, we also introduce application examples of a nitrogen isotope proxy to corals alive in coastal reef and open ocean.

キーワード: 造礁性サンゴ骨格, 窒素同位体比, 硝酸, 陸起源物質の供給, 窒素固定, 湧昇

Keywords: coral skeleton, nitrogen isotope, nitrate, terrestrial input, nitrogen fixation, upwelling