

PEM006-P01

会場:コンベンションホール

時間:5月26日 10:30-13:00

下部電離圏および雷モニタリングのためのアジア VLF/LF 波観測ネットワーク (AVON) システム

Asia VLF/LF wave observation network (AVON) system for monitoring of the lower ionosphere and lightning

大矢 浩代^{1*}, 山下 幸三², 土屋 史紀², 高橋 幸弘³, 塩川 和夫⁴, 三好 由純⁴, 中田 裕之¹

Hiroyo Ohya^{1*}, Kozo Yamashita², Fuminori Tsuchiya², Yukihiro Takahashi³, Kazuo Shiokawa⁴, Yoshizumi Miyoshi⁴, Hiroyuki Nakata¹

¹ 千葉大学大学院工学研究科, ² 東北大学大学院理学研究科, ³ 北海道大学大学院理学研究院, ⁴ 名古屋大学太陽地球環境研究所

¹Graduate School of Eng., Chiba Univ., ²Graduate School of Science, Tohoku Univ., ³Graduate School of Sci., Hokkaido Univ.,

⁴STE Laboratory, Nagoya Univ.

We explain the scientific goals and instrumentations of Asia VLF Observation Network (AVON) system for monitoring the lower ionosphere and lightning. The system consists of three observation sites: Tainan site (23.08N, 120.12E) in Taiwan, Saraburi site (14.53N, 101.03E) in Thailand, and Pontianak site (0.00N, 109.37E) in Indonesia. At each site, we use a monopole antenna and a dipole antenna for the electric field measurements and an orthogonal loop antenna for the magnetic field measurements. The signals detected through these antennas are split into three PCs and used for the monitoring of broadband lightning atmospherics (0.1-40.0 kHz), tweek atmospherics (0.1 ~ 10.0 kHz), and transmitter signals (40.0 and 60.0 kHz etc). Analyzing the VLF/LF data obtained at three sites, we can monitor the lower ionosphere and lightning in Southeast Asia. This network system is utilized in cooperation with other ground-based and satellite-based observation projects to clarify the meteorological aspects of lightning activity and their effects on the middle/upper atmosphere, ionosphere, and magnetosphere. In the presentation, we introduce the AVON system and show the initial results.

PEM006-P02

会場:コンベンションホール

時間:5月26日 10:30-13:00

磁気嵐・サブストーム時における放射線帯電子の大気降下損失

Loss of high-energy electrons into the atmosphere during the magnetic storm and sub-storm

土屋 史紀^{1*}, 森岡 昭¹, 三澤 浩昭¹, 三好 由純², 塩川 和夫², 菊池 崇², 小川 泰信³

Fuminori Tsuchiya^{1*}, Akira Morioka¹, Hiroaki Misawa¹, Yoshizumi Miyoshi², Kazuo Shiokawa², Takashi Kikuchi², Yasunobu Ogawa³

¹ 東北大学, ² 名古屋大学太陽地球環境研究所, ³ 国立極地研究所

¹Tohoku University, ²STEL, Nagoya University, ³NIPR

Man-made VLF/LF radio wave observation at Ny-Alesund in Norway was used to study precipitations of high-energy ($>100\text{keV}$) electrons into the atmosphere during the geomagnetic storm and substorm. The observation system was installed at the NIPR station on March 2010 and measures transmitter signals operated at UK (60.0 kHz) and Germany (77.5 kHz). During three magnetic storms occurred on 5 Apr., 2 May, and 29 May, strong phase variations in the received signals were detected. Comparison of the phase change with the precipitation electron flux observed by the MEPED instrument onboard the NOAA/POSE satellites above the LF wave propagation paths showed the good correlation between them. It is expected that plasma waves excited in the magnetosphere are responsible for the energetic electron precipitation into the atmosphere through the pitch angle scattering. Therefore, local time distribution of the plasma wave would cause the local time dependence of the electron precipitation region. To investigate this process, local time distribution of the electron precipitation events was examined by using the LF wave observation and was compared with the substorm onset timing. At present, the substorm onset timing was determined by the Kakioka Pi2 and the positive bay observed by low-latitude magnetometers located near the local midnight. In the morning and noon sectors, it is found that onset of the precipitation is delayed ten to several tens minutes from the substorm onset. The delay time is consistent with the drift time of energetic electron with energy of $>100\text{keV}$. On the other hand, the electron precipitation onset found in the dusk sector occurred at or just before the substorm onset. This implies that different types of plasma waves are responsible for the dependence of the electron precipitation characteristics on local time.

PEM006-P03

会場:コンベンションホール

時間:5月26日 10:30-13:00

全球雷雲活動とOLR及びそれらの太陽活動依存性 Global thunderstorm activities and OLR, and their dependence on solar cycle

高橋 幸弘^{1*}, 佐藤 光輝¹, 宮原 ひろ子², 星野 直哉³, 山下 幸三¹

Yukihiro Takahashi^{1*}, Mitsuteru Sato¹, Hiroko Miyahara², Naoya Hoshino³, Kozo Yamashita¹

¹ 北海道大学 宇宙理学専攻, ² 東京大学 宇宙線研究所, ³ 東北大学 地球物理学専攻

¹Hokkaido University, ²University of Tokyo, ³Tohoku University

Global ELF observation network, GEON, constructed and operated by Hokkaido University, provides information of each cloud to ground lightning discharge, CG, that is, GPS time, location and energy, as well as Schumann resonance, SR, power, a proxy of global energy proxy of lightning discharge. From the standpoint of the relationship between the effect of solar activity to the climate, lightning activity estimated from the ELF measurement in the frequency range between 1 and 100 Hz and the outgoing longwave Radiation, OLR, an indicator of cloud amount, are examined for their periodicity in the periodic range of about one month. SR power shows about 27 day periodicity in solar maximum years and it becomes elongated toward solar minimum. On the other hand, OLR shows same kind of 27 day periodicity in solar maximum years, but only in the Western Pacific Warm Pool area. Both the spectra of SR and OLR have a peak around 35 days in solar minimum years. The average spectrum of OLR in solar maximum years also shows an enhancement in the range of 50 or 60 days corresponding to the main MJO period. In this paper the relationship between the global lightning distribution based on GEON measurement and OLR are discussed in detail, considering one-month periodicity. Especially synchronization of thunderstorm activity between different longitudes is focused.

キーワード: 太陽活動, 気候, 雷放電, OLR, ELF

Keywords: solar cycle, climate, lightning, thunderstorm, OLR, ELF

PEM006-P04

会場:コンベンションホール

時間:5月26日 10:30-13:00

電離圏加熱現象が極域下部熱圏風へ与える影響 Ionospheric heating effects on the polar lower thermospheric wind

津田 卓雄^{1*}, 野澤 悟徳¹, 大山 伸一郎¹, 小川 泰信², 藤井 良一¹

Takuo Tsuda^{1*}, Satonori Nozawa¹, Shin-ichiro Oyama¹, Yasunobu Ogawa², Ryoichi Fujii¹

¹ 名古屋大学 太陽地球環境研究所, ² 極地研究所

¹STEL, Nagoya Univ., ²NIPR

Of vital importance is to qualify significance of the magnetospheric forcing (such as the Joule heating and the ion drag) to the polar lower thermospheric wind dynamics, in order to obtain better understanding of the Magnetosphere-Ionosphere-Thermosphere (MIT) coupling process. Several measurements by Incoherent Scatter (IS) radars and Fabry-Perot Interferometers (FPIs) demonstrated neutral winds with speeds exceeding 300 m/s in the polar lower thermosphere during geomagnetically active intervals. The wind speeds are significantly larger than a typical wind speed (less than 200 m/s). This suggests that the magnetospheric forcing can accelerate the neutral wind. While a number of observations demonstrated relationships between the neutral wind variations and magnetospheric forcing, there are a few studies on examining quantitatively the forcing on the neutral wind based on observational data. A case study determined contributions of the Joule heating and the ion drag on the acceleration of neutral wind at 118 km, and suggested that the Joule heating was a major important factor. At lower heights (below 110 km), however, the heating effects on the neutral wind dynamics are little known. A couple of studies suggest that an anomalous heating on the atmosphere become important for the wind dynamics in the lower heights due to cooling of the heated electrons induced by the Farley-Buneman instability during the strong electric field.

We have investigated such ionospheric heating effects on the wind dynamics at 100-120 km using data obtained with the European Incoherent SCATter (EISCAT) Svalbard Radar (ESR) located in Longyearbyen (78.2N, 16.0E in geographic coordinates, 75.2 in invariant latitude). As a case study, we have determined contributions of the Joule heating, the ion drag, and also the anomalous heating on the neutral wind acceleration at 100-110 km during a strong electric field. The ESR result shows that electron temperatures in the cusp electrojet reach up to about 4000 K. The heat is transferred to the neutral gas by collisions (i.e., cooling of the heated electrons). This anomalous heating effect can be more important at 101-109 km, compared with that at higher altitude. We have found that the anomalous heating effect at 101 km was comparable to the Joule heating effect and occasionally became much more effective. On the other hand the ion drag contribution became less effective. During the strong electric field, at 101 km, the wind speed increase of 60 m/s was found, while the wind speed increase was decreasing with decreasing altitudes. These results suggest that main contributor to the wind acceleration at 101 km would be a total of the anomalous heating and the Joule heating.

キーワード: EISCAT, 極域, 下部熱圏, 中性風, 電離圏電場, ジュール加熱

Keywords: EISCAT, Polar region, Lower thermosphere, Neutral wind, Ionospheric electric field, Joule heating

PEM006-P05

会場:コンベンションホール

時間:5月26日 10:30-13:00

赤道プラズマバブルに伴う TEC 変動・ロック損失の時空間変動

Spatial and temporal variations of TEC fluctuations and losses of lock associated with equatorial plasma bubbles

菊地 隼人¹, 中田 裕之^{1*}, 津川 卓也², 西岡 未知³, 大塚 雄一³, 鷹野敏明¹, 島倉 信¹, 塩川 和夫³, 小川 忠彦²

Hayato Kikuchi¹, Hiroyuki Nakata^{1*}, Takuya Tsugawa², Michi Nishioka³, Yuichi Otsuka³, Toshiaki Takano¹, Shin Shimakura¹, Kazuo Shiokawa³, Tadahiko Ogawa²

¹ 千葉大学大学院工学研究科, ² 情報通信研究機構, ³ 名古屋大学太陽地球環境研究所

¹ Graduate School of Eng., Chiba Univ., ² NICT, ³ STE Lab., Nagoya Univ.

Equatorial plasma bubbles (EPBs) are depletion of the plasma density in the ionosphere and, inside of EPBs, there are field-aligned irregularities (FAIs) which affects wide-band radio waves. Therefore, EPB causes scintillations on GPS signals because of rapid variations of signal amplitude and phase, and limit the availability of carrier phase measurements. The spatial scale of FAIs that causes the scintillation is determined by Fresnel scale, which is about 2-300 m for GPS signals. This means that loss of phase lock (LOL) on GPS signals is a good proxy for hundred-meter-scale FAIs. It is also widely known that rate of TEC change index (ROTI) enhances around EPBs. Assuming that the altitude of the ionosphere is about 400 km, the velocity of the pierce point of GPS satellites at the ionospheric altitude is approximately 70 m/s around the zenith. ROTI averaged during 5 minutes is a reference of ten-kilometer-scale fluctuations.

In this study, we analyzed LOL and 5-min ROTI to examine the spacial and temporal variations of electron density disturbances associated with EPBs. Examining LOL and ROTI, the developments of two different-scale irregularities are identified. LOL and ROTI data are obtained from GPS data from GPS Earth Observation Network (GEONET) of Japan. From 630-nm airglow images obtained by all-sky imager at Sata, Japan, in 2001, we selected 11 EPBs where the EPBs reach to the geographic latitude of 30 degrees. This is because we compare the distributions of LOL and ROTI determined by GEONET with the airglow imager at Sata whose field of view is 26-34N in geographic latitude.

Both LOL and the enhancement of ROTI were observed in 7 events out of 11 events. The distributions of occurrence of LOLs are approximately in accordance with the depleted region of the airglow intensity, namely the ionospheric electron density. The distribution of the enhancement of ROTI spreads in the vicinities of EPBs, which is wider than that of LOL. The hundred-meter-scale irregularities are distributed within EPBs while the ten-kilometer-scale disturbances are located around EPBs.

In the events associated with LOLs, the kilometer-scale disturbances can be more developed as showing the value of ROTI since there are lacks of observation due to LOL and ROTI would be smaller than the case where there is no lack of observation. In the events without LOL, on the other hand, the fresnel-scale disturbances are not developed enough to cause the scintillation in GPS receivers. The mean values of ROTI in the events associated with LOLs are larger than those not associated with LOLs. Therefore, the kilometer-scale and hundreds-meter-scale disturbances tend to grow and decay simultaneously.

Keywords: Ionosphere, plasma bubble, GPS, loss of lock, TEC, scintillation

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



PEM006-P06

会場:コンベンションホール

時間:5月26日 10:30-13:00

Study of equatorial Spread-F with GNU Radio Beacon Receiver (GRBR) network in Asia, Pacific and Africa Study of equatorial Spread-F with GNU Radio Beacon Receiver (GRBR) network in Asia, Pacific and Africa

山本 衛^{1*}, 津川 卓也², 長妻 努², 大塚 雄一³, Roland Tsunoda⁴, Sri Kaloka⁵, Le Truong Thanh⁶, Ha Duyen Chau⁶, Pornchai Supnithi⁷, Paul Baki⁸

Mamoru Yamamoto^{1*}, Takuya Tsugawa², Tsutomu Nagatsuma², Yuichi Otsuka³, Roland Tsunoda⁴, Sri Kaloka⁵, Le Truong Thanh⁶, Ha Duyen Chau⁶, Pornchai Supnithi⁷, Paul Baki⁸

¹RISH, Kyoto University, ²NICT, ³STEL, Nagoya University, ⁴SRI International, ⁵LAPAN, Indonesia, ⁶HIG, Vietnam, ⁷KMITL, Thailand, ⁸KPUC, Kenya

¹RISH, Kyoto University, ²NICT, ³STEL, Nagoya University, ⁴SRI International, ⁵LAPAN, Indonesia, ⁶HIG, Vietnam, ⁷KMITL, Thailand, ⁸KPUC, Kenya

Equatorial spread F (ESF) is intense ionospheric irregularity that occurs around the geomagnetic equator. It can cause intense scintillation to satellite-ground communications, and serious error in the GPS measurements. The ESF has been a hot research topic of the equatorial/low-latitude ionosphere for long time. However, its day-to-day variability is not well understood. In the southeast Asian region, Japanese researchers developed a network of ground-based observations with the Equatorial Atmosphere Radar (EAR) of RISH, Kyoto University, the ionosonde network SEALION (SouthEast Asia Low-latitude IOnospheric Network) of NICT, and optical instrument network OMTI (Optical Mesosphere Thermosphere Imager) of STEL, Nagoya University. SRI International deploys a VHF radar, an ionosonde and several satellite beacon receivers on Pacific islands. In addition to these, we are deploying the digital satellite beacon receivers named "GNU Radio Beacon Receiver (GRBR)" to fulfill observation gaps. The GRBR-TEC with C/NOFS successfully shows longitudinal large-scale wave structure that is in good relationship to the ESF occurrence. In 2010 we further expanded the network in Asia, Pacific, and African regions. In presentation, we review current status of the wide network of GRBR, and achievement from the observations.

キーワード: 赤道スプレッドF, 衛星ビーコン観測, 電離圏全電子数

Keywords: Equatorial Spread-F, Satellite beacon experiment, Total electron content

PEM006-P07

会場:コンベンションホール

時間:5月26日 10:30-13:00

プラズマバブルに伴う電離圏不規則構造の衛星航法に対する影響とその発生の日々変動に関する研究

Study of impacts of ionospheric irregularities associated with plasma bubbles on GNSS and its day-to-day variability

斎藤 享^{1*}, 松永 圭左¹, 吉原 貴之¹, 星野尾 一明¹, 坂井 丈泰¹, 大塚 雄一²

Susumu Saito^{1*}, Keisuke Matsunaga¹, Takayuki Yoshihara¹, Kazuaki Hoshino¹, Takeyasu Sakai¹, Yuichi Otsuka²

¹ 電子航法研究所, ² 名古屋大学太陽地球環境研究所

¹Electronic Navigation Research Institute, ²STEL, Nagoya University

GNSS (Global navigation satellite system) has been widely used for various applications. Ionospheric irregularities are one of the most serious issues that prevent advanced use of GNSS. Plasma bubbles are among those irregularities that have a great impact on GNSS. Characterization of the ionospheric irregularities are demanded by GNSS applications to realize the high level of services with reliability.

For differential GNSS applications, spatial variability of the total electron content (TEC) is most important, because it directly leads to positioning errors and threatens safety. However, it has not been well studied. Scintillation of GNSS signals due to small-scale irregularities is another aspect of plasma bubble's impact on GNSS degrading the quality of signals.

Indeed, importance of the characterization of the ionospheric irregularities in the low latitude regions has been recognized in the GNSS community, especially in the field of air navigation that requires extremely high level of safety.

Electronic Navigation Research Institute (ENRI) is contributing to International Civil Aviation Organization (ICAO) to provide ionospheric information needed by their own applications. ENRI's activities on this issue are (1) ionospheric data (TEC, TEC gradient, and scintillation) collection in the low latitude regions and (2) leading coordination of the ionospheric data collection in the Asia-Pacific region collaborating with ICAO.

At the meeting, these activities of ENRI including the plan of observation network will be presented more in detail. We believe that these studies should be an important part of the CAWSES-II or ISWI activities and that more attentions should be paid to.

キーワード: 電離圏, プラズマバブル, 観測ネットワーク, 衛星航法, 宇宙天気利用

Keywords: ionosphere, plasma bubble, observation network, GNSS, space weather application

PEM006-P08

会場:コンベンションホール

時間:5月26日 10:30-13:00

タスマニア-ニュージーランド地域における新しい地磁気観測計画：初期解析結果について

A New Project for Constructing a Magnetometer Array in Tasmania and New Zealand: A Preliminary Result

尾花 由紀^{1*}, 塩川 和夫², 吉川 顕正³, 田中 良昌⁴, 才田 聡子⁵, Frederick. W. Menk⁶, Colin. L. Waters⁶, Brian. J. Fraser⁶, Craig. J. Rodger⁷

Yuki Obana^{1*}, Kazuo Shiokawa², Akimasa Yoshikawa³, Yoshimasa Tanaka⁴, Satoko Saita⁵, Frederick. W. Menk⁶, Colin. L. Waters⁶, Brian. J. Fraser⁶, Craig. J. Rodger⁷

¹ 大阪電気通信大学工学部基礎理工学科, ² 名古屋大学太陽地球環境研究所, ³ 九州大学理学研究院地球惑星科学部門, ⁴ 国立極地研究所, ⁵ 新領域融合研究センター, ⁶ The University of Newcastle, ⁷ The University of Otago

¹ Osaka Electro-Communication University, ² Nagoya University, ³ Kyushu University, ⁴ National Institute of Polar Research, ⁵ The Institute of Statistical Mathematics, ⁶ The University of Newcastle, ⁷ The University of Otago

タスマニア-ニュージーランド地域に計画中の、新しい地磁気観測網について、計画の概要を紹介するとともに、初期観測結果を報告する。

観測計画の主たる目的は、磁力線共鳴振動の 1/4 波長モードおよび高調波モードの空間構造を明らかにすることである。太陽風から流入する電磁気エネルギーは電離圏に投影され、磁気嵐時には磁気圏と電離圏の間で巨大な電流系が形成される。磁気圏-電離圏 (M-I) 結合系における電磁気エネルギーの入射と伝搬の解明は、重要な課題であり、精力的な研究が行われているところである。従来の M-I 結合モデルでは、電離圏がエネルギー散逸を引き起こす境界条件としてのみ取り扱われ、磁気圏以外の電磁的環境の効果は基本的に無視されてきた。しかし 90 年代後半から、電離圏からのエネルギー変換過程へのフィードバックを積極的に評価する動きが進みつつある。これは理論やシミュレーション研究が先行しており、観測サイドからのサポートは必ずしも十分ではない。

磁力線共鳴振動の諸モードは、現象それ自体が未知の興味深い研究対象であると同時に、最新の M-I 結合モデルを検証する上での高感度な指標となりうる現象でもある。たとえば観測で得られた磁力線共鳴振動 1/4 波長モードの発生条件を説明するためには、電離圏から M-I 結合系へのエネルギーフィードバックを組み込んだ磁気圏モデルが必要であると言われている。また、1/4 波長から 1/2 波長へのモード遷移は、電離圏上におけるアルフヴェン波の反射係数が、符号を反転させる事に関連して生じると考えられており、沿磁力線電流の反射に対して実効的なアルフヴェン電導度と、それを与えるプラズマ密度について、世界で初めて具体的な観測値を得ることが期待できる。

我々は 2011 年 2 月にニュージーランドのダニーデン近郊、ミドルマーチに地磁気観測点を開設した。将来的にはさらに観測点を増やし、先行する地磁気観測プロジェクト等との連携により、タスマニア-ニュージーランド地域とその共役点における地磁気多点観測体制を構築する計画である。磁力線共鳴振動現象の三次元的空間構造を解明し、観測データと最新の磁気圏モデルを用いた数値計算結果を比較することで、M-I 結合系における諸問題の解明を目指すものである。

キーワード: 地磁気観測, ULF 波動, 磁力線共鳴振動, 磁気圏-電離圏結合, プラズマ圏, 磁気圏

Keywords: geomagnetic field, ULF wave, field line resonance, magnetosphere-ionosphere coupling, plasmasphere, magnetosphere

PEM006-P09

会場:コンベンションホール

時間:5月26日 10:30-13:00

SuperDARN contributions to CAWSES-II SuperDARN contributions to CAWSES-II

行松 彰^{1*}, 細川 敬祐², 元場 哲郎¹, 佐藤 夏雄¹

Akira Sessai Yukimatu^{1*}, Keisuke Hosokawa², Tetsuo Motoba¹, Natsuo Sato¹

¹ 国立極地研究所, ² 電気通信大学

¹ ROIS/NIPR, ² Univ. Electro-Communications

Super Dual Auroral Radar Network (SuperDARN) is an international collaborative HF-radar network originally designed to obtain global large scale two-dimensional polar ionospheric plasma convection patterns in both hemispheres with a temporal resolution of 1 to 2 minutes since 1995. SuperDARN is a powerful tool to be applied to many scientific issues, which can be used not only to deduce dynamics of global large-scale convection patterns, but also to study dynamics of transient meso-scale phenomena like flux transfer events (FTEs), magnetospheric responses to solar wind dynamic pressure like travelling convection vortices (TCVs) and polar cap boundary or open-closed field line boundary (OCB), to detect reconnection sites and to deduce reconnection rates, to study substorms, storms and phenomena related to subauroral regions like sub-auroral polarisation stream (SAPS), to deduce field aligned electric currents (FACs), to study MHD waves in a variety of frequency ranges, and also to study ionospheric irregularities in D-, E-, and F-regions. Moreover, it can be utilised not only to ionospheric researches but also to neutral atmospheric studies, e.g., on atmospheric waves e.g., traveling ionospheric disturbances (TIDs), tides and gravity waves, deducing neutral winds around mesopause region, and also detecting and studying polar mesospheric summer echoes (PMSEs), etc. These days, the fields-of-view (FOVs) of SuperDARN have been expanded to higher latitude (PolarDARN) and mid-latitude (StormDARN) which covers considerable portions of mid- and polar latitudes of earth's ionosphere in both hemispheres and enables us to address much wider ranges of scientific questions (including inner magnetospheric physics). SuperDARN has extensively evolved successfully and has been extremely productive by strong cooperation and competitions within the community and also by collaborative studies with other ground-based and satellite/rocket observations and theoretical research groups.

We present what SuperDARN has done so far and what could not be done by SuperDARN so far, and what SuperDARN will be able to do by recent technical development, e.g., by increasing spatial and temporal resolution and combining with other ground based and satellite observations, and then, discuss how SuperDARN can contribute to CAWSES-II program in terms of the main CAWSES-II themes, especially on the effect of short-term solar variability on the geospace environment (TG3), the geospace response to an altered climate (TG2), and the geospace response to variable inputs from the lower atmosphere.

キーワード: CAWSES-II, SuperDARN, magnetosphere-ionosphere coupling, MLT region dynamics, aurora, neutral winds

Keywords: CAWSES-II, SuperDARN, magnetosphere-ionosphere coupling, MLT region dynamics, aurora, neutral winds

PEM006-P10

会場:コンベンションホール

時間:5月26日 10:30-13:00

昭和基地 - アイスランド共役点におけるオーロラ活動の長期変動 (1) Long-term variation of auroral activity at Syowa-Iceland conjugate stations (1)

門倉 昭^{1*}, 佐藤 夏雄¹

Akira Kadokura^{1*}, Natsuo Sato¹

¹ 国立極地研究所

¹National Institute of Polar Research

Long-term variation of auroral activity at Syowa Station (SYO) (S69.00 deg) in Antarctica and Leirvogur (LRV) (N64.18 deg) in Iceland is investigated. Both SYO and LRV are located at auroral latitudes and in a unique geomagnetically conjugate relationship with each other. Geomagnetic variation data from 1958 and 1966 at LRV and SYO, respectively, are used for this analysis. Using those almost four solar cycle data, similarity and dissimilarity in the solar cycle variation, seasonal variation, and daily variation of geomagnetic activity at those conjugate stations are investigated to understand interhemispheric difference in auroral activity responding to the variation of the solar wind input and solar activity.

キーワード: オーロラ, 地磁気活動, 太陽活動, 長期変動

Keywords: aurora, magnetic activity, solar activity, long-term variation

PEM006-P11

会場:コンベンションホール

時間:5月26日 10:30-13:00

Altitudinal response of global ionosphere to short-period recurrent geomagnetic activity during extreme solar minimum Altitudinal response of global ionosphere to short-period recurrent geomagnetic activity during extreme solar minimum

Tulasi Ram Sudarsanam^{1*}, M. Yamamoto¹, C. H. Liu², H. Liu¹, S.Y. Su³, J. Lei⁴
Tulasi Ram Sudarsanam^{1*}, M. Yamamoto¹, C. H. Liu², H. Liu¹, S.Y. Su³, J. Lei⁴

¹RISH, Kyoto University, Kyoto, Japan, ²Academia Sinica, Taipei, Taiwan, ³National Central University, Taiwan, ⁴University of Colorado, Boulder, USA

¹RISH, Kyoto University, Kyoto, Japan, ²Academia Sinica, Taipei, Taiwan, ³National Central University, Taiwan, ⁴University of Colorado, Boulder, USA

The deep solar minimum of solar cycles 23/24 was exceptionally quiet, with sunspot numbers at their lowest in at least 75 years. During this unique solar minimum epoch, however, solar wind high speed streams emanating from near-equatorial coronal holes occurred frequently and are the primary contributor to the continuous geomagnetic activity at the Earth. These conditions enable the isolation of forcing by geomagnetic activity on the preconditioned solar minimum state of the upper atmosphere caused by Corotating Interaction Regions (CIRs). Global observations of vertical electron density profiles by Formosat3/COSMIC provided a unique opportunity to study the altitudinal response of global ionosphere to this recurrent geomagnetic force caused by CIRs during the extreme solar minimum. The results indicate that the topside ionospheric response (above 350 km) appears to be dominated by changes in the plasma temperature and/or scale height and exhibits concurrent enhancements with the oscillations in geomagnetic activity during both day and nighttime. However, the electron density response at altitudes between 200 and 350 km is dominated by changes in the neutral composition and exhibits significant latitudinal, local time, and seasonal variations. The results are discussed in light of equatorward wind perturbations during enhanced geomagnetic activity and summer to winter transequatorial neutral wind patterns.

キーワード: Co-rotating Interaction Regions, recurrent geomagnetic activity, ionosphere, solar minimum
Keywords: Co-rotating Interaction Regions, recurrent geomagnetic activity, ionosphere, solar minimum