

2020年代の木星系探査計画 (EJSM) Towards EJSM (Europa Jupiter System Mission) in 2020s

佐々木 晶^{1*}, 藤本 正樹², 高島 健², 矢野 創², 笠羽 康正³, 木村 淳⁴, 高橋 幸弘⁴

Sho Sasaki^{1*}, Masaki Fujimoto², Takeshi Takashima², Hajime Yano², Yasumasa Kasaba³, Jun Kimura⁴, Yukihiro Takahashi⁴

¹ 国立天文台 R I S E 月探査プロジェクト, ² 宇宙航空研究開発機構宇宙科学研究所, ³ 東北大学, ⁴ 北海道大学

¹National Astronomical Observatory, ²ISAS/JAXA, ³Tohoku University, ⁴Hokkaido University

国際共同木星探査計画は、2020年代の木星探査を目指して、2006年より日欧のグループが、検討をはじめてきた。2007年に LAPLACE という名称で ESA の COSMIC VISION に提案され、1次選考を通過した。その後、アメリカ NASA で計画していた Europa Explorer 計画と合同する形で、EJSM (Europa Jupiter System Mission) へと展開している。EJSM においては、NASA が JEO (Jupiter Europa Orbiter) (エウロパ周回機)、ESA が JGO (Jupiter Ganymede Orbiter) (ガニメデ周回機)、JAXA が JMO (Jupiter Magnetospheric Orbiter) (磁気圏観測機) を担当する。JEO と JGO は、2年間、木星の周囲を (イオヤカリストを含む) 衛星フライバイを繰り返して観測した後、エウロパ、ガニメデの周回軌道に入る。両衛星では、地下海の存在が確実視されており、生命の存在まで議論されている。レーダーなどで地下海の姿を明らかにすることが大きな目的である。現在もカッシーニが質の高い大量のデータを提供する土星系と比べると、木星とその衛星系の科学データは限定されている、ガリレオ探査機が周回機として長期間観測を行ったが、アンテナや観測機器のトラブルにより取得データが少なかった。木星極周回軌道により木星大気や重力を観測する JUNO 計画では、衛星や木星磁気圏の詳細探査は行われない。

現在、NASA, ESA は探査機を 2020 年に打ち上げて 2026 年に木星周回軌道に投入、2028 年に衛星周回ミッションへと移行する予定である。日本の JMO は、おそくとも 2020 年に打ち上げることができれば、2028 年に木星周回軌道に投入ができ、JEO-JGO の衛星周回時に外側の磁気圏を観測して重要な境界条件を与えることができる。また、軌道計画の検討の結果、カリストによる重力スイングバイを繰り返すことで、JMO の軌道傾斜角を上げて、木星本体のイメージングを行えることがわかった。このときに、カリストの表面、重力、磁場といった情報も詳しく調べることが可能である。カリストは、エウロパ、ガニメデといった天体と比べると未分化であると考えられている。慣性率、磁場を正確に求めることで、内部の分化度や地下海の存否を解明することが可能である。

JAXA では、ソーラー電力セイルを用いて探査機を木星経由トロヤ群小惑星へと送り込む探査計画が検討されている。これに相乗りする形で、木星に行くミッションに、JMO を搭載する検討も行っている。トロヤ群小惑星も、太陽系・木星系の起源を論じる上で非常に重要なターゲットであり、トロヤ群小惑星を組み合わせることで、木星系探査の科学価値は高くなる。ソーラー電力セイルは、太陽電池薄膜で発電してイオンエンジンで推進する機構である。2010 年には、金星ミッション「あかつき」に相乗りの形で、イカロス (Interplanetary Kite-craft Accelerated by Radiation Of the Sun) という技術実証衛星が行われた。

キーワード: 木星磁気圏, 木星探査, 氷衛星, トロヤ群小惑星, エウロパ, カリスト

Keywords: Jovian Magnetosphere, Exploration of Jupiter, Icy satellites, Trojan asteroids, Europa, Callisto

Proposal for the survey of Enceladus by high energy neutrinos Proposal for the survey of Enceladus by high energy neutrinos

庄司 大悟^{1*}, 栗田 敬¹, 田中 宏幸¹
Daigo Shoji^{1*}, Kei Kurita¹, Hiroyuki Tanaka¹

¹ 東京大学 地震研究所

¹ Earthquake Research Institute

Enceladus is a small icy satellite of Saturn orbiting between Mimas and Tethys. In 2005 Cassini has identified characteristic surface features at the south pole of Enceladus such as high albedo and paralleled lineaments called “ tiger stripes ”, which emanate vapor plume [1]. These features suggest that Enceladus has young and active surface around the south pole region. On the other hand, at the north pole, Enceladus has surface whose albedo is lower than the south pole. Such asymmetry of the surface has aroused strong interests on the internal structure as well as its evolution as the origin.

To investigate the surface layer of Enceladus, electrical conductivity is an important information to constraint for the internal structure. Electrical conductivity depends on primarily on temperature [2]. If we can determine the value of conductivity, we can specify the property of ice such as temperature distribution of icy layer. In this presentation we propose a new method to determine the electrical conductivity of ice layer by detecting the radio waves induced by interaction between cosmic neutrino and ice.

When cosmic neutrinos flying in the outer space traverse through Enceladus, Cherenkov radiation induced by the weak interaction of neutrino with Enceladus is emitted. Radiations whose frequency is between a few hundreds of MHz and a few GHz (radio wave) become coherent and have such strong intensity that orbiting probe can detect. The number of detectable emissions depends on attenuation level of radio wave. The attenuation level of water ice can be approximated as $A=0.0009s$ (dB/m) where A and s are attenuation level and electrical conductivity (in μ S/m) respectively [3]. Thus, if we can count the number of emissions and determine their intensity level, electrical conductivity and temperature can be estimated. Since radio waves induced by neutrino interaction come from subsurface area of icy layer (~10 km in depth) local temperature distribution can be obtained by latitude-dependent summation for the emission in polar orbits.

To evaluate this method, we have performed a simulation about interaction of neutrinos with the icy layer and obtained that the number of detectable number of emissions and the shape of intensity distribution changed with the electrical conductivity of the layer. The strong point of this method is the passive detection of radio wave. Further more, accurate determination of electrical conductivity can make the rader system inspect the inner structure more precisely. We consider this radio detection method can be an useful tool to constrain the subsurface temperature of Enceladus.

References

[1] Porco et al. (2006), Cassini Observes the Active South Pole of Enceladus, *Science*, 311, 1393-1401

[2] Corr et al. (1993), Radar absorption due to impurities in Antarctic ice, *Geophys. Res. Lett.*, 20, 1071-1074

[3] Moore (2000), Model of Rader Absorption in European Ice, *Icarus*, 147, 292-300

Keywords: Enceladus, neutrino, electrical conductivity, temperature, passive detection

PPS001-P03

会場:コンベンションホール

時間:5月27日 14:00-16:30

Whistler-mode chorus enhancements and anisotropic electrons in the Jovian inner magnetosphere

Whistler-mode chorus enhancements and anisotropic electrons in the Jovian inner magnetosphere

加藤 雄人^{1*}, 土屋 史紀², 三好 由純³, 森岡 昭², 三澤 浩昭², 氏家 亮², W. S. Kurth⁴, A. T. Tomas⁵, N. Krupp⁶
Yuto Katoh^{1*}, Fuminori Tsuchiya², Yoshizumi Miyoshi³, Akira Morioka², Hiroaki Misawa², Ryo Ujiie², W. S. Kurth⁴, A. T. Tomas⁵, N. Krupp⁶

¹ 東北大学・理・地球物理学, ² 東北大学・理・惑星プラズマ, ³ 名古屋大学太陽地球環境研究所, ⁴University of Iowa, ⁵GeoForschungsZentrum Potsdam, ⁶Max Planck Institute
¹STPP, Grad. Sch. Sci, Tohoku Univ., ²PPARC, Grad. Sch. Sci., Tohoku Univ., ³STEL, Nagoya Univ., ⁴University of Iowa, ⁵GeoForschungsZentrum Potsdam, ⁶Max Planck Institute

We reveal a close relationship between enhancements of whistler-mode chorus and development of energetic electron anisotropies in the Jovian inner magnetosphere by conducting a statistical survey of both wave and particle observations of the Galileo spacecraft. We studied the spatial distribution of intense chorus emissions in the Jovian magnetosphere and identified 104 chorus enhancements by analyzing plasma wave data in the frequency range from 5.6 Hz to 20 kHz obtained from the entire Galileo mission in the inner Jovian magnetosphere during the time period from December 1995 to September 2003. Enhanced chorus emissions with integrated wave power over 10^{-9} (V/m)^2 were observed around the magnetic equator in the radial distance range from 6 to 13 R_J. A survey of energetic particle data in the energy range of 29 to 42 keV reveals that all of the identified chorus events were observed in the region of pancake pitch angle distributions of energetic electrons. Using empirical plasma and magnetic field models, we estimate that the ratio of the electron plasma frequency to the electron cyclotron frequency in this region is in the range from 1 to 10 which is suitable for efficient whistler-mode wave generation. The present study reveals the statistically significant correspondence between intense chorus and flux enhancement of energetic electrons having pancake pitch angle distributions in the Jovian magnetosphere.

Keywords: Jovian inner magnetosphere, whistler-mode waves, energetic electrons, Galileo observations

PPS001-P04

会場:コンベンションホール

時間:5月27日 14:00-16:30

InSb 赤外撮像装置電気回路系の開発 Development of the electronics for an infrared camera with InSb array

北見 拓也^{1*}, 坂野井 健¹, 宇野 健¹, 市川 隆¹, 笠羽 康正¹

Takuya Kitami^{1*}, Takeshi Sakanoi¹, Takeru Uno¹, Takashi Ichikawa¹, Yasumasa Kasaba¹

¹ 東北大学大学院・理学研究科

¹ Graduate School of Science, Tohoku Univ

We present the current status of development of the electronics on our infrared imaging camera with InSb 256x256 array. Infrared remote sensing of planetary atmosphere is one of the most powerful measurement tools to understand the dynamical and chemical processes in the atmosphere since there are many emission and absorption lines in the near-infrared range (1-5 μm), and the solar flux becomes smaller compared to visible range. Further, it is essential to carry out continuous measurement with our own instrument since it is necessary to clarify the time variation of those phenomena with long-term data. In particular, we aim to clarify the Jupiter's H_3^+ auroral response to solar wind variation with statistical approach. We are therefore developing our own 1-5 μm infrared

imager. This imager has a 256x256 InSb array detector, a field of view is 110arcsec with a F12 telescope with a plate scale of 0.43arcsec/pixel. In the case of 3.4 μm Jovian H_3^+ auroral measurement, we estimate S/N of the acquisition of data to be about 33 with 1 minute exposure using 60cm/F12? telescope.

In this presentation, we focus on the electronics to control the detector.

Functions of the electronics are summarized as follows. [1] Generating the timing of three kinds of clocks, selecting horizontal vertical lines and reading the frame of the detector. [2] Converting it into the voltage that adjusted an above clock timing to the detector. [3] Constant voltage (Bias) generation . [4] The amplification of the detector output, and A/D conversion.

To satisfy the function [1] we adopt the digital circuit system with FPGA (Field-Programmable Gate Array) and one-board computer which had a characteristic of incorporated Linux. Concerning on the functions [2] [3] [4], the analog circuit system are used .

This camera will be installed on the 60 cm telescope of Iitate observatory, Tohoku University, and other overseas facilities, and used to monitor the Jupiter's H_3^+ aurora.

PPS001-P05

会場:コンベンションホール

時間:5月27日 14:00-16:30

ガニメデ磁気圏の非MHD的特徴について：ガリレオ探査機データに基づく極域波動 - 粒子相互作用の調査

Non-MHD Aspects of Ganymede's Magnetosphere: Investigation of Polar Wave-Particle Interaction Based on Galileo's data

木村 智樹^{1*}, 笠原 慧¹, 藤本 正樹¹

Tomoki Kimura^{1*}, Satoshi Kasahara¹, Masaki Fujimoto¹

¹ 宇宙航空研究開発機構 宇宙科学研究所

¹JAXA/ISAS

Jovian satellite Ganymede has small magnetosphere with characteristic scale lengths comparable to those of Mercury: e.g., size of the solid body, spatial expansion of the magnetosphere, and electron/ion gyroradii of ambient plasma. Comparative study of Ganymede's magnetosphere with Mercury will provide insights on the process universally existing in small planetary magnetospheres.

Basic characteristics of Ganymede's magnetosphere were revealed based on in-situ measurements by Galileo spacecraft during six encounters. Williams et al. (1997a, b, 1998, 2001, 2004) investigated particle dynamics (e.g., loss and pitch angle diffusion) in Ganymede's magnetosphere based on energetic electron/ion observations. Gurnett et al. (1996) and Kurth et al. (1997) indicated that Ganymede's magnetosphere is emitting radio and local plasma waves similar to planetary magnetospheres. Recently, global configuration of the magnetosphere and interaction with Jovian magnetosphere are also intensively investigated based on MHD simulations (Jia et al., 2009, 2010). However, non-MHD characteristics of Ganymede's magnetosphere have not been discussed in detail yet. For example, wave-particle interactions, ion kinetics, and polar field aligned particle accelerations.

This study addresses wave-particle interaction process in the polar cap region based on multi-instrumental observations during Galileo G02 flyby. Observations of high and low frequency wave, particle energy spectra, and pitch angle distribution revealed two kinds of magnetospheric regions: one where strong particle anisotropy by satellite surface loss is accompanied by electron and ion-related electromagnetic waves, and the other where there are weak surface interactions with electrostatic electron wave and no ion-related waves. The latter region corresponds to the ion upflow region (Jia et al., 2009) and locates near the open-closed boundary region of Ganymede's magnetosphere. We found that ion-related low frequency waves have significant energy flux into the Ganymede's polar ionosphere which is comparable to Jovian magnetospheric electron's energy input. This suggests the polar ionospheric heating by the energy from ion-related waves and subsequent ion upflow.

キーワード: 木星, ガニメデ, 磁気圏, 波動, 高エネルギー粒子

Keywords: Jupiter, Ganymede, magnetosphere, wave, energetic particle

PPS001-P06

会場:コンベンションホール

時間:5月27日 14:00-16:30

木星・土星の赤外オーロラ発光モデリングと観測データ解析への適用 Modeling of infrared auroral emission from Jupiter and Saturn and its applicability for observation data analysis

埜 千尋^{1*}, Sarah V. Badman¹, 宇野 健², 藤本 正樹¹
Chihiro Tao^{1*}, Sarah V. Badman¹, Takeru Uno², Masaki Fujimoto¹

¹ISAS/JAXA, ²東北大学・理・地球物理

¹ISAS/JAXA, ²Tohoku Univ.

Aurorae represent plasma environments around a planet. Outer planetary aurorae are observed in various wavelength from radio emission, infrared (IR), visible, ultraviolet, to X-ray. Since IR wavelength is observable from the ground with spatial information, it would be good tool for monitoring to investigate the variable environment. Recent observation by Cassini spacecraft provides spatially-resolved Saturn's IR auroral image. Since the IR emission relates with thermally excited H_3^+ ion, it reflects atmospheric temperature in addition to ionization by auroral electron and solar EUV. Previous modeling study relates the IR emission and H_3^+ column density and atmospheric temperature. This study newly attempts to test its applicability for monitoring not only atmospheric condition but also auroral electron. We investigate the dependence of IR emission spectrum on temperature and electron energy spectrum using an IR emission model accounting for ionization by auroral electron with various energy flux, ion chemistry, and H_3^+ non-LTE effects. IR emission increases with increasing electron energy for <10 keV and then decreases. This decrease reflects low temperature at low altitude and hydrocarbons which reduces H_3^+ by dissociative recombination. Emission line ratio varies by a few 10s% depends on electron energy and by a factor depends on atmospheric temperature. We will discuss its applicability to observed data analysis and requirement for observations.

Keywords: Jupiter, Saturn, infrared, aurora, spectrum