

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P01

Room:Convention Hall

Time:May 25 16:30-17:30

A new method for evaluating fault activity based on fault gouge properties -Occurrences and colors of fault gouges from

Yukari Miyashita^{1*}, Kenta Kobayashi², Atsushi Kamei³, Jun'ichi Itoh⁴, Mitsuo Manaka⁵, Keisuke Fukushi⁶

¹Geological Survey of Japan, AIST, ²Niigata University, ³Shimane University, ⁴Geological Survey of Japan, AIST, ⁵Geological Survey of Japan, AIST, ⁶Kanazawa University

In order to develop a new method for evaluating fault activities, we have analyzed a relationship between fault activities and fault gouge properties. In particular, we focused on the colors, clay minerals and chemical compositions of fault gouges.

We report occurrences of fault gouges in the western part of Tottori Prefecture, southwest Japan, particularly attentive for the colors of fault gouges derived from granite. Many minor faults accompanying fault gouges are distributed in the area. The fault gouges from the aftershock area of 2000 Tottori-ken Seibu earthquake are characterized by pale-green, white and yellowish brown. In contrast, gouges from the Nichinanko lineament and Komachi-Ohdani lineament area are characterized by yellowish brown, orange and pink. On the a*-b* diagram, these two area are clearly distinguished. The difference of fault gouge color might be corresponding to the difference of fault activity.

Keywords: fault gouge, fault rocks, 2000 Tottori-ken Seibu earthquake, active fault, fault activity

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P02

Room:Convention Hall

Time:May 25 16:30-17:30

A new method for evaluating fault activity based on fault gouge properties-Comparison of fault gouges in the aftershock

Mitsuo Manaka^{1*}, Keisuke Fukushi², Yukari Miyashita¹, Kenta Kobayashi³, Atsushi Kamei⁴, Jun'ichi Itoh¹

¹AIST, ²Kanazawa University, ³Niigata University, ⁴Shimane University

We compared to examine mineralogical and geochemical studies of fault gouges in the aftershock area and the neighborhood of 2000 Tottori-ken Seibu earthquake, to establish a new method for evaluating fault activity of low activity faults. The fault gouges were conducted by powder X-ray diffraction analysis, sequential selective extraction tests and color measurements. As the results, the fault gouge in the aftershock area is mainly composed of illite and chlorite, and the gouge in the neighborhood is mainly composed of halloysite. Iron in the gouge in the aftershock area is mainly contained in illite, and iron in the gouge in the neighborhood is mainly contained in amorphous and crystalline iron oxide. Results of color measurements showed that negative a^* values from the gouge in the aftershock area indicated the presence of chlorite, and that positive a^* values from the gouge in the neighborhood indicated the presence of crystalline iron oxide. These results indicated that mineralogical and geochemical characteristics can distinguish clearly the fault gouges in the aftershock area and the neighborhood of 2000 Tottori-ken Seibu earthquake and that color measurements can be effective to distinguish these gouges.

Keywords: 2000 Tottori-ken Seibu earthquake, fault gouge, powder X-ray diffraction analysis, sequential selective extraction test, color measurement, crystalline iron oxide

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P03

Room:Convention Hall

Time:May 25 16:30-17:30

A new method for evaluating fault activity based on fault gouge properties - Color measurement of fault gouges -

Kenta Kobayashi^{1*}, Yukari Miyashita², Mitsuo Manaka², Atsushi Kamei³, Keisuke Fukushi⁴, Jun'ichi Itoh²

¹Dep. Geol., Fac. Sci., Niigata Univ., ²AIST, ³Shimane Univ., ⁴Kanazawa Univ.

We measured colors of fault gouge zones to estimate the fault activities. First, we made powder containing hematite and/or goethite which have been weighed correctly, and measured colors of the powder. Second, the colors were compared with those of the fault gouge zones along the Nichinan-ko SE lineament and with those in the epicentral area (aftershock zone) of the 2000 Tottori-ken Seibu earthquake (Mj 7.3). Dark-red gouge zones along the Nichinan-ko SE lineament contain 0.1-0.5 % of hematite. Precipitations of the hematite are observed along the margins of the gouge zones, may have formed in the inter-active period.

Keywords: Tottori Prefecture, fault gouge, color, active fault, fault activity

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P04

Room:Convention Hall

Time:May 25 16:30-17:30

Estimation of causative faults producing crustal upwarping in the Nishi-tsugaru Coast, Northeast Japan

Yuumi Ogata^{1*}, Takahiro Miyauchi¹

¹Earth Sci. Dept., Chiba University

The Nishi-tsugaru Coast, Northeast Japan, has experienced co-seismic shoreline uplifts associated with two historic earthquakes (either M6.9) which occurred in 1704 AD and 1793 AD (Imamura, 1920; Usami, 2003). Although each offshore causative fault model was proposed to explain the height distribution of co-seismically emerged abrasion platforms by Nakata et al. (1976) and the small tsunami generation (Sato, 1980), neither models did not coincide with active tectonic structures and topography. Re-recognizing Holocene emerged tidal topography and their dating, we aim to estimate actual causative faults of which movements match the upwarping of paleoshoreline features and active geologic structure in the surrounding submarine areas. Based on the concordance among the deformation modes indicated by co-seismic records and late Quaternary marine terrace records, the causative fault of 1793 earthquake is assigned to the Kita-kanegasawa fault developing the Odoose anticline described in the geological map by Hirayama and Kamimura (1985), and similarly that of 1704 earthquake is likely the Omagoshi fault in Osawa (1963). In addition, the most northeast area of this coast has been probably up-warped by the active movement of Ajigasawa fault, and the Henashi peninsula in the central part of the coast by that of a submarine active fault which is estimated below the steep tectonic scarp several kilometer offshore

Keywords: emerged shoreline topography, upwarping, coseismic uplift, causative fault, Nishi-tsugaru Coast

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P05

Room:Convention Hall

Time:May 25 16:30-17:30

Offshore source fault modeling using late Quaternary paleoshoreline records, Northeast Japan

Takahiro Miyauchi^{1*}

¹Earth Sci. Dept., Chiba University

The eastern margin of Japan Sea is a tectonically active zone of Northeast Japan back-arc and a fold and thrust belt under compressive stress field in Quaternary, which is characterized by inversion tectonics succeeding to the middle Miocene opening of the sea. The differential uplift showing tilting and warping of late Quaternary marine terraces along a 500 km long coast of this eastern margin is understood due to offshore causative fault movements and related large earthquakes. Holocene terrace morphology with steps, historically documented coastal uplift at large earthquakes along the Japanese coast, and harmonious displacement patterns among MIS 5e and Holocene paleoshoreline heights, indicate the occurrence of intermittent and repeated coastal uplift produced by co-seismic distinctive crustal movements. Shallow submarine reverse fault segments with either west dip or east dip, close to the coastlines, are undoubtedly responsible for coastal uplift and accumulative marine terrace tilting and warping demonstrated as fault-related fold structures. Calculating co-seismic displacement distribution to be best fit to marine terrace records, by the use of crustal dislocation model, individual fault parameters range in length from 20-60 km, in dip angle from 30-50 degrees, in slip from 2-7 m (in uplift from 1-3 m), suggesting earthquake magnitude in the range of Mw7.0-7.5 and recurrence time of 1000-4000 years. This implies that the eastern marginal coastal areas of Japan Sea have high probability of large magnitude earthquakes accompanied by coastal uplift, which will happen somewhere in near future. Particularly in the area where present abrasion platforms widely develop, the next large earthquake seems imminent.

Keywords: Late Quaternary, paleoshoreline records, dislocation, source fault modeling, Japan Sea-eastern marginal tectonic zone

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P06

Room:Convention Hall

Time:May 25 16:30-17:30

Seismic reflection profiling survey across the Tengmori-Dedana Faults, the southern Kitakami lowland fault zone

Kohei Abe^{1*}, Dai Nobuoka², Yoshikazu Matsubara², Kyoko Kagohara³, Hideki Kosaka⁴, Takanobu Kamataki², Toshifumi Imaizumi⁵

¹M. T. Brain Co.,Ltd., ²Oyo Co., ³AFERC, AIIT, ⁴Kankyo Chisitu Co.,Ltd., ⁵Tohoku Univ.

The Kitakami lowland fault zone (KLFZ) is an active thrust zone that extends for about 70 km along the Quaternary volcanic front of the northeast Japan arc. The activity of the thrust zone after Quaternary is mostly recognized as a fault reactivation of Miocene normal faults in the area. Tengmori fault group which comprise the southern portion of the KLFZ is composed by several active faults. These faults deform late Quaternary fluvial terraces and debris flow deposits in Kitakami lowland. We present seismic reflection data acquired along the Kitakami - Kanegasaki profile, 12.8 km-long, to define the geological structure of the Tengmori fault group. In seismic lines, the vibrator truck (IVI ENVIRO VIB) is used as the seismic source. Source and geophone spacing are 10-m. Seismic reflection data was processed by using the standard CMP stacking method. The seismic profiles correlated with surface geologic mapping clearly.

Keywords: Tengmori-Dedana Faults, Kitakami Lowland, seismic reflection profiling, subsurface structure

SSS032-P07

Room:Convention Hall

Time:May 25 16:30-17:30

Fault outcrop and tectonic landform of the western margin fault zone of the Kitakami lowland, northeast Japan

Hideki Kosaka^{1*}, Kohei Abe², Kyoko Kagohara³, atsushi Miwa⁴, Toshifumi Imaizumi⁵, Yutaka Mamada⁶, Jun-ichi Uchida⁶

¹Graduate student, Tohoku Univ., ²M. T. Brain Co.,Ltd., ³AIST, ⁴Oyo. Co., ⁵Tohoku Univ., ⁶JNES

The western margin fault zone of the Kitakami Lowland (KLFZ) is a 70-km-long, reverse faults that consist of multiple faults that run parallel to mountain front of northeast Japan. The older paleoseismic events are difficult to explain by only a single trench investigation. Understanding the distribution and activity of the branched faults is important to identify the older paleoseismic events in the KLFZ. We describe paleoseismic events and distribution and activity of the branched faults based on two fault outcrops and the tectonic landform.

Paleoseismic events are identified in the first fault outcrop of the Uwandaira fault group (UFG). We identified at least 4 late Pleistocene-Holocene paleoseismic events, based on the upward termination of fault and angular unconformity, in the fault outcrop. Studies of tectonic landform of fluvial terraces indicate 8-20 late Pleistocene-Holocene paleoseismic events, assuming slip per event obtained from the fault outcrop and the minimum height of the scarplet. The pronounced disparity of paleoseismic events between the fault outcrop and tectonic landform suggests that multiple events have been recorded beneath each angular unconformity. The results imply that the study of the tectonic landform is indispensable to interpretation of paleoseismic events in the region where unconformities have been formed in the footwall of the active thrust.

Distribution and activity of the branched faults are identified in the second fault outcrop of the Nanshozan active fault group (NFG; F1 fault, F2 fault), based on geomorphic feature and surface geology (Plio-Pleistocene Siwa Formation), including the active reverse fault passing through the second fault outcrop (F3 fault). The Miocene strata have been thrust over the Siwa Formation along the mountain front (F1 fault). Distribution of fluvial terraces indicates that the F1 fault has been inactive. The F2 fault deforms fluvial terraces in the footwall of the F1 fault. Deformations of fluvial terraces and arrangement of valley spread of alluvial terraces provide that the F3 fault runs through east side of the hills on the footwall of the F2 fault.

Keywords: Paleoseismic event, angular unconformity, branched fault

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P08

Room:Convention Hall

Time:May 25 16:30-17:30

Active fault traces along the Hanawa higashi fault zone, northeastern Japan

Masayoshi Tajikara^{1*}, Takashi Nakata², Toshifumi Imaizumi³, Yasutaka Ikeda⁴, Takahiro Miyauchi⁵, Hiroyuki Tsutsumi⁶, Hideaki Goto⁷, Takashi Azuma⁸, Tokihiko Matsuda¹, Tadaki Mizumoto¹, Ritsuko S. Matsu'ura¹

¹ADEP, ²Professor Emeritus, Hiroshima Univ., ³Tohoku Univ., ⁴Tokyo Univ., ⁵Chiba Univ., ⁶Kyoto Univ., ⁷Hiroshima Univ., ⁸AFRC, AIST

Hanawa higashi fault zone is a NS-trending, 19-km-long fault and is located along the eastern margin of the Hanawa Basin, northeastern area of the Akita prefecture. Based on detailed interpretation of air photographs, we mapped fault traces in the northern part of the Hanawa Basin where no active faults have been mapped, despite existence of clear topographic boundary between the Hanawa Basin and the Ou Backbone Range. As a result, length of the fault zone changed from 19 km to 26 km. This fact indicates that reevaluation of seismic hazard from the Hanawa higashi fault zone is needed.

This research was funded by grants from the Ministry of Education, Culture, Sports, Science and Technology.

Keywords: active fault, inland earthquake, Hanawa Basin, Hanawa higashi fault zone, air photograph interpretation

SSS032-P09

Room:Convention Hall

Time:May 25 16:30-17:30

Tectonic Geomorphology of the southern part of eastern marginal fault zone of Aizu Basin, Northeast Japan

Tadaki Mizumoto^{1*}, Masayoshi Tajikara¹, Tokihiko Matsuda¹, Takashi Azuma², Hiroyuki Tsutsumi⁶, Toshifumi Imaizumi³, Yasutaka Ikeda⁴, Takahiro Miyauchi⁵, Ritsuko S. Matsu'ura¹

¹Earthquake Reserch Center, ADEP, ²AFERC, AIST, ³Tohoku Univ., ⁴Univ. of Tokyo, ⁵Chiba Univ., ⁶Kyoto Univ.

At the eastern edge of the Aizu basin close to the foot of the Ou backbone range, several active faults constitute a fault zone trending north to south, named of the eastern marginal fault zone of Aizu Basin. These fault traces were found to prolong towards a mountainous area constituting a fault zone extending to about 49 km in length. In the mountainous area, the two faults, Ouchikuramura fault and Simogo fault constituting the southern part of the fault zone, are mapped by Nakata and Imaizumi (2002).

In the vicinity of Otokane, at the southeast part of Shimogo town, the Katodani River flows westward and forms several terraces along them. The fault produces westward facing scarps across the late Pleistocene terraces (probably formed in and around 17,000 years ago) to be displaced in flexure scarp facing the westward and having a vertical deformation of 5 m. At the opposite bank of the river, the lower terrace is also formed with a vertical component of 1.7 m. The progressive amounts of displacement on the late Pleistocene terraces suggest that at least two faulting events have occurred during past 17,000 years. This fault having a length of about 9 km trending north to south in the mountainous area where no active faults have been mapped are newly named to be Otogane fault and also constitutes a southern edge of the fault zone. The length of the fault zone is extended by 4km to southward including the new fault.

This research was funded by grants from the Ministry of Education, Culture, Sports, Science and Technology.

Keywords: Active fault, Aizu Basin, Simogo town, Otogane fault

SSS032-P10

Room:Convention Hall

Time:May 25 16:30-17:30

Quantitative analysis of tectonic landforms along the Nagamachi-Rifu fault segment by using LiDAR-based 2-m-grided DEM

Yasuo Awata^{1*}, Sakae Mukoyama², Hisashi Sasaki²

¹Geological Survey of Japan, AIST, ²Kokusai Kogyo Co., Ltd.

The location and slip-distribution of the Nagamachi-Rifu fault segment is quantitatively analyzed using 2-m-grided DEM that detected by airborne LiDAR. 0.5-2 m interval contour maps and 150 topographic cross sections generated from DEM permit identification and measurement of 13 river terraces and a 5-km-wide flexure and fault zone in and around the urbanized Sendai city.

The narrowly-defined Nagamachi-Rifu Line fault is a 22-34 km-long and 2.5-km-wide active flexure and fault zone, which consists of the right-stepping Miyagino and Nigatake flexures, and the Dainenji-yama back thrust. Each of flexures and fault displace the terrace surfaces and recent alluvial plain cumulatively. The Dainohara terrace of 100 ka or older has received a maximum 70 m vertical by the movement of the Miyagino and Nigatake flexures. The total deformation of the flexures consist of 70 % of uplifting relative to the alluvial plain. 30-50 % of uplifting of the flexures attribute the deformation by the Dainenji-yama back thrust.

The Yagiyama flexure is an active structure longer than 10km, which developed on the upthrown side of the Nagamachi-Rifu Line fault. A 1-1.5 km-wide asymmetric anticline of the terrace surfaces are newly recognized in the downtown of the city. A 2-km-long scarplet on the river terraces continues northward from 3-km-north of the Yagiyama flexure. The active Tsubonuma-Enda fault is located on the SW of the flexure.

The displacements of river terraces, normalized by the number of faulting event, depict the pattern of slip-distribution along the active fault and flexure with high clarity. The normalized displacements along the Nagamachi-Rifu Line fault are constantly large along the 15-km-long central part of the fault and gradually decrease toward the both ends. On the other hand, the displacement along the Yagiyama flexure increase in the southwestern part of the flexure seems continue to the Tsubonuma-Enda fault. When the 4.5 m of uplifting of lowest terrace has experienced 2 faulting events caused by the Nagamachi-Rif Line fault, the recurrence interval of the fault is estimated to be 5-6 ky or longer based on the age and cumulative displacement of the Dainohara terrace. This estimation is consistent with the paleoseismicity in Holocene revealed at the NE part of the fault.

Keywords: Nagamachi-Rifu Line fault, airborne LiDAR, DEM, tectonic landform, active fault, slip distribution

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P11

Room:Convention Hall

Time:May 25 16:30-17:30

Fault geomorphology identified by the interpretation of stereoscopic images produced from digital elevation model

Hideaki Goto^{1*}, Tomohiro Tatemichi¹

¹Hideaki Goto

Detailed digital elevation model (DEM) data distributed from Geospatial Information Authority of Japan (GSI) has been stored steadily since the Basic Act on Promotion of Utilization of Geographical Information was published in 2007. We produced stereoscopic images from all files of 5m-mesh-DEM made by GSI, and interpreted fault topography on the fluvial plains. The small fault scarps are newly identified on the Kyoto basin, Toyama and Niigata plains. It shows that stereoscopic images from detailed DEM are applied materials for active fault research.

Keywords: active fault, digital elevation model, stereoscopic image, Kyoto basin, Isuguri Fault, Kakuda-Yahiko fault

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P12

Room:Convention Hall

Time:May 25 16:30-17:30

Geologic structure in the epicentral area of the 2008 Iwate-Miyagi Nairiku earthquake

Keiichi Ueta^{1*}, Toshinori Sasaki¹, Daiei Inoue¹, Yasuhira Aoyagi¹, Kazuo Mizoguchi¹, Keishi Nunohara², Toshitaka Yokoyama², Osamu Hasegawa³, Koichi Okuzawa⁴

¹CRIEPI, ²Techno Hase, ³DIA Consultants, ⁴CERES

We performed geologic reconnaissance in the epicentral area of the 2008 Iwate-Miyagi Nairiku earthquake to understand the relation between geological structure and the seismogenic faulting. Geologic mapping reveals that the Miocene Maekawa and Orose Formation form a monoclinial flexure-thrust belt and a fold belt trending NNE-SSW in the epicentral area. The Miocene strata thrust over the terrace deposits at outcrops within the fold and thrust belts. The monoclinial flexures and thrusts may have been developed during formation of the major folds propagating from the reverse basement fault, which corresponds to the source fault responsible for the main shock.

Keywords: Iwate-Miyagi Nairiku earthquake, Fold, Monoclinial flexure, Source fault

SSS032-P13

Room:Convention Hall

Time:May 25 16:30-17:30

Re-Examination of the 1762 Horeki Sado Earthquake

Hisashi Ishii^{1*}, Tomiichi Uetake², Tatsuo Usami³, Hideki Nagumo¹, Ryoichi Nakamura¹

¹TEPCO CO.,LTD.,, ²TEPCO CO., LTD.,, ³Tokyo Univ. Prof. Emeritus

Re-Examination of the 1762 Horeki Sado Earthquake

Hisashi Ishii¹, Tomiichi Uetake², Tatsuo Usami³, Hideki Nagumo¹, Ryoichi Nakamura¹

1 TEPCO CO., LTD., 2 TEPCO CO., LTD., 3 Tokyo Univ. Prof. Emeritus

We found new historical documents of the 1762 Horeki Sado earthquake and re-examined seismic intensity distributions. In this study, we reported new documents and estimation of the source location and magnitude.

The earthquake that occurred off Sado Island caused tsunami damages on the northern coast of Sado Island. The epicenter was estimated by Imamura(1947), Usami(1975,1987,1996,2003), Hatori(1990)and Kawauchi(2000).Usami(1996) estimated the magnitude as 7.0 ,whereas Hatori estimated as 7.2 from tsunami damages and the area of seismic intensity 5.

Seismic intensities were large at Sado Island and the coast of Niigata, Yamagata prefecture. Imamura (1947) estimated epicenter E off Sado Island, somewhere near the midway between Sado and Niigata.

Usami (1975, 1987, 1996) estimated epicenter E off Sado Island, 138.7deg E 38.1deg N and suggested 38.35deg N by the change possibility of disaster area (2003). Hatori (1990) drew refraction diagrams of the hypothetical tsunami to see the shoaling effects around Sado island and concluded epicenter was N off Sado Island, the northern peninsula Hajikizaki,138.3deg E 38.4deg N. Kawauchi (2000) inferred the epicenter W off Awashima Island off Sado Island,138.8deg E 38.6deg N. We investigated historical documents of Niigata, Yamagata and Akita pref. and found 5 new documents that were unpublished in the compiled historical earthquake documents. With these new documents and published historical earthquakes, we tried to estimate the epicenter of this earthquake. The epicenter was estimated N off Sado Island.

Keywords: 1762 Horeki Sado earthquake, Historical earthquake

References: Tatsuo Usami, Materials for comprehensive List of Destructive Earthquakes in Japan. [416]-2001
Akitune IMAMURA: Seismic Activity of Both Sides of the Fossa Magna, Proc, Imp. Acad, 22(1947), 314-321.




図1 宝暦十二(1762)年佐渡地震の震央・震度分布
Fig1 The epicenter, Intensity distribution of the 1762 Horeki Sado Earthquake
ゴシック文字は新史料を示す、字匠(2003) *は、精島を北精島と考
えたときの位置を示す。

Keywords: 1762 Horeki Sado Earthquake, Historical Earthquake

SSS032-P14

Room:Convention Hall

Time:May 25 16:30-17:30

Timing of the last faulting event on the Sekidosan fault of the Ouchigata fault zone, Central Japan

Takashi Azuma^{1*}, Nobuhiko Sugito², Hiroyuki Tsutsumi³, Daisuke Hirouchi⁴, Takashi Hosoya⁵, Koji Magara⁵, Taku Ito⁵

¹AFERC, AIST, ²RSVD, Nagoya Univ., ³Department of Geophysics, Kyoto Univ., ⁴Faculty of Education, Shinshu Univ., ⁵Chuo Kaihatsu Corporation

Sekidosan fault is southeast dipping reverse fault with trending NE-SW, and bounding between the Sekido mountains and Ouchigata plain at the neck of the Noto Peninsula, Ishikawa Prefecture. Sugito et al. (2007) pointed out that the last faulting event had occurred between 850-250 cal.yrBP, whereas the Earthquake Research Committee recognized its timing as between 3200 cal.yrBP - 9 century (Earthquake Research Committee, 2005). We conducted trench excavation and drilling survey for the paleoseismological study at four sites, Mijiro, Sakai, Hongo and Shikinami, on Sekidosan fault in order to obtain the new data to identify the last faulting event of this fault. In this abstract we will show the result of survey and tentative interpretation to them.

A trench and 6 boreholes were excavated at Mijiro site. Trench is excavated on the foot of the flexure on the valley bottom. On the trench walls, humus, silt, sand and gravels deposited after 7000 cal.yrBP were observed. There was no fault in these sediments. Drilling core shows deformation of sediments below the trench floor.

At Sakai site, a trench was excavated on a small scarp on a fan, but this scarp was formed not by faulting, but by erosion. Sediments on the surface of the higher side of this scarp shows they were deposited in wetland. It indicates that trench site could be located on the up-thrown side of the fault concealed in the plain.

At Hongo site, 8 boreholes and a pit were excavated on the both sides of small scarp, which is located on the plain-side of the previous trench site. 2 boreholes were drilled with an angle of 45 degree and others are vertical. From the section and age data of sediments, this scarp could be formed by faulting, but more age data are required to identify the timing of it.

3 boreholes and a pit were excavated at Shikinami site. From the geological section of this site, the scarp was identified as formed by erosion. But existence of humic silt with age of around 7000 cal.yrBP and those elevation of 6-7 m above sea-level, indicate this site could be tectonically uplifted. Based on the age data from the pit, the last uplift event seems to be occurred in historical age.

Keywords: active fault, faulting history, paleoseismological trench excavation, drilling survey, Sekidosan fault, Ouchigata fault zone

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P15

Room:Convention Hall

Time:May 25 16:30-17:30

Paleoseismicity on the Kajiya, Sekigahara and Miyashiro faults in the Yanagase-Sekigahara fault zone, central Japan

Toshikazu Yoshioka^{1*}, Kouichi Sago², Hiroshi Yamane²

¹Active Fault and Eq. Res. Ctr, AIST/GSJ, ²Dia Consultants

The Yanagase-Sekigahara fault zone truncating from Japan Sea to western Gifu prefecture, central Japan, is one of the major fault zones in Japan. The Earthquake Research Committee evaluated that the probability of the earthquake occurrence in the future on the southern part of this fault zone is unknown because of the lack of paleoseismological data. We carried out paleoseismological studies on the Kajiya, Sekigahara and Miyashiro faults in this fault zone to evaluate the rupture probability in the future of these faults, using the fund of the Ministry of Education, Culture, Sports, Science and Technology. A high-angle fault exposed on the trench walls at the Kajiya A site on the Kajiya fault. Radiocarbon dates indicate that at least one faulting event occurred in these 3,300 years. At the Kajiya B site, a fault cutting the bedrock and terrace deposit was observed, and radiocarbon dates indicate that the last faulting event occurred in these 1,000 years. At the Akiba site on the Sekigahara fault, the bedrock covered by the slope deposit and no fault was observed. On the Miyashiro fault, boring surveys show that the top of the Tokai Group and the bottom of young gravel layer are vertically displaced about 35 meters and 3 meters respectively.

Keywords: Yanagase-Sekigahara fault zone, Kajiya fault, Sekigahara fault, Miyashiro fault, trench excavation, paleoseismology

SSS032-P16

Room:Convention Hall

Time:May 25 16:30-17:30

Stress inversion method from fault slip senses and its application to active fault data

Katsushi Sato^{1*}, Hiroyuki Tsutsumi¹, Atsushi Yamaji¹

¹Div. Earth Planet. Sci., Kyoto Univ.

Stress inversion methods from fault-slip data usually require to observe fault surface orientation, slip orientation and sense of shear. However, the descriptions of fault geometry made by many researchers are sometimes incomplete for lack of slip orientations. In such cases, Andersonian fault model may be applied in spite of the existence of oblique slip faults. This presentation introduces a stress inversion method for incomplete fault-slip data, which is applied to the active fault data compiled by Tsutsumi et al. (this meeting) in the Kinki and Chubu districts, central Japan.

The stress inversion method used in this study was developed by Sato (2006). A fault-slip datum has constraint on stress state through the assumption that a fault slips in the direction of resolved shear stress (Wallace-Bott hypothesis). The constraint can be expressed as a region in the deviatoric stress space which is geometrically the surface of five-dimensional unit sphere. If a datum is incomplete, the area of constraint should be large. The inversion method superposes the constraint regions from all observed faults to compose a fitness distribution on the sphere. Since the deviatoric stress space is a metric space with the measure of difference between stress tensors (Yamaji and Sato, 2006), the area of constraint region can be used as the weights in the superposition process. Finally, the stresses which give the maxima of fitness are picked up as optimal solutions.

The active fault data set from about 200 locations were analyzed, and a reverse faulting stress with WNW-ESE compression was obtained. Although the data included no information about slip orientations, stress could be constrained in a small area in the deviatoric stress space (within 20 degrees around the optimal solution) due to the variation of fault surface orientations. Most of fault data are concordant with the optimal stress, while a small number of outliers deviate by only several degrees in the stress space. The fact that single stress can explain almost all of active fault slip senses shows the uniformity of stress state in the district.

References

Sato, K., 2006. *Tectonophysics* 421, 319-330.

Yamaji, A. and Sato, K., 2006. *Geophys. Jour. Int.* 167, 933-942.

Keywords: stress tensor inversion, fault-slip data, active fault, Hough transform

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P17

Room:Convention Hall

Time:May 25 16:30-17:30

Regional stress field across Kinki and Chubu regions derived from stress inversion analysis of active fault data

Hiroyuki Tsutsumi^{1*}, Katsushi Sato¹, Atsushi Yamaji¹

¹Kyoto University

We revealed regional stress field across Kinki and Chubu regions based on stress inversion analysis of active fault data. We compiled fault slip data including fault plane orientation and sense of slip (right-lateral, left-lateral, reverse, normal and combination of strike-slip and vertical-slip). A stress field composed of WNW-ESE-oriented σ_1 with almost vertical σ_3 was detected by the analysis. This suggests that Kinki and Chubu regions have been under a fairly uniform stress field in the late Quaternary.

Keywords: active faults, fault slip data, regional stress field, stress inversion analysis

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P18

Room:Convention Hall

Time:May 25 16:30-17:30

Seismic reflection profiling across the Shufuji fault, Kinki district

Shigeru Toda^{1*}, Mai Morioka¹, Shigehiro Katoh², Atsumasa Okada³

¹Aichi University of Education, ²Division of Natural History, Hyogo Museum, ³Ritsumeikan University Global Innovation

We present high-resolution seismic reflection profiling acquired by the hammer and 24-channel recording system. A 1.0 meter source and geophone spacing give a 0.5 meter CMP spacing on the profile. The normal CMP stacking fold had 24 traces used by the same shot point. The location and continuity of active faults comprising a part of the Kinki region are clearly expressed in terms of topography. This area is the so-called Kinki Triangle. This study presents the results of seismic reflection surveys across the Fukurojou Maiseki Valley in the northern part of the Kinki Triangle. The subsurface configurations of the active faults are correlated with geomorphological fault trace and are related with the fault strikes.

Keywords: Seismic Reflection Survey, Shufukuuji Fault

SSS032-P19

Room:Convention Hall

Time:May 25 16:30-17:30

Tectonic topography and active fault outcrop in the eastern focal area of the 1943 Tottori Earthquake

Toshinori Sasaki^{1*}, Keiichi Ueta¹, Yasuhira Aoyagi¹, Daiei Inoue¹, Takahiro Nagata², Kota Shimogama², Osamu Hasegawa², Makoto Yanagida³, Norihisa Goto³

¹CRIEPI, ²Dia consultants Co. Ltd., ³Hanshinconsultants Co. Ltd

The magnitude (Mj) 7.2 Tottori earthquake occurred on 1943 in the Tottori Province of Japan, resulting in great damage throughout Tottori City urban area mainly. In the hill or mountain area west Tottori plane, there were two echelon surface ruptures called Yoshioka fault and Shikano fault associated with this earthquake known by previous studies (Tsuya, 1944). On the other hands, in the hill or mountain area east Tottori plane, surface rupture had not been confirmed.

Recently, the active fault was found by aerial photo investigation in this area (Takada et al., 2003), and there was not surface rupture because of the vertical slip distribution of this area was estimated to be deeper than the west area (Nakata, 2009). We have been studied geomorphological and geological survey for the purpose of seismic hazard assessment. In this presentation, we conclude the results of high resolution topographic investigation and active fault outcrop observation along the tectonic topography in the eastern part of the focal area.

Aerial photographic survey is using 1/20,000 scale (CG-75-2X) and 1/10,000 scale by U.S. military forces. We interpret DEM using shaded relief map and stereoscopic bird's-eye view made from 2m mesh topographic data which is obtained by airborne laser scanner of Kokusai Kogyo Co., Ltd. The outcrop survey and survey of the small trench excavated by man power are conducted by observation and sketch of the wall surface after the wall has been made flat and smooth.

As a result of topographic survey, we found consecutive tectonic topography which is right lateral displacement of several ridge and valley lines along the ENE-WSW lineament in the mountain area from Momodani, Tottori city to Kujira, Fukube town. We interpret four other relatively short lineaments which are parallel in this lineament east of Tottori City urban area. These short lineaments are composed of cols and rectilinear valleys, and we found an active fault outcrop (Takiyama outcrop) along the southernmost part of the short lineament. Takiyama outcrop have the active fault which cut the layer of DKP (Daisen-Kurayoshi tephra: 55ka) which apparent vertical displacement is ca.90cm, and we confirm that the fault has been repeatedly moved during late Quaternary. A small trench is excavated to study the latest event in focus just behind the outcrop in the col topography. As a result of this survey, the age of the latest event is limited between before depositional age of K-Ah (Kikai-Akahoya tephra: 7.3ka) after AT (Aira-Tanzawa tephra: 26-29ka).

Keywords: 1943 Tottori earthquake, tectonic geomorphology, active fault outcrop, DEM investigation, aerial photo investigation

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P20

Room:Convention Hall

Time:May 25 16:30-17:30

Study on Late Pleistocene to Holocene activity of the eastern part of Shinji Fault

Shuji Hirokane^{1*}, Kohei Kurooka¹, Yuji Ito¹, Junji Fujiki², Shuji Hata², Takenobu Tanaka³

¹The Chugoku Electric Power Co.,Inc, ²Chuden Engineering Consultants, ³HANSHIN CONSULTANTS Co., Ltd.

We investigated the eastern part of the Shinji Fault to clarify the activity during Late Pleistocene to Holocene by arrayed boring and trenching surveys at the Shimoubeo in Matsue city.

A fault is confirmed by one of arrayed borings crossing the extended line of the lineament based on our geomorphological study. It is presumed that the fault deformed the layer considered to be MIS7 or older, which is weathered and distributed below the layer that contains grains originated from the Daisen Matsue Pumice (DMP), and not deformed upper layer.

Another fault confirmed on the trench wall crossing the active fault shown on Nakata et al.(2008) deforms "layer A" considered as MIS6 or older that based on tephra and pollen fossil analysis shows some time gap with "layer B" contains grains originated DMP.

The activity after the fall of DMP is not confirmed on both faults at the site.

Keywords: Shinji Fault, trenching survey

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P21

Room:Convention Hall

Time:May 25 16:30-17:30

Study on Late Pleistocene to Holocene activity of the western part of Shinji Fault

Yuji Ito^{1*}, Shuji Hirokane¹, Kohei Kurooka¹, Msahiro Nagase², Junji Fujiki², Takenobu Tanaka³

¹The Chugoku Electric Power Co.,Inc., ²Chuden Engineering Consultants, ³HANSHIN CONSULTANTS Co., Ltd.

We investigated the western part of the Shinji Fault to clarify the activity during Late Pleistocene to Holocene by arrayed boring survey at the Sadahongo-Sakoya in Matsue city.

The result of survey shows systematic vertical displacement on the Daisen Matsue Pumice (DMP) layer and the upper layers.

The vertical displacement of key beds is estimated to be 1.3m at about 25,000¹⁴C yBP, 0.8m at about 10,000¹⁴C yBP and 0m at 7,000 to 10,000¹⁴C yBP.

The latest faulting event is presumed to be between 10,000¹⁴C yBP and 25,000¹⁴C yBP.

The degree of activity on the western part of Shinji fault is lower than previous studied sites located at the Minamikoubu, Kashima town along the central part of the Shinji fault.

Keywords: Shinji fault, displacement of fault, arrayed boring

SSS032-P22

Room:Convention Hall

Time:May 25 16:30-17:30

Seaward extension of the Nishiyama Fault Zone off Fukuoka, western Japan

Koji Ito^{1*}, Shintaro Abe², Ryoyu Arai³, Yukinobu Okamura⁴, Azusa Nishizawa¹

¹JCG, ²ADEP, ³KGE, ⁴AFERC

The Nishiyama Fault Zone is an active left-lateral strike-slip fault existing between Fukuoka City and Kitakyusyu City and is estimated to make a M7.3(Richter scale) earthquake (the Headquarters for Earthquake Research Promotion, 2004). This fault zone has been rated an overland fault. The Off O-shima Fault was discovered in the northwest extension sea area of the Nishiyama Fault Zone, but it had not seemed a sequence of the Nishiyama Fault Zone because they were away by 5 kilometers. On the other hand, the Kego Fault is 20 kilometers southeast of the Nishiyama Fault Zone and also had seemed to be an overland fault. But the 2005 West Off Fukuoka Prefecture Earthquake (M7.0 in Richter scale) happened in the northwestern extension sea area of the Kego Fault and the fault seemed to extend to the northern sea area. In response to this, Abe et al.(2010) conducted reflection seismic survey and vibratory coring and revealed that the Nishiyama Fault Zone extended to the northwest sea area, too. Then, Japan Coast Guard conducted accurate topographic survey in the same area and discovered tectonic reliefs on the seafloor. In this study, the fault distribution is investigated by these survey analyses. Survey vessel and multibeam echo sounder used by Japan Coast Guard are "Kaiyo" and EM302, respectively.

The depth of this sea area is between 60 and 100 meters and northern area is deeper. Sand waves with long frequency and gentle slope are found in southern area and sharp and small-scale reliefs are distributed in northern area. This geomorphological difference is caused by the agglomeration degree of sediments. The new and soft sediments with sand waves cover the old and hard bed.

Lineament composed of channels and bulges in direction of northwest-southeast was found in the northwestern extension sea area of the Nishiyama Fault Zone and the Off O-shima Fault. The bulges are formed by the old and hard bed uplifted by the faults of both sides and crop out in the new and soft sediments. The maximum relative elevation is 5 meters. The channels deform both new sediments and old bed. The maximum depth is 2 meters. The flower structure is found in the seismic profile under the channels and this shows that the channels were formed by a lateral strike-slip fault. The main lineament runs for about 30 kilometers from O-shima to Oki-no-shima and extends further north. Some short faults are shown in the seismic profiles and on the seafloor and indicate that the fault is splitting toward north.

The surveyed sea area was a land in the Last Glacial Maximum about 20,000 years ago. The soft sediments with sand waves laid down after the submergence.

Keywords: Nishiyama Fault Zone, active fault, lateral fault, sea area, seafloor topography, seismic reflection survey

SSS032-P23

Room:Convention Hall

Time:May 25 16:30-17:30

Extension position and continuity of the Nishiyama fault in Chikuzen-Oshima Island, Fukuoka Prefecture

Tatsuya Sugiyama^{1*}, Shoichi Shimoyama², Hiroyuki Tsutsumi¹, Daisuke Ishimura¹

¹Dept. Geophysics, Kyoto Univ., ²Earth and Planetary Sci., Kyushu Univ.

The Nishiyama fault, a left-lateral strike-slip fault, locates from Fukutsu City to Iizuka City (Pref. Fukuoka, 1996). In the nearby Genkainada Sea, the 10-km long Off Oshima fault, was identified in Genkainada Sea, where is the offshore northwestern extension of the Nishiyama fault (e.g. Iwabuchi, 1996). Tsutsumi et al. (2008) suggested, on the basis of geomorphology, that the Off Oshima fault belonged to the Nishiyama fault zone. In August 2010 the Japan Coast Guard re-surveyed the Off Oshima fault and identified a 30-km long northwesterly extension.

However, there is a gap of longer than 10 km between the northern end of the Nishiyama fault and the southern end of the Off Oshima fault. Chikuzen-Oshima Island locates between these faults. The aims of this study are to identify the position of an active fault in Chikuzen-Oshima Island and the continuity of the Nishiyama fault and the Off Oshima fault.

Aerial photograph interpretation was carried out to identify tectonic geomorphology such as fault scarps or the displacement of valleys or mountain ridges. Geological and topographical field surveys were conducted.

The results were as follows. We revealed topographical features of recent fault activity in the central part of the island. We found two outcrops of a fault, low fault scarps and sinistral displacements of the valley and a mountain ridge. It was therefore concluded that this was an active fault. Tephra analysis of deposits of fluvial terrace, which is displaced 1.5 m vertically by the latest faulting, in the Tani district was carried out. It is clarified that these deposits contained Kikai-Akahoya tephra (K-Ah: 7.3ka yBP) and Aira-Tn tephra (AT: 26-29ka) tephras [Machida and Arai(2003)]. We concluded that this terrace age is younger than 7.3ka.

In conclusion, an active fault was identified in the central part of Chikuzen-Oshima Island and the Holocene terrace surface was displaced 1.5 m vertically by the latest faulting.

Keywords: active fault, Chikuzen-Oshima Island, Nishiyama fault, Off Oshima fault

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P24

Room:Convention Hall


Time:May 25 16:30-17:30

Fault exposure along the west foot of the Kodaiji mountain, Munakata-City, Fukuoka Prefecture.

Yosuke Nakamura^{1*}, Kiyohide Mizuno¹

¹GSJ, AIST

In the Fukuoka Coastal zone, a new outcrop of reverse active fault was observed along the on the west foot of the Kodaiji mountain, Munakata-City, Fukuoka Prefecture. The strike and dip of the fault surface are N20W and 28W, respectively. In this outcrop, terrace gravels and Tertiary rocks were deformed and displacement of the bottom terrace gravels on this thrust fault is about 3.2m in the vertical component. Also, we reveal the nature of the active fault located the Fukuoka Coastal zone on the basis of fault outcrops and geomorpic features.



Keywords: active fault, fault exposure, Fukuoka Coastal zone, C class fault

SSS032-P25

Room:Convention Hall

Time:May 25 16:30-17:30

Tsunami deposit investigation in the Fukushima coastal area

Kenji Oikawa¹, Makoto Takao¹, Mitsunobu Usami^{2*}, Riichiro Miyawaki³

¹TOKYO ELECTRIC POWER COMPANY, ²DIA CONSULTANTS co.,ltd, ³HANSHIN CONSULTANTS co.,ltd

Jogan tsunami deposit which was caused by AD869 Jogan earthquake, is distributed in the Fukushima coastal area (from Sendai plain to Soma city: e.g., Sawai et al., 2007; Sugawara et al., 2002), and this information has been used to construct and verify the fault models (Satake et al., 2008). The detailed distribution of the Jogan Tsunami deposit, however, has been not clear. Therefore, in order to obtain the more detailed distribution of the Jogan tsunami deposit, we investigated in the five investigation points (south Matsukawaura, Urajiri, Hotokehama, Shimo-Asamigawa and Takaku) of the Fukushima coastal area. Here we report investigation results of the hand breaker boring core sample by geological and geochronological methods.

In south Matsukawaura, we found two tsunami deposits. The upper tsunami deposit is possibly thought to be ca. AD1500 tsunami deposit (Sawai and Shishikura, 2010), and the lower one is the Jogan tsunami deposit determined by the carbon dating. The run up height by the Jogan tsunami is possibly 0.5m above sea level.

In Urajiri, we found four tsunami deposits. These tsunami deposits were estimated to be AD1500 tsunami deposit, AD970 to AD1160 tsunami deposit, the Jogan tsunami deposit and BC730 to AD420 deposit, respectively. The run up height of the Jogan tsunami deposit is estimated less or equal 4m above sea level.

In Hotokehama, Shimo-Asamigawa and Takaku, we did not find the tsunami deposits.

Paleo-sea depth changes and tsunami deposits due to the Kanto earthquakes in Ena Bay, south coast of Miura Peninsula

Takashi Chiba^{1*}, Takeo Ishibe², Kenji Satake², Kunihiko Shimazaki³, Toshihiko Sugai¹, Akihito Nishiyama², Tomoya Harada², Kentaro Imai⁵, Yuichi Namegaya⁴, Toshihiro Ueno²

¹Grad.Sch.of Frontier Sci.,The Univ.Tokyo, ²Earthquake Res. Inst., Univ. of Tokyo, ³Assoc. for Earthquake Disaster Prev., ⁴AFERC, ⁵Tohoku Univ. DCRC

Paleo-earthquake studies by geological and geomorphological surveys are important for earthquake forecast because they provide information not only of earthquake occurrence time and magnitude but also on environmental changes during co-seismic and inter-seismic intervals.

The recurrence interval of the great interplate earthquake along the Sagami Trough, Kanto earthquake, is estimated to be 200-400 years (Earthquake Research Committee, 2004). However, earthquake histories prior to the 1703 Genroku Kanto earthquake have not been revealed from historical literature. Miura Peninsula, which locates in northeastern part of the Sagami Trough, has been uplifted (Kanie et al., 1989) and tsunamis attacked along the coast of the Kanto region accompanied by the previous Kanto earthquakes (Hatori et al., 1973). Shimazaki et al. (2009) conducted Geo-slicer surveys in Koajiro Bay, Miura Peninsula and investigated histories of the previous Kanto earthquakes. As a result, they suggested that the 1293 earthquake causing destructive damage in and around Kamakura was the Kanto earthquake prior to the 1703 Genroku earthquake, as pointed out by Ishibashi (1991).

The purpose of this study is to reveal histories and identify tsunami deposits of previous Kanto earthquakes, and processes of uplift and subsidence from a reconstruction of paleo-sea depth using diatom and grain size analyses in Ena Bay, Miura Peninsula. In May and November, 2009, we conducted 3m length handy Geo-slicer surveys at Ena Bay, south coast of Miura peninsula. We have basically analyzed 3 cores (ENA-C, ENA-E and ENA-F).

As a result, three (ENA-E) or four (ENA-F) coarse layers including shell fragment and gravel are recognized. These event deposits erode a subsurface layer indicating that they accompanied with a strong current. Diatom analysis indicates an increase or a decrease of relative abundance of marine species, suggesting a change of sea depth. Namely, marine benthic species gradually decrease prior to the deposition of tsunami deposits indicating coastal uplift or sea level fall, and benthic species increase above tsunami deposits indicating coastal subsidence or sea level rise. It is revealed that Miura Peninsula uplifted about 1.5 m at the time of the 1923 Kanto earthquake and now subsides with a rate of about 3.7 mm/year from tide gauge record at Aburatsubo (Geospatial Information Authority of Japan, 2010). The characteristics of diatom analysis suggest that environmental changes corresponding to these co-seismic and inter-seismic crustal movements. This is consistent with the results in Koajiro Bay (Shimazaki et al., 2008). We concluded that these event layers are tsunami deposits accompanied with the previous Kanto earthquakes, named as T1, T2, T3, T4 and T5 unit from the top to the bottom, respectively.

The T1 unit is concluded as a tsunami deposit accompanied with the 1923 Taisho Kanto earthquake using Pb-210 dating. However, radiocarbon ages indicate that the T2, T3, T4 and T5 unit deposited about 2000 cal. yBP, 3000 cal. yBP, 3300 cal. yBP and 3700 cal. yBP, respectively. This is consistent with histories of previous Kanto earthquakes inferred from marine terraces in Boso Peninsula (Shishikura, 2003). Moreover, at least three coarse-grained layers are recognized between T1 and T2 at ENA-C. These layers are possibly identified as tsunami deposits from similar tendencies of grain size distribution and diatom species.

Keywords: Kanto earthquake, Tsunami deposit, Ena bay, Paleo-sea depth change, Diatom analysis

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SSS032-P27

Room:Convention Hall

Time:May 25 16:30-17:30

Classifications of the earthquake type and the recurrence interval for the Kanto earthquake

Haeng Yoong Kim^{1*}

¹ERI, University of Tokyo

Classification of past Kanto earthquake is examined as follows.

- A. 1923-type; poorly known
- B. 1703-type; 950 to 2,500 years [Seno (1977)], 2,000 to 2,700 years [Shishikura (2003)].
- C. 1923-1703 combination-type; 800 to 1500 years [Matsuda et al. (1974, 1978)], 1,450 to 2,600 years [Nakata et al. (1980)].
- D. 1923-1703 addition-type; 260 to 320 years [Kanamori (1973)], 200 to 300 years [Ishibashi (1977)], 180 to 400 years [Seno (1977)], 470 to 1,143 years [Matsuda (1985)], 300 years [Kumaki (1982)], two patterns of ~600 years and ~900 years [Kumaki (1988)], 380 to 400 years [Shishikura (2003)].

Keywords: Kanto earthquake, earthquake type, recurrence interval, occurrence time, earthquake cycle, long term prediction

SSS032-P28

Room:Convention Hall

Time:May 25 16:30-17:30

Re-evaluation of the Thoen fault activity in the Lampang basin, northern Thailand

Weerachat Wiwegwin^{1*}, Yuichi Sugiyama², Ken-ichiro Hisada¹, Punya Charusiri³, Suwith Kosuwan⁴, Preecha Saithong⁴

¹University of Tsukuba,Japan, ²AIST, Japan, ³Chulalongkorn University, Thailand, ⁴Department of Mineral Resources,Thailand

We applied remote-sensing technique and geomorphic index analysis to a study of the NE-SW-striking Thoen fault, Lampang basin, northern Thailand. Significant morphotectonic landforms caused by normal fault in the Lampang basin are represented by fault scarps, triangular facets, wine-glass canyons and linear mountain front. Along the Thoen fault, the SL index indicates the steeper slope near the mountain front. These SL indices possibly relate to a normal fault system. Moreover, most of Vf and Smf values at Ban Mai and Sop Prap segments are low (0.44 to 2.75 of Vf and 1.11 to 1.82 of Smf). These geomorphic indices may indicate slightly active tectonic area that results from vertical slip on the normal fault. Geomorphological features and geomorphic indices of the study area envisage active normal faulting. However, stratigraphic units of trench at Ban Don Fai indicate no clear-cut evidence of any recent fault movement. At Ban Don Fai trench No. 2, AMS radiocarbon and OSL ages suggested that the sediments of the lowest unit were deposited between 960 to 910 years ago. Therefore, the last movement of the Ban Don Fai segment might have occurred earlier than 960 years ago. High resolution seismic survey data for coal investigation in the Mae Tha sub basin (adjacent to the present trenching site) reveal that the Thoen fault is a concealed fault. Thus, the Ban Don Fai segment of the Thoen Fault might be a concealed fault.

In the northern Thailand, there are six major faults that have been regarded as an active fault: Mae Chan, Mae Tha, Pua, Thoen, Mae Hong Son and Mae Ping faults. This study proved that the Thoen fault might be not so active at least during the latest several thousand years. More researches on the other active faults are necessary to evaluate the fault activity and very recent tectonics in northern Thailand.

Keywords: Thoen fault, Normal fault, Morphotectonic landforms, Geomorphic index analysis, OSL and AMS radiocarbon datings, Lampang basin, Northern Thailand