Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



AHW26-P10

会場:コンベンションホール

中部山岳地域における降水の安定同位体比の時間変動特性 Temporal variation of stable isotope ratio in precipitation on Chubu-mountainous areas: case study of Mt. Ontake in cent

浅井 和由^{1*}, Fantong Wilson Yetoh², 辻村 真貴³ ASAI, Kazuyoshi^{1*}, Fantong Wilson Yetoh², TSUJIMURA, Maki³

¹ 株式会社地球科学研究所, ²Institute of Mining and Geological Research, Cameroon, ³ 筑波大学生命環境系 ¹Geo-science Laboratory Inc, ²Institute of Mining and Geological Research, Cameroon, ³Graduate School of Life and Environmental Sciences, University of Tsukuba

In order to clarify the temporal variation of the stable isotope ratio in precipitation on the mountainous area in Japan, monthly precipitation was collected at 11 sites on Mt.Ontake from Jan 2003 to Dec 2005, and determined their delta-values. d-excess values ranged from 5 to 34 per mil, and showed clear seasonal variations in which values are low in rainy season from spring to fall and high in snow season (winter), reflecting the change of monsoon. The isotope ratios showed high delta-values in spring, low delta-values in winter and intermediate delta-values in summer to fall. These seasonal variations reflect neither the influence of the amount effect nor the temperature effect, indicating that the values variation in precipitation in Mt. Ontake do not have a close relation with a local weather condition. Rain-bearing weather condition and spatial variation of rainfall amount around Mt. Ontake were investigated, and discussed a factor controlling the seasonal variation of delta-values. In the warm rainy season, the low delta-value in precipitation were observed in the period at which the rainfall amount of windward (south-east) area of Mt. Ontake was high, suggesting that delta-values variation on Mt. Ontake during the warm rainy season depend on the rainfall amount during the vapor mass transpiration from Pacific coast to Mt. Ontake. Moreover, it is indicated that the low delta-values of snow cover in winter was caused by the inland effect, considering the existence of heavy snowfall area in windward. These variations of delta-value associated with the rainfall amount on surrounding area is considered to be reflect the geographical location of Chubu-mountainous area which is away from the ocean generating a vapor mass.

Keywords: Chubu-mountains areas, Mt. Ontake, precipitation, stable isotope, vapor mass transpiration,, inland effect