

## 八ヶ岳南麓における湧水中の硝酸イオンの起源と水文条件が濃度分布に与える影響について

### Analysis of nitrate variation by source and hydrologic factors in spring water of Mt. Yatsugatake

中村 高志<sup>1\*</sup>, 浅井和由<sup>2</sup>, 安原正也<sup>3</sup>, 藪崎志穂<sup>4</sup>, 鈴木裕一<sup>4</sup>

NAKAMURA, Takashi<sup>1\*</sup>, Kazuyoshi Asai<sup>2</sup>, Masaya Yasuhara<sup>3</sup>, Shiho Yabusaki<sup>4</sup>, Yuichi Suzuki<sup>4</sup>

<sup>1</sup> 山梨大学・国際流域環境研究センター, <sup>2</sup> 株式会社地球科学研究所, <sup>3</sup> 産総研・地質調査総合センター, <sup>4</sup> 立正大学・地球環境科学部

<sup>1</sup>ICRE, University of Yamanashi, <sup>2</sup>Geo Science Laboratory Inc., <sup>3</sup>Geological survey of Japan, AIST, <sup>4</sup>Rissho University

Concerns regarding ground-water contamination in the Mt. Yatsugatake on the west slope have arisen due to a steady increase in averaged nitrate concentrations (from 1.2 to 3.5 NO<sub>3</sub>-mg/l) during t

he past 15 years (from 1987 to 2002) in spring waters (Shimizu et al. 2007). Mt. Yatsugatake is volcano in central Japan having elevations of 2899m. The artificial and local resort towns are locating blow 1100m elevation. The stock farms are expanding from upper 1100m until 1400m elevation.

Twenty-nine water samples were collected from 750m to 2380m elevation. Multiple isotopic tracers were used to identify the sources and extent of nitrate dynamics in groundwater under the recharging systems.

The ranges of nitrate-N concentration were from <0.05 to 17.8mg/l. Few samples of high nitrate-N concentrations (5.8 and 17.8 mg/l) were located in locally concentrated of agricultural and local resort town. And 80% of collected samples were <0.9 mg/l. Although delta 15N-NO<sub>3</sub> values (from -0.4 to 10.5permil) were highly variable both spatially and vertically. The high delta 15N-NO<sub>3</sub> values are decreasing with the averaged recharge elevation of springs which was estimated by elevation effects of water hydrogen isotope values on Mt. Yatsugatake reason. Relatively high isotope value samples (4.4-10.1permil) were detected blow 2000m elevation indicated that nitrate in springs mixed from human/animal wastes consistent with landuses. The relationship between NO<sub>3</sub>-N concentration and delta 15N values was insignificant. However the high NO<sub>3</sub>-N concentration (>0.9 mg/l) springs having relatively small amount of discharges (850 m<sup>3</sup>/day) indicated the effects for the distribution of NO<sub>3</sub>-N concentration is nitrogen sources rather than quantity of the groundwater.

キーワード: 湧水, 八ヶ岳, 硝酸イオン, 硝酸イオンの窒素・酸素安定同位体比, 水の水素・酸素安定同位体比

Keywords: Spring water, Mt. Yatsugatake, Nitrate, Nitrate nitrogen and oxygen isotopes, Water hydrogen and oxygen