

火星におけるCO₂, CH₄ハイドレートの放射線分解による有機物生成の可能性 Possibility of organic matter formations in irradiated CO₂ and CH₄ hydrates on Mars

大島 基^{1*}, 谷 篤史¹, 北野勝久², 菅原武³, 大垣一成³

OSHIMA, Motoi^{1*}, TANI, Atsushi¹, Katsuhisa Kitano², Takeshi Sugahara³, Kazunari Ohgaki³

¹ 大阪大学大学院理学研究科, ² 大阪大学大学院工学研究科, ³ 大阪大学大学院基礎工学研究科

¹Science, Osaka Univ., ²Engineering, Osaka Univ., ³Engineering Science, Osaka Univ.

To look for evidence of life on Mars, explorations of organic matter is important. There are some possibilities of the formations of organic matter on Mars; for example, by biological activity and by radiolysis and photolysis of mixtures of icy materials. If organic matter exist on Mars, to identify the detected organics as the matter by biological activity, it is necessary to investigate reaction products and their amount by radiolysis and photolysis of the ice mixtures, and to distinguish organics by biological activity from ones by radiolysis and photolysis. As the icy materials on Mars, it is suggested that gas hydrates would exist on Mars, based on the discussion of temperature and pressure conditions. Gas hydrates are crystalline inclusion-compounds, which are composed of hydrogen bonded water molecules encaging gas molecules (e.g. CO₂, CH₄). Most of them are stable at high pressures and/or low temperatures. Average temperature of Martian surface is about 210 K, the Martian atmosphere consists of about 95% CO₂ and the average atmospheric pressure is about 0.56 kPa (Kieffer et al., 1993). From the observations of the Mars Express spacecraft, methane with 10-40 ppmv concentrations has been detected in the Martian atmosphere and on surface of the ground (Formisano et al., 2004; Mumma et al., 2009). In Martian conditions, it is suggested that CO₂ and CH₄ hydrates are formed in about 10 m below the ground and on the polar caps in winter (e.g. Max and Clifford 2001; Prieto-Ballesteros et al., 2006). Since water ice exists on Martian surface, the environment on the Martian surface should be enough to form CO₂ and CH₄ hydrates. In this case, Martian gas hydrates should be irradiated by natural radiation from radioisotopes in sediments as well as cosmic rays, which may cause radical formation in CO₂ and CH₄ hydrates.

Radiation-induced radicals in CH₄ and CO₂ hydrates have been investigated by electron spin resonance (ESR) measurements. Methyl radical is mainly formed by gamma-ray irradiation at 77 K in CH₄ hydrate (Takeya et al., 2004), and not stable above 180 K at 0.1 MPa (Tani et al., 2006). Carboxyl radical, hydrogen atom and hydroxyl radical are formed by gamma-ray irradiation at 77 K in CO₂ hydrate, hydrogen atom and the hydroxyl radical are not stable above 120 K at 0.1 MPa and the carboxyl radical quickly disappears at 180 K (Oshima et al., submitted). In thermal conditions on Mars, these radicals in the hydrates are unstable, and some products may form through the radical reactions. In this study, to investigate the organic products from radicals induced in CO₂ and CH₄ hydrates on Martian conditions, we analyzed the aqueous solution after dissociation of the hydrates irradiated at 195 K by ion chromatography and gas chromatography-mass spectrometry (GC-MS).

Formic acid and oxalic acid are observed in aqueous solution after dissociation of irradiated CO₂ hydrate. Methanol and formaldehyde are observed in irradiated CH₄ hydrate. If CO₂ and CH₄ hydrates exist on Mars, these organics will be formed in the hydrates and accumulated on the polar caps and below subsurface.

キーワード: ガスハイドレート, 有機物, ラジカル, イオンクロマトグラフ, ガスクロマトグラフ質量分析, 火星

Keywords: Gas hydrates, Organic matter, Radicals, Ion chromatography, Gas chromatography-mass spectrometry (GC-MS), Mars