## **Japan Geoscience Union Meeting 2012**

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



BAO01-03

会場:202

時間:5月24日09:30-09:45

### たんぽぽ:たんぽぽ計画:有機物・微生物の宇宙曝露と 宇宙塵・微生物の捕集 Tanpopo: Astrobiology exposure and micrometeoroid capture experiments

山岸 明彦  $^1$ , 横堀 伸一  $^{1*}$ , 橋本 博文  $^2$ , 今井 栄一  $^3$ , 河合 秀幸  $^4$ , 小林 憲正  $^9$ , 三田 肇  $^5$ , 中川 和道  $^6$ , 奥平 恭子  $^7$ , 田端 誠  $^2$ , 太刀川 純孝  $^2$ , 薮田 ひかる  $^8$ , 山下 雅道  $^2$ , 矢野 創  $^2$ , たんぽぽ WG  $^2$ 

YAMAGISHI, Akihiko<sup>1</sup>, YOKOBORI, Shin-ichi<sup>1\*</sup>, HASHIMOTO, Hirofumi<sup>2</sup>, IMAI, Eiichi<sup>3</sup>, Hideyuki Kawai<sup>4</sup>, KOBAYASHI, Kensei<sup>9</sup>, MITA, Hajime<sup>5</sup>, NAKAGAWA, Kazumichi<sup>6</sup>, OKUDAIRA, Kyoko<sup>7</sup>, Makoto Tabata<sup>2</sup>, Sumitaka Tachikawa<sup>2</sup>, YABUTA, Hikaru<sup>8</sup>, Masamichi Yamashita<sup>2</sup>, YANO, Hajime<sup>2</sup>, Tanpopo WG<sup>2</sup>

 $^1$ 東京薬大・生命科学,  $^2$ JAXA/ISAS,  $^3$  長岡技大・生物,  $^4$  千葉大・理,  $^5$  福岡工大・工,  $^6$  神戸大・院人間発達環境学,  $^7$  会津大,  $^8$  大阪大・院理,  $^9$  横浜国大・院工

<sup>1</sup>Sch. Life Sci., Tokyo Univ. Pharm. Life Sci., <sup>2</sup>JAXA/ISAS, <sup>3</sup>Dept. Bioeng., Nagaoka Univ. Tech., <sup>4</sup>Facl. Sci., Chiba Univ., <sup>5</sup>Facl. Eng., Fukuoka Inst. Tech., <sup>6</sup>Grad. Sch. Human Develop. Environ., Kobe Univ., <sup>7</sup>Univ. Aizu, <sup>8</sup>Grad. Sch. Sci., Osaka Univ., <sup>9</sup>Grad. Sch. Eng., Yokohama Natl. Univ.

For the origin of life on Earth emerged within a short period after the end of heavy bombardment, Panspermia hypothesis was proposed (e.g. Arrhenius 1908; Crick 1981). Recent findings of the Martian meteorite suggested possible existence of extraterrestrial life, and interplanetary migration of life as well.

Microbes have been collected from high altitude using balloons, aircraft and meteorological rockets since 1936, even it is not clear how could those microbes be ejected up to such high altitude. Spore forming fungi, spore forming Bacilli, and Micrococci (probably Deinococci) have been isolated in these experiments. We have also isolated novel deinococcal species high altitude (Yang et al. 2010, 2011). These spores and Deinococci are known by their extremely high resistance against UV, gamma ray, and other radiation. If microbes could be found present even at the higher altitude of low earth orbit (400km), the fact would endorse the possibility of interplanetary migration of terrestrial life.

On the other hand, from the viewpoints of chemical evolution for study of origin of terrestrial life, where is the home of organic compounds which might have become precursors of materials such as protein and nucleic acids. Recent studies suggest that the some of such organic compounds were created in space. Then, they reached the surface of Earth via meteorites, cosmic dusts, and so on. One of problems to study such materials of extraterrestrial origin is contamination of materials of terrestrial origin. Avoiding contamination of terrestrial materials from the extraterrestrial materials is quite important issues for this kind of study. Capturing such extraterrestrial materials before falling down on the surface of Earth might be one of possible solutions.

TANPOPO, Japanese name of dandelion, is a plant species, whose seeds with floss are spread by wind. We propose this mission to examine possible interplanetary migration of microbes, and organic compounds on Japan Experimental Module (JEM) of the International Space Station (ISS) (Yamagishi et al. 2008). Ultra low-density aerogel will capture micrometeoroid and space debris. Particles captured by aerogel will be analyzed after the initial curation of the aerogel and tracks in it. Careful curation of the tracks in the aerogel will provide information on the size and velocity of meteorites captured. The particles will be characterized in terms of mineralogical, organic and microbiological properties. The aerogel with low density and layered structure is ready for production in Japan.

In addition to particle-capture on ISS, we also proposed direct exposure experiments of microorganisms and organic compounds with/without model-clay materials that might protect microorganisms and organic compounds from UV and cosmic ray. Spore of *Bacillus* sp., *Deionococcus radiodurans*, and novel Deionococoal species isolated from specimen collected from high altitude by us are candidate subjects for exposure. Amino acids and complex organic compounds that can be formed in space are planed to be exposed.

All the analytical techniques are ready to conduct the TANPOPO mission. Our proposal was accepted as a candidate experiments on Exposed Facility of ISS-JEM. In this paper, we discuss current status of exposure/capture experiments of microorganisms defined for the TANPOPO mission.

#### References

Arrhenius, S. (1908) Worlds in the Making-the Evolution of the Universe (translation to English by H. Borns) Harper and Brothers Publishers, New York.

Crick, F. (1981) Life Itself. Simon & Schuster, New York.

Yamagishi, A., et al. (2008) TANPOPO: astrobiology exposure and micrometeoroid capture experiments. International Symposium on Space Technology and Science (ISTS) Web Paper Archives. 2008-k-05

# **Japan Geoscience Union Meeting 2012** (May 20-25 2012 at Makuhari, Chiba, Japan)

## ©2012. Japan Geoscience Union. All Rights Reserved.



BAO01-03

会場:202

時間:5月24日09:30-09:45

キーワード: 国際宇宙ステーション Keywords: International Space Station