

雲の特性がもたらす気候変化についての数値シミュレーション A numerical simulation study of climate change induced by changes in cloud properties

大淵 済^{1*}, 力石 祐介², 鈴木 遼平², 草野 完也³, 田口 文明¹, 片岡 龍峰², 宮原 ひろ子⁴, 丸山 茂徳², 浜野 洋三¹
OHFUCHI, Wataru^{1*}, CHIKARAISHI, Yusuke², SUZUKI, ryohei², KUSANO, Kanya³, Bunmei Taguchi¹, KATAOKA, Ryuho²,
MIYAHARA, Hiroko⁴, MARUYAMA, Shigenori², HAMANO, Yozo¹

¹ 海洋研究開発機構, ² 東京工業大学, ³ 名古屋大学, ⁴ 東京大学

¹Japan Agency for Marine-Earth Science and Technology, ²Tokyo Institute of Technology, ³Nagoya University, ⁴The University of Tokyo

The terrestrial climate has been changing since the Earth's birth. There are some important factors to trigger the climate change. The faint young sun, land-sea configuration and atmospheric composition are examples of the important factors. Another important factor may be cloud properties. Cloud droplet size, cloud water/ice content, and the height and amount of clouds can affect the climate. It has been pointed out that changes in the intensity of galactic cosmic ray may change the cloud properties. The intensity of cosmic ray may be affected by the solar activity.

We have conducted a set of numerical simulations of a three-dimensional global atmosphere-ocean model and a vertically one-dimensional radiative-convective equilibrium model in order to address the effect of cloud properties on the climate. According to our results, the climate cools (warms) as cloud droplet size is reduced (increased) in the atmospheric radiation computations.