Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.

PEM07-11



時間:5月22日13:45-14:00

## 東南アジア及びアフリカの衛星ビーコン観測から明らかにされた赤道スプレッドF 現象の性質 Characteristics of Equatorial Spread-F (ESF) observed with GNU Radio Beacon Receiver (GRBR) in southeast Asia and Africa

山本 衛<sup>1</sup>\*, 萩行憲輔<sup>1</sup>, Sudarsanam Tulasi Ram<sup>2</sup>, Tsunoda Roland<sup>3</sup> YAMAMOTO, Mamoru<sup>1</sup>\*, Kensuke Hangyo<sup>1</sup>, SUDARSANAM, Tulasi Ram<sup>2</sup>, Roland Tsunoda<sup>3</sup>

<sup>1</sup> 京都大学生存圈研究所, <sup>2</sup>EGRL, Indian Institute of Geomagnetism, <sup>3</sup>SRI International <sup>1</sup>Research Institute for Sustainable Humanosphere, Kyoto University, <sup>2</sup>EGRL, Indian Institute of Geomagnetism, <sup>3</sup>SRI International

Equatorial spread F (ESF) is intense ionospheric irregularity that occurs around the geomagnetic equator. It can cause intense scintillation to satellite-ground communications, and serious error in the GPS measurements. The ESF has been a hot research topic of the equatorial/low-latitude ionosphere for long time. However, its day-to-day variability is not well understood. Now we deploy a very wide network of GNU Radio beacon receivers (GRBR) at low latitudinal regions over east Africa, southeast Asia, and Pacific region, and observe 150MHz/400MHz beacon signal from C/NOFS and other polar-orbiting satellites. In this paper, we use data from Bac Lieu, Vietnam (9.29N, 105.71E, Dip Lat. 1.67N, observations started in January 2009) and Bahirdar, Ethiopia (11.56N, 37.38E, Dip Lat. 3.93N, observations started in March 2011). We discuss relationships between day-to-day variability of ESF with ionospheric structures, i.e., large-scale wave structure (LSWS), meridional symmetry of the ionosphere density distribution, and VHF/UHF scintillation intensity. Occurrence of ESF is well correlated with enhanced LSWS events in the evening time. At Bahirdar, scintillation level is high, and very intense LSWS events are found. From statistical comparison between Africa and southeast Asia, we found that scintillation level is higher in Africa than in southeast Asia, inferring more enhanced occurrence of the ESF over African region.

キーワード: 衛星ビーコン観測, 赤道スプレッド F 現象, 低緯度電離圏, アフリカ, 東南アジア Keywords: Satellite beacon experiment, Equatorial spread-F, Low-latitude ionosphere, Africa, Sroutheast Asia