Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



PEM07-13

会場:304

## negative SIの数値シミュレーション A numerical simulation of a negative solar wind impulse

藤田 茂<sup>1\*</sup>, 田中 高史<sup>2</sup>, 村田 健史<sup>3</sup>, 田 光江<sup>3</sup> FUJITA, Shigeru<sup>1\*</sup>, TANAKA, Takashi<sup>2</sup>, MURATA, Ken T.<sup>3</sup>, DEN, Mitsue<sup>3</sup>

## 1 気象大学校, 2 九州大学, 3 情報通信研究機構

<sup>1</sup>Meteorological College, <sup>2</sup>Kyushu University, <sup>3</sup>National Institute of Information and Communication Technology

Response of the magnetosphere-ionosphere system to the negative impulse of the solar wind dynamic pressure (the negative SI) are studied again with foci of periodic variations of the ionospheric convection and appearance of the overshielding potential. When the negative impulse impinges on the magnetopause, the Region 1 (R1)-type field-aligned current (FAC) and R2-type FAC appears alternatively in the lower-latitudes ( $^{\sim}$  70 deg) of the dayside ionosphere ( $^{\sim}$ 10hLT and 14hLT). These current systems shift nightward and poleward. This alternative appearance of FACs invokes positive and negative ionospheric potential patterns switching alternatively. In addition, the R2-type FAC induced by the negative SI and that by the positive SI tend to yield the shielding electric potential in the ionosphere. This shielding potential has short duration than that for the northward turn of the interplanetary magnetic field does. The duration is longer for the negative SI than for the positive SI.

キーワード: negative SI, 数値シミュレーション, 磁気圏電離圏複合系, 対流変動

Keywords: negative SI, numerical simulation, magnetosphere-ionosphere compound system, convection oscillation