

## Recurrent and propagating SAPS/ SAID structures observed by the SuperDARN Hokkaido radar

## Recurrent and propagating SAPS/ SAID structures observed by the SuperDARN Hokkaido radar

西谷 望<sup>1\*</sup>, 小川 忠彦<sup>2</sup>, Sasha Koustov<sup>3</sup>, 菊池 崇<sup>1</sup>, 海老原 祐輔<sup>4</sup>, 片岡 龍峰<sup>5</sup>, 堀 智昭<sup>1</sup>, 細川 敬祐<sup>6</sup>, 寺本 万里子<sup>1</sup>  
NISHITANI, Nozomu<sup>1\*</sup>, OGAWA, Tadahiko<sup>2</sup>, Sasha Koustov<sup>3</sup>, KIKUCHI, Takashi<sup>1</sup>, EBIHARA, Yusuke<sup>4</sup>, KATAOKA, Ryuhō<sup>5</sup>,  
HORI, Tomoaki<sup>1</sup>, HOSOKAWA, Keisuke<sup>6</sup>, TERAMOTO, Mariko<sup>1</sup>

<sup>1</sup>Solar-Terrestrial Environment Laboratory, Nagoya University, <sup>2</sup>National Institute of Information and Communications Technology, <sup>3</sup>University of Saskatchewan, <sup>4</sup>Research Institute of Sustainable Humanosphere, Kyoto University, <sup>5</sup>Tokyo Institute of Technology, <sup>6</sup>University of Electro-Communications

<sup>1</sup>Solar-Terrestrial Environment Laboratory, Nagoya University, <sup>2</sup>National Institute of Information and Communications Technology, <sup>3</sup>University of Saskatchewan, <sup>4</sup>Research Institute of Sustainable Humanosphere, Kyoto University, <sup>5</sup>Tokyo Institute of Technology, <sup>6</sup>University of Electro-Communications

We study two events of the recurrent propagating SAPS/SAID structures at magnetic latitudes of  $\sim 60^\circ$ , which were observed by the Hokkaido radar at 0830 - 0930 UT (1730 - 1830 MLT) on April 23, 2008 and at 1100-1500 UT on March 1, 2011. Both events occurred during the main phases of moderate geomagnetic storms with minimum Dst about -37 nT and -75 nT respectively. The IMF was southward and turned to northward just prior to the first event, whereas it was constantly southward for the second event. For both events the period, velocity of westward progression and wavelength are 30 min, 700 m/s and 1200 km respectively. They have similar characteristics to those of the 'auroral giant undulation' at the equatorward edge of the auroral oval although some numbers, for example the periods, are slightly different. The radar data presented strongly support an idea that the recurrent SAPS / SAID structures are generated through internal processes in the inner magnetosphere, due to probably some kind of plasma instability, rather than due to externally-driven processes.

キーワード: 北海道-陸別 HF レーダー, SuperDARN, SAPS / SAID, 内部磁気圏プラズマ不安定

Keywords: Hokkaido HF radar, SuperDARN, SAPS / SAID, plasma instability in the inner magnetosphere