

不均一ヘイローとスプライトストリーマの関係

Relationship between structures of sprite streamers and inhomogeneity of preceding halos

三宮 佑介^{1*}, 高橋 幸弘¹, 佐藤 光輝¹, 工藤 剛史¹, 小林 縫², 山田 大志², 島 侑奈¹, NHK 宇宙の渚プロジェクト³, Hans Nielsen⁴, Matthew G. McHarg⁵, Ryan Haaland⁶, Takeshi Kammae⁷, Steven Cummer⁸, Yoav Yair⁹, Walter Lyons¹⁰, SANMIYA, Yusuke^{1*}, TAKAHASHI, Yukihiko¹, SATO, Mitsuteru¹, KUDO, Takeshi¹, KOBAYASHI, Nui², Taishi Yamada², SHIMA, Yuuna¹, NHK Cosmic Shore Project³, Hans Nielsen⁴, Matthew G. McHarg⁵, Ryan Haaland⁶, Takeshi Kammae⁷, Steven Cummer⁸, Yoav Yair⁹, Walter Lyons¹⁰

¹ 北海道大学大学院理学院宇宙理学専攻, ² 北海道大学理学部地球惑星科学科, ³ 日本放送協会, ⁴ Univ. of Alaska, Fairbanks, AK, United States, ⁵ US Airforce Academy, Colorado Springs, CO, United States, ⁶ Fort Lewis College, Durango, CO, United States, ⁷ Univ. of Alaska, Fairbanks, AK, United States, ⁸ Duke University, Durham, NC, United States, ⁹ Open University of Israel, Ra'anana, Israel, ¹⁰ FMA Research, Ft Collins, CO, United States

¹ Department of CosmoSciences, Hokkaido University, ² Department of Earth Sciences, School of science, Hokkaido University, ³ Japan Broadcasting Corporation, ⁴ Univ. of Alaska, Fairbanks, AK, United States, ⁵ US Airforce Academy, Colorado Springs, CO, United States, ⁶ Fort Lewis College, Durango, CO, United States, ⁷ Univ. of Alaska, Fairbanks, AK, United States, ⁸ Duke University, Durham, NC, United States, ⁹ Open University of Israel, Ra'anana, Israel, ¹⁰ FMA Research, Ft Collins, CO, United States

Transient Luminous Events (TLEs) in the stratosphere, mesosphere and lower thermosphere is phenomena of high time resolution associated with lightning. The diffuse glows such as elves and sprite halos progresses first and discrete structure of sprite streamers progresses next. Elves and sprite halos are considered to have a relationship with structure of sprite streamers. Therefore, spatial distribution and time evolution of these phenomena is key parameters of the condition of generation and mechanism of TLEs.

In the period of June 27 - July 10, 2011, aircraft and ground-based campaign in support of NHK Cosmic Shore project was carried out under collaboration between NHK, Japan Broadcasting Corporation, and universities. Two aircrafts carried the high-speed camera, and observed the TLEs.

The observation from an aircraft makes it possible to capture TLEs without influence of atmospheric absorption, and detailed structures of TLEs over 40 events was able to observe with the high-speed camera with frame rate of 8300 /sec.

In the TLEs observed by the campaign, sprite streamers and sprite halos are always showing some inhomogeneity, and it seemed that sprite streamers is clearly related to inhomogeneity of sprite halos. At this presentation, we will show the relationship between inhomogeneity of halo and sprite streamers with these velocities of progress estimated from the high-speed images.

キーワード: スプライト, ヘイロー, ストリーマ, ハイスピードカメラ

Keywords: sprite, halo, streamer, high-speed camera