

Akebono 衛星で観測される THR 放射の出現頻度について The occurrence rate of THR emissions observed by the Akebono satellite

佐藤 由佳^{1*}, 小野 高幸², 熊本 篤志¹
SATO, Yuka^{1*}, ONO, Takayuki², KUMAMOTO, Atsushi¹

¹ 東北大学大学院理学研究科惑星プラズマ・大気研究センター, ² 東北大学大学院理学研究科地球物理学専攻
¹Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, ²Department of Geophysics, Graduate School of Science, Tohoku University

This is a report on the occurrence rate of Terrestrial Hectometric Radiation (THR), auroral radio emissions emanating from the topside ionosphere in the MF and HF ranges. Data shown in this paper were obtained by the Plasma waves and sounder experiment (PWS) mounted on the Akebono satellite. The Akebono/PWS measurements show that THR emissions are sometimes observed in two frequency bands near 1.5-2.0 MHz and 3.0-4.0 MHz when the satellite passes over the auroral latitudes; however, their occurrence rate has not yet been investigated. Statistical studies using the Akebono/PWS data show the spatial distribution of THR occurrence; THR is detected at all magnetic local times and most often during premidnight hours (2100-2400 MLT) in a wide magnetic latitude range ($|\text{MLAT}| > 30\text{deg}$). During daytime hours (0600-1500 MLT), the distribution of its occurrence is confined in higher latitude ranges ($|\text{MLAT}| > 70\text{deg}$). The explanation of this spatial distribution is that THR is favorably generated in the night-side auroral latitudes near 1000-km altitude.

キーワード: オーロラ現象, オーロラ電波, 電波伝搬, オーロラ電離圏

Keywords: auroral phenomena, auroral radio emissions, radio propagation, auroral ionosphere