

## Magnetic conjugacy of northern and southern auroral beads

MOTOBA, Tetsuo<sup>1</sup>, HOSOKAWA, Keisuke<sup>2\*</sup>, Akira Kadokura<sup>1</sup>, Natsuo Sato<sup>1</sup>

<sup>1</sup>National Institute of Polar Research, <sup>2</sup>University of Electro-Communications

Auroral beads, i.e. azimuthally arrayed small-scale bright spots resembling a pearl necklace, have recently drawn the attention of researchers as a possible precursor of explosive activation of the aurora. Here we used simultaneous, ground-based, all-sky camera observations from a geomagnetically conjugate Iceland-Syowa Station pair to demonstrate that small-scale auroral beads evolve synchronously in the northern and southern hemispheres and have good magnetic conjugacy for  $\sim 7$  min before an auroral breakup. The synchronous conjugate auroral beads undergo a two-step evolution: in the first  $\sim 4$  minutes, well-organized bead structures move eastward with an almost constant speed of 1 km/s or less, and subsequently they develop dramatically into brighter and larger auroral forms with faster propagation speed of 2-6 km/s. Our observations strongly suggest that, for the auroral beads, the magnetosphere plays a fundamental role in the determining their temporal evolution, while the ionospheric contribution that can induce asymmetric auroral behavior in the two hemispheres is minor or not significant.

Keywords: auroral beads, magnetic conjugacy, substorm