

Volatile partitioning and formation of core, mantle, and atmosphere on early Titan Volatile partitioning and formation of core, mantle, and atmosphere on early Titan

関根 康人^{1*}, 渋谷 岳造², 桑谷 立¹, 洪 鵬¹, 鈴木 勝彦²

SEKINE, Yasuhito^{1*}, SHIBUYA, Takazo², KUWATANI, Tatsu¹, HONG, Peng¹, SUZUKI, Katsuhiko²

¹Dept. Complexity Sci. & Engr., Univ. of Tokyo, ²PEL, JAMSTEC

¹Dept. Complexity Sci. & Engr., Univ. of Tokyo, ²PEL, JAMSTEC

Titan is a complex body with a thick atmosphere composed mainly of N₂ and CH₄, relatively young surface with lakes of liquid hydrocarbons, ice mantle that holds an interior liquid ocean, and low density core. The core may consist of either a mixture of ice and rock or hydrous silicates, suggesting that Titan's interior would have been much colder than previously thought (Iess et al., 2009; Fortes et al., 2012). However, such a view of a cold interior seems to be inconsistent with evidence for extensive geological activities near the surface inferred from the young surface age (Neish and Lorenz, 2012) and presence of radiogenic Ar degassed from the interior (Niemann et al., 2005). In addition, the findings of significant amounts of CO₂ in Enceladus plumes (Waite et al., 2009) indicate that the chemical composition of the building blocks of Titan may have also been a comet-like one with a few percents of CO₂ content relative to H₂O. Nevertheless, the presence of such CO₂ in Titan's proto-atmosphere and mantle should have resulted in the formation of large amounts of atmospheric CO in addition to N₂ through photolysis (Atreya et al., 1978), shock heating (McKay et al., 1988; Ishimaru et al., 2011), and hypervelocity impacts (Sekine et al., 2011). Although the Cassini spacecraft has provided important clues to understand the origin and evolution of Titan, there is no unified view that accounts for the complex nature of the core, mantle, and atmosphere consistently.

Here we investigate partition and chemical evolution of primordial CO₂, NH₃ and CH₄ in a water magma ocean that is considered to have been formed during Titan's accretion (Kuramoto and Matsui, 1994). On the basis of laboratory experiments, we show that primordial CO₂ would have been readily converted to carbonate minerals in the water magma ocean through hydrothermal reactions with primitive minerals, such as olivine. Thus, CO₂ would have been partitioned mainly into the core by forming a low density carbonate during accretion. The proto-atmosphere and mantle conversely would have been highly depleted in CO₂, preventing the formation of a thick CO atmosphere. On the other hand, our experimental results indicate that NH₃ would not have been converted into N₂ in the water magma ocean due to the efficient formation of H₂ through serpentinization, suggesting that NH₃ has been partitioned mainly in the mantle and proto-atmosphere. Given the low solubility of CH₄ in the water magma ocean, CH₄ is considered to have been partitioned mainly in the proto-atmosphere. Because of the thermodynamical stability of carbonate at high temperatures (e.g., 2000 K) under high pressure conditions corresponding to Titan's core (Isshiki et al., 2004), the carbonate-rich, low density core allows a warm interior of Titan, consistent with the evidence for extensive surface geological activities. Our model provides a new hypothetical starting point for the evolution of interior and atmosphere-surface systems of Titan and other large icy bodies.

キーワード: Titan, partitioning, hydrothermal reactions, icy satellites, core, atmosphere

Keywords: Titan, partitioning, hydrothermal reactions, icy satellites, core, atmosphere