Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.

PPS02-11



時間:5月24日16:30-16:45

木星探査計画 Jupiter Magnetospheric Orbiter に向けた X 線撮像・分光 装置の検討 Study of the Jupiter X-ray imaging spectrometer on Jupiter Magnetospheric Orbiter

木村 智樹¹*, 江副 祐一郎², 笠原 慧¹, 三好 由純³, 山崎 敦¹, 藤本 正樹¹ KIMURA, Tomoki¹*, EZOE, Yuichiro², KASAHARA, Satoshi¹, MIYOSHI, Yoshizumi³, YAMAZAKI, Atsushi¹, FUJIMOTO, Masaki¹

¹ 宇宙航空研究開発機構宇宙科学研究所,² 首都大学東京理工学研究科,³ 名古屋大学太陽地球環境研究所 ¹JAXA/ISAS, ²Graduate school of science and engineering, Tokyo Metropolitan University, ³Solar-Terrestrial Environment Laboratory, Nagoya University

In 2000's, the new generation X-ray observatories (Chandra, XMM-Newton and Suzaku) have revealed various new X-ray phenomena in the Jupiter system. The detected objects include Jupiter's aurorae, disk (middle and low-latitude emission), Io, Europa, the Io Plasma Torus, and radiation belts. For example, Jupiter's aurorae emit time variable X-rays via bremsstrahlung by keV electrons and charge exchange by MeV ions (Gladstone et al. 2002 Nature). A diffuse X-ray emission associated with the Jupiter's radiation belts suggests an inverse Compton scattering of tens MeV electrons (Ezoe et al. 2010 ApJ). Hence, the X-ray emission can be a unique diagnostic tool to investigate key fundamental problems on the Jupiter system such as the relativistic particle acceleration and the Jupiter-satellite interaction. However, since these observations have been done with the X-ray astronomy satellites orbiting the Earth, the photon statistics of X-ray spectra and light curves, and the angular resolution of X-ray images were severely limited. In this context, we have started to study design of an X-ray imaging spectrometer for JMO (Jupiter Magnetospheric Orbiter) which is a Jupiter exploration mission led by JAXA descended from international collaboration with JUICE (JUpiter ICy moon Explorer) mission led by ESA in 2020's. JMO is expected to perform high-latitude (10-30 deg inclination) measurements of the Jupiter system and overview the magnetospheric activities. The in-situ measurements by JMO provide us with an unprecedented opportunity to observe Jupiter with extremely high photon statistics, high time and angular resolution. To realize the in-situ X-ray instrument for JMO, stringent mass and power limitations must be fulfilled. Furthermore, the radiation and the contamination of optical lights and debris must be taken care. The base line is a combination of an original ultra-light weight X-ray telescope with the mass to area ratio of 10 kg/m^2 based on the micromachining technologies (Ezoe et al. 2010 Microsystem technologies), and a radiation-hard and low-power DepFET imaging detector (Struder et al. 2010 SPIE). In this presentation, the scientific objectives and current status of instrumental study will be presented.

キーワード:木星,磁気圏,探査,X線,粒子加速

Keywords: Jupiter, magnetosphere, exploration, X-ray, particle acceleration