

奄美大島に分布する中生代付加体の碎屑性ジルコン年代から見た後背地の変遷 Temporal Change of Provenance detected by detrital zircon chronology for the basement complex in Amami-Oshima Island, Ry

木林 和美^{1*}

KIBAYASHI, Kazumi^{1*}

¹ 広島大学大学院理学研究科

¹graduate school of science.hiroshima university

Basement geology of Amami-Oshima Island is composed of Yuwan Complex and Naon Complex of Southern Chichibu Terrane as Late Jurassic to earliest Cretaceous accretionary prism, Shinkogachi Unit, Yakugachi Unit and Naze Unit of Shimanto Terrane as Early Cretaceous accretionary complex, and Wano Formation as Paleogene fore-arc cover. Detrital zircon U-Pb chronology by LA-ICP-MS reveals the temporal change of their provenance. Age populations of detrital zircon from each of the above units are summarized as bellows.

Yuwan Complex shows the predominant age clusters at around 185 Ma, 250 Ma, 1890 Ma and 2300 Ma with the youngest age cluster around 160 Ma. Naon Complex shows the predominant age clusters at around 175 Ma, 220 Ma, and subordinate cluster at around 140 Ma, 420 Ma, 1850 Ma, and 2200-2300 Ma. Shinkogachi Unit shows the predominant age cluster at around 100 Ma, 175 Ma, and 190 Ma with subordinate cluster at around 370 Ma and 1850-2300 Ma. Yakugachi Unit shows the predominant age clusters at around 120 Ma and 165 Ma, and subordinate clusters around 95 Ma, 270 Ma and 1605-1900 Ma. Naze Unit shows the predominant age cluster at around 65 Ma, 120 Ma, 170 Ma, and 215 Ma, and subordinate cluster of 1630-1950 Ma. Wano Formation shows predominant age clusters at around 55 Ma, 75 Ma, 185 Ma and 255 Ma, and subordinate clusters around 220 Ma and 1850 Ma.

Appearance and absence of each age clusters are different with respect to each unit, while the all unites predominate the early to middle Jurassic age clusters. This shows that the provenance change was happened to occur during the period of their formation. As a wholly, clear appearance of early Proterozoic zircon and the absence of late Proterozoic zircon may show the provenance of North China craton. Combining with whole rock chemistry of sandstones reveals that their provenances changed from continental arc side for the Southern Chichibu Terrane to island arc side for the Shimanto Terrane, and then after back to the continental arc side again for the Naze Unit. The island arc can be postulated to form during Late Triassic time shown by the predominance of Ca. 220 Ma zircons in the Naon complex.

キーワード: 奄美大島, 秩父帯, 四万十帯, 碎屑性ジルコン, レーザーアイシーピーマス

Keywords: Amami-Oshima, Chichibu Terrane, Shimanto Terrane, detrital zircon, LA-ICP-MS