

隠岐島後・吉備の捕獲岩ジルコン年代から推定される西南日本下部地殻の発達過程 Evolution of the lower crust under southwest Japan constrained from ages of zircon in the xenolith

越田 溪子^{1*}, 岩森 光¹, 平田 岳史²

KOSHIDA, Keiko^{1*}, IWAMORI, Hikaru¹, HIRATA, Takafumi²

¹ 東京工業大学地球惑星科学専攻, ² 京都大学大学院理学研究科地球惑星科学専攻

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²Graduate School of Science, Kyoto University

The present-day continental crust contains a very large proportion, about 20-70%, of the incompatible elements by mass in the Earth (Rudnick and Fountain, 1995). Therefore, it should have enormous effects on evolution of the Earth both chemically and physically. Although the upper crust has been investigated extensively in terms of its composition, structure, and evolution process, those of the lower crust which extends to 35 km depth on average has been poorly investigated.

The main purpose of this study is to decipher process of formation and evolution of the lower crust, aiming at understanding the evolution of continental crust and the Earth. As a first case study, we try to constrain the evolution process of the lower crust in SW Japan, the Oki-Dogo and Kibi areas, based on zircon ages and geothermometers in the xenoliths.

These two areas are located on the continental margin, and may provide useful insights regarding how the lower crust is formed or destructed associated with subduction.

At Oki-Dogo, peridotite, pyroxinite, gabbro and granulite were found as xenoliths in alkali olivine basalts (Takahashi, 1978) erupted at 3.61Ma (whole rock K-Ar age, Kaneoka et al., 1977). At Kibi, peridotite, pyroxinite and granulite are found as xenoliths in alkali olivine basalts (Iwamori, 1985) erupted at around 9 Ma (Uto, 1989).

To estimate the equilibrium temperature and the U-Pb age of zircon, we sampled gabbros from Oki-Dogo and granulites from Kibi, and analyzed them by using EPMA and LA-ICP-MS.

Major constituent minerals in the gabbros from Oki-Dogo are olivine (ol), clinopyroxene (cpx) and plagioclase (pl), and they have a diameter of 1~2mm. Equilibrium temperature estimated by ol-cpx geothermometer (Loucks, 1996) is approximately 1100^o+50^oC. Zircon grains have been obtained from only one gabbro (out of total seven gabbros processed), and are almost anhedral and homogeneous. The total forty grains have been dated to give ages approximately ranging from 2 to 4 Ma, which are broadly the same with that of the host alkali basalt and indicates that zircon grains lost almost all Pb during the magmatic event exceeding the closure temperature for zircon U-Pb dating (i.e., above 900^oC).

Major constituent minerals in the granulite from Kibi quartz, K-feldspar, garnet, kyanite and spinel are the major constituent minerals, and they have a diameter of 500~1000 um. Based on the stability of kyanite and assuming the maximum depth of 30 km based on the present-day Moho depth estimated from the seismic profile (Ito et al., 2010), the equilibrium temperature is constrained to be less than 800^oC. Zircons obtained from the granulites are various in shape, exhibiting a wide age range from 420 to 10 Ma. We classified these zircons into igneous and metamorphic origins based on the U/Th ratio, and distinguish the overlapping events recorded in the grains. Bulk rock composition of this granulite from Kibi is aluminous and pelitic (Kushiro, 1987). From these results, we propose a model that a sedimentary material subducted with an oceanic plate, and accreted the material to the continental crust. Then at 28Ma, a part of the Philippine Sea Plate started spreading to create the Shikoku Basin and the spreading ridge had subducted. Because of this, the subduction angle became gentler and accrete more materials and push the formerly accreted sedimentary material further to the reararc region at the same time, accreted prism was metamorphosed by the heat.

In summary, these two examples from Oki-Dogo and Kibi suggest that (1) the lower crust beneath Oki-Dogo were heated by magmatic events that erupt alkali basalts and may reset the U-Pb age of zircon now found in the xenolith, and (2) subducted sedimentary materials can accrete to the lower crust at deep levels, which may be promoted by ridge subduction. Therefore, at this area, a part of lower crust develop independently of the upper crust.

キーワード: 下部地殻, ジルコン, 年代, 捕獲岩, 西南日本

Keywords: Lower Crust, zircon, age, xenolith, southwest Japan