## **Japan Geoscience Union Meeting 2012**

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



SGD24-02 Room:105 Time:May 25 09:15-09:30

## Gravity field determination around Syowa station, Antarctica, by combining GOCE and in-situ gravity data

FUKUDA, Yoichi<sup>1\*</sup>, NOGI, Yoshifumi<sup>2</sup>, MATSUZAKI, Kazuya<sup>1</sup>

<sup>1</sup>Graduate School of Science, Kyoto University, <sup>2</sup>National Institute of Polar Research

GOCE (Gravity field and steady-state Ocean Circulation Explorer) satellite launched in March 2009 by ESA (European Space Agency) aims at improving static gravity fields, in particular at short wavelengths. In addition to its low-altitude orbit (250km), the sensitive gravity gradiometer installed is expected to reveal 1 mgal gravity anomaly and 1cm geoid at the spatial resolution of 100km (half wavelength). On the other hand, due to instrumental drifts, lack of reference points, and other reasons, the accuracy of in-situ gravity data (land, surface ship and airborne gravity data) is decreasing toward the longer wavelength more than several tens km. In particular in Antarctica where very few gravity reference points are available, the long wavelength accuracy and/or consistency among the data sets are quite limited. The Japanese Antarctic Research Expedition (JARE) has been conducting in-situ gravity measurements around the Japanese Antarctic stations for a long period. These measurements also suffered from such influences and they cause large errors in the long wavelength gravity fields, and, consequently, errors in geophysical and geodetic applications. This study aims at improveing the accuracy of the JARE gravity data using GOCE gravity models (level 2 EGMs). There are three different approaches for estimating the GOCE gravity models, namely, direct solution (DIR), timewise solution (TIM) and space-wise solution (SPW). Among these, TIM never uses any a-priori information other than GOCE. Therefore we mainly employed TIM models (RL 1-3). We also employed EGM2008 as a reference. In this talk, we present the comparisons between the gravity models and each of in-situ gravity data sets, and a preliminary result of the improved gravity field around Syowa station, Antarctica.

Keywords: GOCE, Gravity Model, Antarctica, Syowa Station