Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



SSS01-03

会場:104

時間:5月25日09:30-09:45

短期クラスタリングの ETAS モデルに基づく長期地震予測 Long-term earthquake forecasts based on the ETAS model for short-term clustering


庄 建倉 ^{1*} ZHUANG, Jiancang ^{1*}

1 統計数理研究所

Based on the ETAS (epidemic-type aftershock sequence) model, which is used for describing the features of short term clustering of earthquake occurrence, this paper presents some theories and techniques related to evaluating the probability distribution of the maximum magnitude in a given space-time window, where the Gutenberg-Richter law for earthquake magnitude distribution cannot be applied directly. It is shown that the distribution of the maximum magnitude in a given space-time volume in long term is determined by the background seismicity rate, the magnitude distribution for all earthquakes, and the structure of earthquake clusters. The introduced techniques were applied to the seismicity in the Japan region in the period from 1926 to 2009. It is found that the regions most likely to have great earthquakes are along the Tohoku (Northeastern Japan) Arc and the Kuril Arc, both much higher probabilities than the offshore Nankai and Tokai regions.

キーワード: ETAS モデル, 常時地震活動, 長期地震予測, 地震確率

Keywords: ETAS model, background seismicity, long term earthquake forecast, earthquake probability



¹Institute of Statistical Mathmematics