

2010年夏における日本の猛暑のメカニズム Mechanisms of the 2010 summer heat wave in Japan

澁谷 亮輔^{1*}, 岡島 悟³, 末善 健太², 仲谷 幸浩², 東塚 知己¹, 西井 和晃³, 宮坂 貴文³, 中村 尚³

SHIBUYA, Ryosuke^{1*}, Satoru Okajima³, SUEYOSHI, Kenta², NAKATANI, Yukihiko², TOZUKA, Tomoki¹, NISHII, Kazuaki³, MIYASAKA, Takafumi³, NAKAMURA, Hisashi³

¹ 東京大学大学院理学系研究科地球惑星科学専攻, ² 東京大学地震研究所, ³ 東京大学先端科学技術研究センター

¹Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, ²Earthquake Research Institute, The University of Tokyo, ³Research Center for Advanced Science and Technology, The University of Tokyo

Japan experienced an extraordinary heat wave in the summer of 2010 with highest average temperature ever since 1898. This may be linked with the strong positive sea level pressure (SLP) anomaly just to the south of Japan. To identify its cause, a series of atmospheric general circulation model simulations are conducted. Although the tropical Pacific is known as one of the major sources of abnormal weather in Japan, it did not play a significant role during this summer. Rather, the sea surface temperature (SST) anomaly in the tropical Indian and Atlantic Oceans played an important role in forming the positive SLP anomaly. It is suggested that anomalously high SST in the tropical Indian Ocean after the 2009/10 El Nino played a role in inducing an anomalous anticyclone over the northwestern Pacific. On the other hand, the positive SST anomaly in the tropical Atlantic induces more convection and diabatic heating, which in turn, becomes a source of Rossby wave-train along the Asian jet in the upper troposphere that generate positive SLP anomalies around Japan.

キーワード: 大西洋熱帯域, ロスビー波, 異常気象, 亜熱帯ジェット, 猛暑

Keywords: tropical Atlantic Ocean, Rossby wave, abnormal weather, Asian jet, heat wave