

南極におけるインフラサウンド波：気候・環境モニタリングの新指標 Infrasound Waves in Antarctica: A New Proxy for Climate and Environmental Monitoring

石原 吉明¹, 山本 真行², 金尾 政紀^{3*}

ISHIHARA, Yoshiaki¹, YAMAMOTO, Masa-yuki², KANAOKI, Masaki^{3*}

¹ 国立天文台, ² 高知工科大学, ³ 国立極地研究所

¹National Astronomical Observatory of Japan, ²Kochi University of Technology, ³National Institute of Polar Research

The Infrasound is defined as sub-audible sound with frequencies ranging from the cut-off frequency of sound (3.21 mHz) to the lowest frequency of the human audible band (20 Hz). This frequency range presents new opportunities for remote sensing of the Earth's atmospheric physical environment (Hedlin et al., 2002). There is an emerging body of published results on infrasound waves generated by diverse sources, such as volcanic eruptions, ocean waves, earthquakes, thunders, sprites and airplane passages (Matoza et al, 2007; Garces et al., 2008). For a remarkable example, the 2011 Tohoku-Oki, Japan earthquake (Mw = 9.0) produced unequivocal infrasound signals associated with the large tsunami (Arai et al., 2011). Free oscillations of the Earth from great earthquakes can also affect even the upper atmosphere. Another striking example is given by infrasound shock waves generated by a large meteorite fall in Japan (Ishihara et al, 2004).

Characteristic features of infrasonic waves associated with environmental changes are clearly identified in Antarctic stations deployed during the International Polar Year (IPY2007-2008). Loading effects from the Southern Ocean are recorded by infrasound sensors deployed at Syowa Stations (SYO; 69S, 39E), in the Lutzow-Holm Bay, East Antarctica. The oceanic effects on infrasound modulated by the presence of sea-ice are examined relating to atmosphere-ocean-cryosphere system. Measurements of these microbaroms are useful tools for characterizing ocean wave climate and global storm intensity, complementing other geophysical measurements. The infrasound data at SYO, moreover, demonstrate unique signals with harmonic over-tones that may be influenced by local near-surface processes, such as katabatic winds or the ice dynamics. Infrasound observations in Antarctica, consequently, could present unique new proxies for monitoring environmental change as well as temporal climate variations in the southern polar region.

キーワード: infrasound, Antarctica, microbaroms, environmental monitoring, global network, atmosphere-ocean interaction

Keywords: infrasound, Antarctica, microbaroms, environmental monitoring, global network, atmosphere-ocean interaction