

氷期中の数千年スケールの急激な気候変化とその氷床への影響に関するモデリング Modelling the Abrupt climate change in millennial scale and its influence upon ice sheets during the middle level glacia

阿部 彩子^{1*}, 斎藤冬樹², 岡頭¹, 大垣内るみ², 高橋邦夫²

ABE-OUCHI, Ayako^{1*}, Fuyuki Saito², Akira Oka¹, Rumi Ohgaito², Kunio Takahashi²

¹ 東京大学大気海洋研究所, ²JAMSTEC

¹AORI, University of Tokyo, ²JAMSTEC

Abrupt climate change in millennial time scale such as D-O cycles and AIM recorded in ice cores occurs more frequent with high amplitude during the middle level glacial climate state than in the interglacial state or the full glacial state. The mechanism of the frequent occurrence of abrupt change through the Atlantic meridional overturning circulation is unclear and the necessity of the high frequency during the middle level glacial state is not known. Here we use a coupled ocean atmosphere model, MIROC, to compare the detailed nature of the response to fresh water release (0 to 0.1 Sv under different initial condition) of AMOC under middle level glacial state with interglacial or full glacial state. Under middle level glacial state, the AMOC is stronger at the basic state (without water release), nearly switches off with small amount of fresh water release of 0.05 Sv and induces larger cooling in the Northern hemisphere than other background climate states. The recovery of AMOC is induced from a nearly switched off AMOC state by reducing the fresh water release to 0.01 Sv or less, lead by the gradual response in the low latitude and followed by the lagged but sudden response in the convection and sea ice area in the North Atlantic. Laurentide and Fenno-Scandian Ice sheets' melt water estimated by an ice sheet model IcIES is consistent to the melt water amount needed for the AMOC change under middle glacial state, showing a possibility of favorable condition of self sustained oscillation between ice sheet and ocean meridional overturning under middle level glacial climate.