

Solar wind control of lunar external magnetic enhancement: A case study

NISHINO, Masaki N.^{1*}, FUJIMOTO, Masaki¹, TSUNAKAWA, Hideo², SHIBUYA, Hidetoshi³, TAKAHASHI, Futoshi², SHIMIZU, Hisayoshi⁴, MATSUSHIMA, Masaki², SAITO, Yoshifumi¹

¹JAXA, ²Tokyo Institute of Technology, ³Kumamoto University, ⁴University of Tokyo

We study an interaction between the solar wind (SW) and the magnetic anomalies on the lunar surface using SELENE (Kaguya) data. It has been known that magnetic enhancements are at times detected near the limb external to the lunar wake, which is thus called lunar external magnetic enhancement (LEME), as a result of direct interaction between the SW and the lunar crustal fields. Previous observational studies, based on statistical trends that stronger interplanetary magnetic field (IMF) and higher SW density favor the LEME in high solar zenith angle (SZA) region, suggested a fluid-type interaction as a candidate for formation mechanism of the LEME. However, neither the IMF orientation nor the crustal field direction has not been taken into account in the previous analyses.

We show evidence that relation between the IMF and crustal field orientation is also one of the key factors that control the extent of LEME, focusing on one-day observations (12 revolutions) that include data above South Pole-Aitken (SPA) basin which is characterized by strong crustal fields in a wide region. Strong LEMEs are detected at 100 km altitude around SPA basin under the stronger and northward IMF condition, while they weakens under southward IMF. We examined the crustal field model (uncompressed by the SW) constructed from the SELENE magnetometer data to know the orientation of the crustal field at 300 km, 100 km, and lower altitude. In the region where the peak of the magnetic enhancement is detected at 100 km altitude, the model crustal field at 300 km altitude is directed northward, while the model field at 100 km and lower altitude had a southward component in some revolutions. This suggests that the lunar crustal field is compressed by the SW dynamic pressure, and that its large scale component is essential to the formation of the LEME. In addition, our results show that pile-up of the IMF above the crustal fields becomes more effective under parallel field configuration, and suggests that magnetic reconnection between the IMF and the lunar crustal field may take place under anti-parallel field configuration.

Keywords: Solar wind-Moon interaction, Lunar magnetic anomalies, Magnetic pile-up, SELENE

Radio scintillation observation of solar corona with Venus orbiter Akatsuki

IMAMURA, Takeshi^{1*}, ANDO, Hiroki², MIYAMOTO, mayu², ISOBE, Hiroaki³, ASAI, Ayumi³, SHIOTA, Daikou⁴, YAJI, Kentaro⁵, TOKUMARU, Munetoshi⁶

¹Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, ²The University of Tokyo, ³Kyoto University, ⁴RIKEN, ⁵Rikkyo University, ⁶STE Lab, Nagoya University

Radio occultation observation of solar corona was conducted in June, 2011 using Japanese Venus probe Akatsuki. The observations covered solar offset distances of 0.5–20.5 solar radii. The amplitude scintillation spectra revealed the nature of disturbances in the inner heliosphere.

Keywords: sun, solar corona, radio wave, Akatsuki, scintillation

Kinematic properties of slow ICMEs and modification of an equation for a drag model

IJU, Tomoya^{1*}, TOKUMARU, Munetoshi², FUJIKI, Ken'ichi²

¹Particle and Astrophysical Science., Nagoya-University., ²Solar-Terrestrial Environment Laboratory, Nagoya-University.

We report kinematic properties of slow interplanetary coronal mass ejections (ICMEs) detected by interplanetary scintillation (IPS) observations, and propose a modified equation for the ICME motion. Understanding of ICME propagation is very important for space weather forecasting. We believe that the radial motion of ICME is governed by a drag force due to interaction with the background solar wind and the magnitude of the force is proportional to the difference between speeds. Earlier studies dealt mainly with propagation of ICMEs moving faster than the solar wind, while observational studies for propagation of slow ICMEs were rare. The IPS observations have been carried out since the early 1980s using the 327MHz radio-telescope system of the Solar-Terrestrial Environment Laboratory, Nagoya University. These observations allow us to probe into the inner heliosphere with a cadence of 24 hours. We take advantage of the IPS observations to determine speeds and accelerations of slow ICMEs. In this study, we identified 59 ICMEs from the IPS observations during 1997-2011. Here, we define that V_{SOHO} , V_{IPS} and V_{bg} are the initial speed of ICME, speed of that in the interplanetary space, and speed of background solar wind, respectively. Using the values of these, we classify the 59 ICMEs into three types of ICMEs, *i.e.*, fast ($V_{SOHO} - V_{bg} > 500 \text{ km s}^{-1}$), moderate ($0 \text{ km s}^{-1} < V_{SOHO} - V_{bg} < 500 \text{ km s}^{-1}$), and slow ($V_{SOHO} - V_{bg} < 0 \text{ km s}^{-1}$). Here, we eliminate ICMEs exhibiting unusual values of $V_{IPS} - V_{bg} > 500 \text{ km s}^{-1}$ for the moderate and of $V_{IPS} - V_{bg} > 100 \text{ km s}^{-1}$ for the slow as events having a strange acceleration, and analyze the remaining 19 fasts, 28 moderates, and 5 slows. Our analyses for slow ICMEs show that $a_{ave} = -k_1(V - V_{bg})$ is more suited than $a_{ave} = -k_2(V - V_{bg})|V - V_{bg}|$ to describes their motion, where a_{ave} is the average acceleration, k_1 and k_2 are coefficients, and V is propagation speed of ICME. This result is the same as that for ICMEs having $V_{SOHO} - V_{bg} > 0 \text{ km s}^{-1}$ (*i.e.* a group of the fasts and moderates). In addition, we also found from examinations of all ICMEs above that the value of the coefficient k_1 has velocity dependence. On the basis of these, we propose a modified equation for the ICME motion.

Keywords: Coronal Mass Ejection, Interplanetary Coronal Mass Ejection, Interplanetary Scintillation

Structure of a termination shock: Parameter survey

MATSUKIYO, Shuichi^{1*}, Manfred Scholer²

¹ESST Kyushu Univ., ²Max-Planck-Institute

Structure of a termination shock is investigated by utilizing one-dimensional full particle-in-cell simulation. Parameter dependence of the shock structure on solar wind plasma beta, distribution function of the pickup ions, Alfvén Mach number, ion-to-electron mass ratio, and electron plasma to cyclotron frequency ratio is discussed, while a relative pickup ion density and shock angle are fixed to 30% and 87 deg., respectively. When the solar wind plasma beta is low ($\beta=0.17$), modified two-stream instability (MTSI) gets excited in the extended foot sustained by reflected pickup ions and both solar wind electrons and ions are heated. If the solar wind plasma beta gets five times higher ($\beta=0.85$), on the other hand, the MTSI is weakened and the pre-heating of the solar wind plasma in the extended foot is suppressed. When the distribution function of the upstream pickup ions are given by Maxwellian, instead of a spherical shell, the size of the extended foot becomes larger and heating of downstream solar wind ions is less efficient. If the Alfvén Mach number becomes high ($M=28$), a self-reformation of the shock front occurs. This results in a wiggled structure of the downstream solar wind ions, but the reformation seems not to contribute to strong acceleration of pickup ions.

Keywords: termination shock, pickup ion

Probing solar coronal fields using the Sun's shadow in cosmic ray intensity observed with the Tibet air shower array

M. Amenomori¹, C. Ding², K. Hibino³, N. Hotta⁴, T. Inaba⁵, A. Ishizak⁵, F. Kajino⁶, K. Kasahara⁷, Y. Katayose⁸, C. Kato⁵, K. Kawata², M. Kozai⁵, T. Masakawa⁵, K. Mizutani⁹, T. Motoyama⁸, MUNAKATA, Kazuoki^{5*}, Y. Nakano⁵, Y. Nakao², H. Nanjo¹, M. Nishizawa¹⁰, M. Ohnishi², I. Ohta¹¹, S. Ozawa⁷, T.Y. Saito¹², T. Saito¹³, M. Sakata⁶, T. Sako⁸, M. Shibata⁸, M. Shiomi¹⁴, T. Shirai³, S. Shishido⁸, H. Sugimoto¹⁵, M. Takita², N. Tateyama³, S. Torii⁷, H. Tsuchiya¹⁶, S. Udo³, Y. Yamamoto⁶, S. Yasue¹⁷, K. Yoshigoe², T. Yuda²

¹Department of Physics, Hirosaki U., ²ICRR, U. of Tokyo, ³Faculty of Engineering, Kanagawa U., ⁴Faculty of Education, Utsunomiya U., ⁵Department of Physics, Shinshu U., ⁶Department of Physics, Konan U., ⁷RISE, Waseda U., ⁸Faculty of Engineering, Yokohama Nat. U., ⁹Saitama U., ¹⁰National Institute of Informatics, ¹¹Sakushin Gakuin U., ¹²Max-Planck-Institut für Physik, ¹³Tokyo Metropolitan College of Industrial, ¹⁴College of Indust. Technology, Nihon U., ¹⁵Shonan Institute of Technology, ¹⁶RIKEN, ¹⁷School of General Education, Shinshu U.

Very high energy cosmic rays travel nearly straight in the interplanetary space between the Sun and the Earth. The Sun shields these particles and casts a tiny shadow in the cosmic ray intensity measured at the Earth, so-called the "Sun's shadow". We continuously observed the Sun's shadow in 10 TeV cosmic ray intensity with the Tibet air shower array over an entire period of the Solar Cycle 23. We find a good correlation between the intensity deficit in the Sun's shadow and the solar activity changing with the 11-year cycle. The intensity deficit decreases (increases) in the solar activity maximum (minimum) period. In this paper, we present a variation of the Sun's shadow observed in a period from 1996 through 2009 and discuss the effect of the large-scale structure of the coronal magnetic field on the shadow by means of numerical simulations. We calculate trajectories of antiparticles ejected from the Earth to the Sun in the model magnetic field and reproduce the Sun's shadow. For the magnetic field in the solar corona, we adopt the PFSS (Potential Field Source Surface) and CSSS (Current Sheet Source Surface) models and examine which model can reproduce better the observed Sun's shadow. The PFSS model ignores effects of the electric current in the solar corona, while the CSSS model takes account of the large-scale horizontal and volume currents. The large-scale magnetic field structures derived from two models are significantly different. We find that the intensity deficit in the simulated Sun's shadow is very sensitive to the coronal field structure. It is clear from the statistical consideration that the Sun's shadow observed by the Tibet air shower array is better reproduced by the CSSS model than by the PFSS model. The Tibet air shower experiment succeeded for the first time in evaluating the coronal field models by using the Sun's shadow observed in the very high energy cosmic ray intensity.

Keywords: Sun's shadow, solar coronal magnetic field, solar cycle variation, galactic cosmic rays, air shower

Spectrum characteristics of solar radio type-I burst by statistical analysis

IWAI, Kazumasa^{1*}, MISAWA, Hiroaki¹, TSUCHIYA, Fuminori¹, MORIOKA, Akira¹, MIYOSHI, Yoshizumi², MASUDA, Satoshi²

¹PPARC, Tohoku University, ²STEL, Nagoya University

Non-thermal electrons accelerated in the corona emit radio waves in the metric range. They are observed as several types of solar radio bursts. Type-I noise storm is one of the most frequently observed solar radio phenomena at a metric frequency range. They are thought to be a plasma emission around local plasma frequency. Type-I contains many complex fine structures in their spectra. They are thought to be caused by some inhomogeneities of particle acceleration, wave generation, radio emission, and/or radio propagation processes. However, the fundamental spectral structures have not been resolved sufficiently because of the limited time and frequency resolutions of the observation system. Now, more detailed analysis of spectrum structures of them are thought to be important to understand plasma processes in the solar corona. The purpose of this study is to extract fundamental spectrum parameters of type-I from high resolution observation and explain its generation processes.

Iitate Planetary Radio Telescope (IPRT) is a ground-based radio telescope developed by Tohoku University. Solar radio observation system of IPRT (AMATERAS) enables us to observe solar radio bursts in the frequency range between 150 and 500 MHz with the 10 ms accumulation time and 61 kHz bandwidth. It is suitable for observing characteristics of fine spectrum structures of solar radio bursts. The observational results for Type I bursts showed that the fundamental spectral structures have a duration of between 100 and 1000 ms. Typical full-width half-maximum of the burst bandwidth is between 1 and 5 MHz. Each element shows symmetric exponential growth and decay phases in time. We have also researched the peak flux distribution of type-I bursts. The observational results suggest that type-I bursts show a power-law distribution with the spectral index of 2 - 3. The index is larger than that of flares and radio type-III bursts (< 2). According to the logistic avalanche model (see Aschwanden et al 1998), the observed soft spectrum can be explained that source region of type-I is localized in a small region or the dominant instability of type-I grows slowly.

Keywords: Sun, corona, radio burst, ground-based observation, particle acceleration

Power spectra of the electron density fluctuation in the solar corona obtained by radio occultation observations

MIYAMOTO, mayu^{1*}, IMAMURA, Takeshi², ANDO, Hiroki¹, ISOBE, Hiroaki³, ASAI, Ayumi³, SHIOTA, Daikou⁴, YAJI, Kentaro⁵

¹Department of Earth Planetary Science, Graduate School of Science, The University of Tokyo, ²Japan Aerospace Exploration Agency, Institute of Space and Astronautical Science, ³Unit of Synergetic Studies for Space, Kyoto University, ⁴RIKEN, ⁵College of Science, Rikkyo University

The Venus orbiter Akatsuki is now in orbit around the Sun. Using Akatsuki, we have conducted radio occultation observations of the solar corona from June 6, 2011 to July 8, 2011. The method utilizes radio waves that are transmitted from the spacecraft and penetrate the solar corona when the spacecraft is occulted by the sun as seen from the Earth. Due to the movement of small-scale density irregularities across the ray path, the frequency (phase) and intensity of the signal received at the ground station change with time. By analyzing these time series we could get information on the power spectrum of the electron density fluctuation. The observations were conducted at solar offset distances of 1.5-20.5 R_s (R_s is solar radius), a region which is not well studied in the previous observations. Coordinated simultaneous observations using the space solar telescope Hinode were also conducted from June 24 to 25 when the ray path from Akatsuki was especially close to the sun.

In this paper we will present the radial variation of the power spectrum of the electron density fluctuation. The spectra show Kolmogorov-like logarithmic slopes at far distances, while knee features are seen at close distances.

Keywords: radio occultation observations, solar corona

Data assimilation of the solar wind in the inner heliosphere to estimate the source function and solar wind variation

TAO, Chihiro^{1*}, Iku Shinohara¹, SHIOTA, Daikou², KATAOKA, Ryuho³, MIYOSHI, Yoshizumi⁴, TOKUMARU, Munetoshi⁴

¹ISAS/JAXA, ²RIKEN, ³Tokyo Institute of Technology, ⁴STEL, Nagoya University

We develop a technique for predicting variations of the solar wind and source functions by incorporating wind velocity data from interplanetary scintillation (IPS) into a three dimensional magneto-hydrodynamic (MHD) solar wind model in the context of data assimilation using the Ensemble Kalman filter. In the data assimilation process, we constrain the solar wind source function which relates the observable magnetic field on the solar surface and terminal solar wind velocity. Previous studies estimated the function statistically, while we estimate the best fit model coefficients in this study.

We perform the "twin experiments" to evaluate the data assimilation method and containment of the source function and obtain results as follows : i) Variations of the solar wind and source function coefficients are well reproduced by the data assimilation. ii) Case for Ensemble number of being larger than 15 shows good estimation for 40 data per day case. iii) IPS data positions do not affect the prediction effectively because the source function affects large structure. The number of the state of a system is $21 \text{ (radial)} \times 360 \text{ (longitude and latitude)} \times 8 \text{ (MHD parameter)} + 2 \text{ (source function coefficients)} = 60,482$, while SOHO/MDI magnetic field data is referred at the inner boundary and IPS observation ~ 40 per day is assimilated. We discuss the applicability of this method to the observed solar wind.

Keywords: solar wind, data assimilation, simulation, MHD

Relationship between solar wind speed and coronal magnetic field parameter through solar cycle 23 and 23/24 minimum

FUJIKI, Ken'ichi^{1*}, Momotaro Noda², TOKUMARU, Munetoshi¹

¹Solar-Terrestrial Environment Laboratory, Nagoya University, ²Division of Particle and Astrophysical Science, Graduate School of Science, Nagoya University

Solar wind acceleration mechanism is one of the most important issues to be solved in solar wind physics. We have been studied the relationship between velocity of solar wind which comes from coronal holes and coronal magnetic field parameter. Precedence researches in our group showed that solar wind speed, V had been in proportion to a parameter Bp/f from solar minimum in cycle 22/23 to solar maximum in cycle 23, where Bp and f are magnetic field strength and expansion factor of the magnetic flux tube, respectively [1]. In solar minimum in the cycle 23/24, however, polar magnetic field gets weaker and solar wind has lower density, lower temperature, and lower mass flux than the previous minimum [2]. For this reason, we examined the relationship between V and Bp/f from 22/23 to 23/24 solar minima. In this analysis we used following data set. Solar wind velocity map in each Carrington rotation was derived from the interplanetary scintillation measurements at Solar-Terrestrial Environment Laboratory. Coronal magnetic field in each Carrington rotation was calculated by potential field source surface model using synoptic photospheric magnetic field data observed at Kitt Peak National Solar Observatory. As results, we found that V always correlates positively Bp/f over the solar cycle. Then we focused on the difference of slopes of the regression lines between two solar minima. The slope in the 23/24 minimum becomes larger, in other words, the fast solar wind in the 23/24 minimum has smaller Bp/f compared to that in the 22/23 minimum. This result was compared with a theoretical model of solar wind acceleration [3] by taking account of the declining of solar wind parameters. We confirmed that the difference of regression lines in two solar minima is consistent with global trend of solar and solar wind variations.

Reference

- [1] Fujiki et al., Adv. Space Res., Volume 35, Issue 12, 2185, (2005)
- [2] McComas et al., Geophys. Res. Lett., 35, 18103, (2008)
- [3] Suzuki, T. K., Astrophys. J. Lett., 640,75, (2006)

Keywords: solar wind, interplanetary scintillation, solar cycle

Long term variation of the solar diurnal anisotropy of galactic cosmic rays over four solar activity cycles

MUNAKATA, Kazuoki^{1*}

¹Physics Dept., Shinshu U.

The galactic cosmic ray (GCR) anisotropy observed with the muon detector network monitoring high-energy GCR intensity provides us with unique information of physical parameters, such as the spatial density gradient and the scattering mean free path of GCRs, which reflect the large-scale magnetic structure governing the GCR propagation in the heliosphere. The solar cycle variation of the anisotropy particularly gives important information on the temporal variation of the GCR propagation in the heliosphere. In this paper, we analyze the solar diurnal anisotropy observed with a network of surface and underground muon detectors monitoring the primary GCRs in a wide energy range of 50-500 GeV. This network includes a detector at Nagoya which has been in operation more than 40 years. The derived anisotropy shows clear 11-year and 22-year variations respectively in clear correlations with the solar activity- and magnetic-cycles. We will discuss the physical mechanisms responsible for these long-term variations.

Keywords: galactic cosmic rays, solar diurnal anisotropy, muon detector networks

Long-term variation of the solar wind acceleration

HAKAMADA, Kazuyuki^{1*}, TOKUMARU, Munetoshi², FUJIKI, Ken'ichi²

¹Chubu University, ²Solar-Terrestrial Environment Laboratory

We calculate the three-dimensional structure of the coronal magnetic field during the time interval of 1800 Carrington rotation (CR 1800) through CR 2075 by the Radial-Field model devised by Hakamada with synoptic maps of Kitt Peak photospheric magnetic field data. This time interval covers almost two solar activity cycles of about 20 years. We constructed synoptic maps of $\text{Log}_{10}|\text{Br}_{\text{sou}}|$, the common logarithm of the radial component of the coronal magnetic field on the source surface of 2.5 solar radii. We also constructed synoptic maps of $\text{Log}_{10}|\text{Br}_{\text{pho}}|$, the common logarithm of the radial component of the photospheric magnetic field projected on the source surface along the coronal magnetic field line. We, further, constructed synoptic maps of the solar wind speed, V , estimated by the CAT method with results of IPS observations. Those three values can be compared directly with each other because of the values on the synoptic maps of the same format. We draw the distribution of V in xyz space; x-axis corresponds to $\text{Log}_{10}|\text{Br}_{\text{sou}}|$, y-axis corresponds to $\text{Log}_{10}|\text{Br}_{\text{pho}}|$, and Z-axis corresponds to V . It is found that the distribution of V has a planar structure in xyz space. We, thus, assume the empirical equation of $V = a + b * \text{Log}_{10}|\text{Br}_{\text{sou}}| + c * \text{Log}_{10}|\text{Br}_{\text{pho}}|$, and calculate the multiple correlation coefficient, r , and the multiple regression coefficients, a , b , and c . It is found that, although the r has smaller value, around 0.3, during the solar minimum phase, r is very high, around 0.7, during the solar maximum phase. These results suggest that the solar wind is accelerated not only by the mechanism related to the coronal magnetic field and photospheric magnetic field but also by an unknown mechanism which shows the long-term variations.

Keywords: solar wind, acceleration, long-term variation

Determination of 3D configuration of magnetic clouds using 2D imaging data of solar wind

MARUBASHI, Katsuhide^{1*}, TOKUMARU Munetoshi², JACKSON Bernard V.³

¹none, ²STEL, Nagoya University, ³CASS, University of California, San Diego

The structure of magnetic cloud (MC) has long been studied by fitting in-situ magnetic field measurements to magnetic flux rope models. Such studies generally provide the size and orientation of the model structure, which are taken to be applicable only locally to the portion where the cloud passed the spacecraft. The obtained geometry often changes depending on the models used for the fitting. For example, for a single observational data set, a cylinder model and a torus model often give different cloud axis orientations. Thus we need further consideration about the global configuration to deduce the 3D structure of the MC. 2D images from heliospheric remote sensing measurements provide reliable constraints about the global MC structure. With the above in mind we attempt to study the global structure of MC by combining the model fitting results and the 3D reconstruction data from the Solar Mass Ejection Imager (SMEI) and interplanetary scintillations (IPS). For this purpose, we first select MC events in which the proton densities are high enough (generally > 20 /cc) in the sheath regions behind the driven shocks and/or in the regions occupied by MCs. Then we examine possible 3D configurations which are consistent with the orientation of the MC axis obtained from model fittings. The global MC structure is finally obtained by applying the constraints from 2D image data. Our preliminary examination shows that the above procedure is helpful for determining the most probable 3D global structures of MCs.

Keywords: solar wind magnetic field, magnetic flux rope, solar wind density, model fitting, magnetic cloud

Source estimation of Electrostatic Solitary Waves (ESWs) observed by Kaguya near the Moon

HASHIMOTO, Kozo^{1*}, OMURA, Yoshiharu², KASAHARA, Yoshiya³, KOJIMA, Hirotsugu², HASHITANI, Maki⁴, ONO, Takayuki⁵, TSUNAKAWA, Hideo⁶

¹Paleological Association of Japan, ²RISH, Kyoto University, ³Kanazawa University, ⁴Kyushu Electric Power Co., ⁵Tohoku University, ⁶Tokyo Institute of Technology

In KAGUYA (SELENE) LRS[1], WFC-L [2] observes waveforms of plasma waves in 100Hz-100kHz and a lot of electrostatic solitary waves (ESWs) have been observed. Some results have been reported [3].

An ESW potential generally has a two-dimensional structure. They have not only a parallel component to the background magnetic field, but also a perpendicular component. Then received ESW fields are distorted from well-known bipolar fields based on the one-dimensional potential. In order to evaluate the effects of the perpendicular components, the received ESW waveforms are fitted to ideal ESW waveforms based on the two-dimensional structures. We examined where ESWs are received near the moon. The source positions of the ESWs will be discussed based on the two-dimensional structures and the background magnetic field directions and magnetic anomalies.

References

[1] Takayuki Ono, Atsushi Kumamoto, Yasushi Yamaguchi, Atsushi Yamaji, Takao Kobayashi, Yoshiya Kasahara, and Hiroshi Oya, Instrumentation and observation target of the Lunar Radar Sounder (LRS) experiment on-board the SELENE spacecraft, *Earth Planets Space*, 60, 321-332, 2008.

[2] Y. Kasahara, Y. Goto, K. Hashimoto, T. Imachi, A. Kumamoto, T. Ono, and H. Matsumoto, Plasma Wave Observation Using Waveform Capture in the Lunar Radar Sounder on board the SELENE Spacecraft, *Earth, Planets and Space*, 60, 341-351, 2008.

[3] K. Hashimoto, M. Hashitani, Y. Kasahara, Y. Omura, M.N. Nishino, Y. Saito, S. Yokota, T. Ono, H. Tsunakawa, H. Shibuya, M. Matsushima, H. Shimizu, and F. Takahashi, Electrostatic solitary waves associated with magnetic anomalies and wake boundary of the Moon observed by KAGUYA, accepted for publication in *Geophys. Res. Lett.*, 37, L19204, doi:10.1029/2010GL044529, 2010.

Keywords: electrostatic solitary wave, moon, Kaguya

Proton entry into the plasma void formed downstream of an insulating, non-magnetized obstacle in the solar wind

NAKAGAWA, Tomoko^{1*}

¹Tohoku Institute of Technology

The proton entry into the plasma void formed downstream of an insulating, non-magnetized obstacle in the supersonic flow of the solar wind is studied by using a two-dimensional, electromagnetic particle-in-cell simulation. The protons are accelerated by the negative electric potential in the plasma wake, and the size of the proton void is much smaller than that estimated from the ratio of the solar wind bulk speed to the proton thermal speed. In the slow solar wind, the proton void shrinks because the protons are well accelerated within a short distance past the obstacle, due to the long time required to pass through the intense electric field near the wake boundary.

Keywords: moon, near moon wake, proton acceleration, solar wind, PIC simulation, electric potential