

Unravelling metamorphic-fluid events in Gondwana collision: U-Pb-REE constraints from Sor Rondane Mountains, Antarctica

HOKADA, Tomokazu^{1*}, HORIE, Kenji¹, ADACHI, Tatsuro², OSANAI, Yasuhito², NAKANO, Nobuhiko², BABA, Sotaro³, TOYOSHIMA, Tsuyoshi⁴

¹National Institute of Polar Research, ²Kyushu University, ³University of Ryukyus, ⁴Niigata University

Assembly of Gondwana supercontinent has been argued in numerous studies. Generally the reported ages of Gondwana collision zones are in the range of 750-500 Ma, and two main age groups of 750-620 Ma and 570-530 Ma are estimated (e.g., Meert, 2003). The first episode (~620Ma) is mainly reported from eastern Africa to East Antarctica, and the second episode (~530Ma) is dominated in southern Africa through East Antarctica-Sri Lanka-southern India to eastern Australia. It is, however, not yet fully understood the superimposition of these two events in the crossing region.

Sor Rondane Mountains in East Antarctica is one such area where 640-630 Ma and 550-520 Ma metamorphic-fluid events are recorded (e.g., Shiraishi et al., 2008. Adachi, 2010). Greenschist-facies through amphibolite-facies to granulite-facies metamorphic rocks occupy the area, and the granulite-facies rocks are dominated in the northeastern-central part of the area (e.g., Shiraishi et al., 1992; Osanai et al., 1992). Multiple leucocratic veins and granitic intrusives are also developed. We present zircon and monazite U-Th-Pb and REE analyses by using ion microprobe and electron microprobe applying to garnet-biotite-sillimanite gneiss and associated leucoveins in the central part of Sor Rondane Mountains. Zircon in garnet-sillimanite-biotite gneiss yields c.640-630 Ma with minor >700 Ma and 550-520 Ma ages, and gives clear age-chemistry relation that HREE/MREE ratios drop in c.640-630 Ma zircon crystals compared with older and younger grains that are presumably controlled by REE partitioning with the coexisting garnet. Zircon and monazite in multiple generations of leucoveins also yield >700 Ma, 640-630 Ma and 550-520 Ma ages. Combined rare earth elements (REE) chemistry with the U-Th-Pb age domains in syn- and post-metamorphic leucoveins suggests contrasting isotopic and chemical signatures, and could provide constraints for decoding Neoproterozoic-Cambrian metamorphic-fluid regimes in the Gondwana collision zone.

Keywords: Gondwana, East Antarctica, Sor Rondane Mountains, metamorphism, zircon, monazite

Evolution of continental lithosphere in the Sor Rondane Mountains, East Antarctica

OWADA, Masaaki^{1*}, KAMEI, Atsushi², HORIE, Kenji³, BABA, Sotaro⁴, OSANAI, Yasuhito⁵

¹Yamaguchi University, ²Shimane University, ³National Institute of Polar Research, ⁴University of Ryukyus, ⁵Kyushu University

The Sor Rondane Mountains is situated within the collision zone between the West and East Gondwana and the time of collision is regarded as the late Proterozoic (650-600 Ma). The mountains consist of greenschist- to granulite-facies metamorphic rocks and various kinds of intrusive rocks. The tonalite complex is exposed in the southern part of the mountains and its magmatic age is considered to be at the middle Proterozoic (990-920 Ma). This tonalite would originally be formed at the subduction related tectonic setting prior to the collision event. Large amounts of microgabbro occur as mafic magmatic enclaves (MMEs; 990 Ma) and dikes (950 Ma) in the tonalite complex. Unmetamorphosed lamprophyre dikes intrude the tonalite complex and gneisses during the late- to post-collisional stages. The intrusive age of the lamprophyre is of 560 Ma. The magma processes of the tonalite complex together with the late- to post collisional lamprophyre dikes, therefore, provide us useful information of the evolution of continental lithosphere during the formation of Gondwana supercontinent.

The microgabbro represents the low-K and tholeiitic series, and is geochemically classified into Low-Ti and High-Ti microgabbros. The MMEs and dikes of microgabbros are equivalent to the Low-Ti and the High-Ti microgabbros, respectively. The Low-Ti and High-Ti microgabbros show geochemical signature similar to the Oceanic Arc Basalts and the Back-Arc Basin Basalts, respectively. The middle Proterozoic magma processes would, therefore, proceed at a subduction zone with back arc spreading in an oceanic arc environment. The lamprophyre corresponds to alkaline rocks in the TAS diagram, and are characterized by high abundances of LILE elements and REE, especially Rb, Ba, Sr and LREE. The trace element abundances normalized to primitive mantle display enrichment of LILE and depression of HFSE with Nb and Ta negative anomalies. The lamprophyre is plotted in the within-plate field and a part of the island arc field that is close to the within-plate field on some discrimination diagrams. Considering the geochemical features, the lamprophyre was formed in a within-plate tectonic setting by the mixing of subduction-related materials. The initial Sr isotopic ratios (SrI) range from 0.7022 to 0.7040 ($\epsilon_{\text{SrI}} = -14$ to 12) for the Low-Ti microgabbro and from 0.7024 to 0.7030 ($\epsilon_{\text{SrI}} = -14$ to 1) for the High-Ti microgabbro. The initial epsilon Nd values for the Low- and High-Ti microgabbros are calculated within the same range ($\epsilon_{\text{NdI}} = -0.1$ to +0.5). On the other hand, the isotopic compositions of the lamprophyre show SrI = 0.7043 - 0.7044 ($\epsilon_{\text{SrI}} = +7.6$ to +9.2) and $\epsilon_{\text{NdI}} = -0.62$ to -0.34.

The geochemical studies including Sr-Nd compositions reveal that the microgabbros have been originated from a depleted source, whereas chemical compositions of the lamprophyre is more enrichment rather than those of the microgabbros. Consequently, the magma processes in the Sor Rondane Mountains reflect the evolution of lithosphere from the middle Proterozoic to the early Paleozoic; the depleted mantle at the initial subduction stage then changing to the enriched mantle at the continental collision stage. This lithospheric evolution can be explained by interaction between the depleted mantle and the enriched materials (e.g., slab-derived fluids, melting product of subducted crustal rocks, or reaction with fossil wedge mantle) during closure of the Mozambique Ocean.

Keywords: Gondwana supercontinent, Antarctica, Sor Rondane Mountains, tonalite, microgabbro, lamprophyre

Occurrence of alkali ~ highly potassic dykes intruded into metamorphic rocks on Lutzow-Holm Complex, East Antarctica

MIYAMOTO, Tomoharu^{1*}, TSUNOGAE, Toshiaki², DUNKLEY, Daniel Joseph³

¹Kyushu University, ²University of Tsukuba, ³Geoscience Australia

The Lutzow-Holm Complex (LHC) of Dronning Maud Land, East Antarctica, is a high grade metamorphic terrane within the East Antarctic Shield, situated to the west of Rayner Complex and to the east of the Yamato-Belgica Complex. Japanese Antarctic Research Expeditions (JARE) have carried out detailed surveys of its geology and tectonics for many coastal exposures, and have identified various kinds of metamorphic rocks. Subsequent igneous rocks as granites and pegmatites that intruded during and after the peak metamorphism, were also recognized.

Mafic dyke rocks, which discordantly intruded the surrounding gneisses, were already found in some localities. A few post-metamorphic mafic - intermediate rock dykes were newly found in Skallevikshalsen, Rundvagshetta, and Niban-Iwa as well as Innhovde and Austhovde in the LHC, during the geological survey by JARE-52. The dykes on Skallevikshalsen and Rundvagshetta were thin sheets with a few centimeters to half meter in thickness, and almost NS (to slightly NNE-SSW) trending with dipping east steeply. On the other hand, dyke from Niban-iwa on the Prince Orav Coast, has a ten to twenty centimeter thickness and dip to NE with WNW-ESE striking. Internal textures as mineral arranging and extension are parallel to the trend of the dyke intrusion. Dykes in Rundvagshetta were strongly related for the origin with post-genetic pegmatites, and partly modified to amphibolite with coarse hornblende by the pegmatitic activity.

The dyke rocks are holocrystalline and aphyric, and grain size is mostly between 0.1 and 2 mm. They consist dominantly of alkali-feldspar and subsequent biotite, augite, hornblende, titanite, apatite and minor amount of plagioclase and quartz: these mineral abundances vary according to their occurrences, and alkali-feldspar, apatite and quartz are commonly included in the dykes by all means more or less, and others are occasionally absence in some rock specimens. Minerals, especially biotite flakes, are commonly aligned to parallel to boundary between the dykes and the host gneisses.

Whole rock composition of dyke rocks are different in the 5 outcrops each; the SiO₂ and MgO contents range from 46.3 to 60.2 wt.% and from 9.48 to 0.69 wt.%, respectively. Their K₂O content reaches 3.42 - 10.83 wt.% with higher K₂O/Al₂O₃ and K₂O/Na₂O values than general igneous rocks. They are classified into tephrite, trachyandesite, and trachyte according to their total alkali versus silica characters. In them, dyke rocks in Skallevikshalsen resemble lamproite for their ultrapotassic characters (K₂O = 8.10 - 8.72 wt.%) with much MgO (= 7.92 - 9.48 wt.%) and abundance of typical minor elements (e.g., Ba, Sr).

In Rundvagshetta, host metamorphic rocks were partly metasomatized by hydrated reaction around the dykes, and garnet was broken down to biotite in the metasomatized domain: this hydration modification was stronger at close to the boundary, and garnet which far from the boundary was commonly survived. Since the dyke contains abundant biotite and subsequent apatite as well as alkali feldspar, it is expected that the dyke supplied fluid for the metasomatism during its activity.

Keywords: alkali rock, ultrapotassic rock, dyke

The stability of sapphirine + quartz in high/low oxygen fugacity rocks: a case study of Southern India/East Antarctica

SHIMIZU, Hisako^{1*}, TSUNOGAE, Toshiaki¹, M. Santosh²

¹Earth Evolution Sciences, Graduate School of Life and Environmental Sciences, University of Tsukuba, ²Faculty of Science, Kochi University

Sapphirine has been the focus of many petrological investigations for the last two decades as the mineral often occurs in Mg-Al rich and pelitic rocks formed at high temperature to ultrahigh temperature (UHT). Particularly, sapphirine coexisting with quartz is considered as one of the most diagnostic mineral assemblages of UHT metamorphism. It is also known that sapphirine often occurs in magnetite-bearing high oxygen fugacity rocks, and, in such cases, the mineral can incorporate considerable quantity of ferric iron as well as Fe²⁺. It is therefore important to evaluate the effect of Fe³⁺ content on the stability of sapphirine-bearing assemblages for estimating peak conditions as well as constructing *P-T* paths. In this study, we compared the stability of sapphirine + quartz in magnetite-bearing high-oxygen fugacity rocks from India (Madurai Block in the southern granulite terrane) with that in magnetite-absent low-oxygen fugacity rocks from Antarctica (Bunt Island in the Napier Complex) using mineral equilibrium modeling technique. The calculations have been done in NCKFMASHTO system using THERMOCALC 3.33 with an updated version of the internally consistent data set.

The Madurai Block is the largest granulite block in the Southern Granulite Terrane, India, which was formed by collisional orogeny related to the assembly of the Gondwana Supercontinent. The block contains granulites with various UHT mineral assemblages including sapphirine + quartz, orthopyroxene + sillimanite + quartz, and Al-rich orthopyroxene. Magnetite-bearing quartzo-feldspathic garnet-sillimanite granulites from Rajapalayam area in the southern part of the block, for example, contain sapphirine + quartz inclusion in garnet as a stable mineral assemblage at the peak of metamorphism. The calculated *T-X* pseudosections suggest that the stability temperature of sapphirine + quartz is lowered from 1000°C at reduced condition ($X_{Fe_2O_3} = 0.02$) to 910°C at oxidized condition ($X_{Fe_2O_3} = 1.0$).

The Napier Complex of Enderby Land, East Antarctica, underwent regional UHT metamorphism at ca. 2.5 Ga. Bunt Island in the Napier Complex, located in the highest-grade region of the complex, contains various kinds of UHT granulites including sapphirine-bearing rocks. Sapphirine + quartz assemblage, probably formed at the peak UHT condition, occur in sapphirine- and osumilite-bearing layers of the granulite. The absence of magnetite in the rocks indicates the sapphirine granulite was formed at reduced condition. *T-X* pseudosection of the rock suggests that the stability field of sapphirine + quartz is $T > 1050^\circ\text{C}$ at $X_{Fe_2O_3} = 0.04$, while it will be lowered in more oxidized condition ($T > 800^\circ\text{C}$ at $X_{Fe_2O_3} = 0.24$).

The results of this study demonstrated that the occurrence of sapphirine + quartz in UHT rocks is highly controlled by the oxidation state of the host rocks, particularly low oxygen fugacity rock is especially strongly influenced. In the case of Bunt Island, if $X_{Fe_2O_3}$ increases by 0.04, the stability temperature of sapphirine + quartz will be lowered by 50°C. It is therefore important to evaluate the effect of Fe³⁺ content of the stability of sapphirine-bearing granulites for estimating peak conditions as well as constructing *P-T* paths even if granulites were formed at reduced condition.

Keywords: sapphirine + quartz, Southern India, East Antarctica, pseudosection, THERMOCALC, oxygen fugacity

Petrology of garnet-clinopyroxene rocks from the Gondwana suture zone in southern India

TSUNOGAE, Toshiaki^{1*}

¹Univ. Tsukuba

The Palghat-Cauvery Suture Zone (PCSZ) in the southern granulite terrane, India, which separates Pan-African granulite blocks (e.g., Madurai and Trivandrum Blocks) to the south and Archean terrane (e.g., Salem Block and Dharwar Craton) to the north is regarded as a major suture zone in the Gondwana collisional orogeny. It probably continues westwards to the Betsimisaraka suture in Madagascar, and eastwards into Sri Lanka and possibly into Antarctica. The available geochronological data including U-Pb zircon and EPMA monazite ages indicate that the rocks along the PCSZ underwent an episode of high-grade metamorphism at ca. 530 Ma that broadly coincides with the time of final assembly of the Gondwana supercontinent. Recent investigations on high-grade metamorphic rocks in this region have identified several new occurrences of garnet-clinopyroxene rocks and associated meta-gabbros from Perundurai, Paramati, Aniyapuram, Vadugappatti, and Mahadevi areas in Namakkal region within the central domain of the PCSZ. They occur as elongated boudins of 1 m to 1 km in length within hornblende-biotite orthogneiss. The garnet-clinopyroxene mafic granulites contain coarse-grained (up to several cm) garnet (Alm30-50 Pyr30-40 Grs10-20) and clinopyroxene (XMg = 0.70-0.85) with minor pargasite, plagioclase (An30-40), orthopyroxene (hypersthene), and rutile. Garnet and clinopyroxene are both subidioblastic and contain few inclusions of clinopyroxene (in garnet) and plagioclase. Orthopyroxene occur only as Opx + Pl symplectite between garnet and clinopyroxene in almost all the localities, suggesting the progress of decompressional reaction: $\text{Grt} + \text{Cpx} + \text{Qtz} \Rightarrow \text{Opx} + \text{Pl}$, which is a dominant texture in the PCSZ. The prograde mineral assemblage of the rocks is therefore inferred to be $\text{Grt} + \text{Cpx} + \text{Qtz}$, although quartz was probably totally consumed by the progress of the reaction. The metamorphic P-T calculations using Grt-Cpx-Pl-Qtz geothermobarometers yield $T = 850\text{-}900\text{C}$ and $P > 13$ kbar, which is consistent with the occurrence of high-pressure Mg-rich staurolite in Mg-Al-rich rocks from this region. Fluid inclusion study of some garnet-clinopyroxene rock samples identified CO₂-rich fluid inclusions trapped as primary phases within garnet, suggesting that prograde high-pressure metamorphism was dominated by CO₂-rich fluids. The results therefore confirmed that the PCSZ underwent regional dry high-pressure metamorphism followed by the peak ultrahigh-temperature event probably associated with the continent-continent collisional and suturing history along the PCSZ.

Mineral equilibrium modeling of incipient charnockite and adjacent garnet-biotite gneiss from southern India

ENDO, Takahiro^{1*}, TSUNOGAE, Toshiaki¹, M. Santosh²

¹Univ. Tsukuba, ²Kochi Univ.

The Southern Granulite Terrane (SGT) in India is known for its classic exposures of regionally metamorphosed granulite-facies rocks formed during the collisional orogeny related to the amalgamation of Gondwana supercontinent. The SGT is composed of a collage of Proterozoic crustal blocks dissected by large Late Neoproterozoic shear/suture zones. The Trivandrum Granulite Block (TGB) is comprised dominantly of metasedimentary sequence with khondalites, leptynites and charnockites with subordinate quartzite, mafic granulite, calc-silicate rocks, and meta-ultramafic rocks. The TGB is known as one of the classic examples for the spectacular development of "incipient charnockites" within orthopyroxene-free felsic gneisses as exposed in several quarry sections in the states of Kerala and Tamil Nadu. The charnockite-forming process in the TGB is considered to have been triggered by the infiltration of CO₂-rich anhydrous fluids along structural pathways within upper amphibolite facies gneisses, resulting in the lowering of water activity and stabilization of orthopyroxene through the breakdown of biotite. However, no quantitative study on the stability of charnockitic mineral assemblage using mineral equilibrium modeling approach has been done so far. In this study, we report a new occurrence of incipient charnockite from Mavadi in the TGB and discuss the petrogenesis of granulite formation in an arrested stage on the basis of petrography, geothermobarometry, and mineral equilibrium modeling. In Mavadi, patches and lenses of charnockite (Kfs + Qtz + Pl + Bt + Grt + Opx + Ilm + Mag) of about 30 to 120 cm in length occur within Opx-free Grt-Bt gneiss (Kfs + Qtz + Pl + Bt + Grt + Ilm) host rocks. The application of mineral equilibrium modeling on charnockite assemblage in NCKFMASHTO system to constrain the conditions of charnockitization defines a P-T range of 800C at 4.5 kbar to 850C at 8.5 kbar, which is broadly consistent with the results from the conventional geothermobarometry (810-880C at 7.7-8.0 kbar) on these rocks. The P-T conditions are lower than the inferred peak metamorphic conditions from the ultrahigh-temperature granulites of the study area (T >900C), which might suggest heterogeneity in peak P-T conditions within this crustal block in relation to local buffering of metamorphic temperature by Opx-Bt-Kfs-Qtz assemblage. The result of T versus mole H₂O (M(H₂O)) modeling demonstrated that Opx-free assemblage in Grt-Bt gneiss is stable at M(H₂O) = 0.3 to 1.5 mol.%, and orthopyroxene occurs as a stable mineral at M(H₂O) <0.3 mol.%, which is consistent with the petrogenetic model of incipient charnockite related to the lowering of water activity and stabilization of orthopyroxene through breakdown of biotite by dehydration caused by the infiltration of CO₂-rich fluid from external sources. We also propose a possible alternative process to form charnockite from Grt-Bt gneiss through slight variations in bulk-rock chemistry (particularly K- and Fe-rich portion of Grt-Bt gneiss) that can enhance the stability of orthopyroxene rather than that of biotite.


Metamorphic reaction to describe local-scale difference in mineral assemblage stable under different conditions

YAMASAKI, Yukiko^{1*}, IKEDA, Takeshi¹, MOTOYOSHI, Yoichi², HIROI, Yoshikuni³, PRAME, Bernard N.⁴


¹Earth and Planet. Sci., Kyushu Univ., ²National Institute of Polar Research, ³Graduate School of Sciences, Chiba Univ., ⁴Geological Survey of Sri Lanka

Mineral assemblage of metamorphic rocks varies according to temperature, pressure and fluid composition. In regional metamorphic belts, we have a common understanding that the difference of mineral assemblage stems from the difference of temperature and pressure. In contrast, it is not so uncommon that the mineral assemblage varies in local scale (several decimeters). It is unlikely that temperature and pressure differ in these scales. Fluid composition, instead, could vary in local scale. Local variation of fluid composition could explain the change of the mineral assemblages that are mutually described by any dehydration reactions. Few researches have estimated the relevant reaction based on the difference of mineral composition and occurrence between the adjacent rocks. This study clarified the reactions to produce orthopyroxene in arrested charnockites from central Sri Lanka based on compositional difference of hornblende and biotite between charnockites and surrounding gneisses.



Biotite in charnockites is enriched in Fe+Mg and depleted in Ti as compared with that in the host gneisses. This compositional difference is expressed as the following substitution with a site vacancy, [],



The charnockites contain hornblende that is rich in K, Fe+Mg and Si, and poor in Na, Ti and Al relative to those in the surrounding gneisses, which is described by the substitution,



Using these substitutions, the reactions responsible for the formation of orthopyroxene can be estimated as,



Each reaction contains either biotite or hornblende. Changes of the modal abundance of biotite and hornblende reflect the amount of orthopyroxene produced by reaction (1) and (2), respectively. The amounts of orthopyroxene in the estimated reactions well agree with the modal abundance of orthopyroxene in charnockites. These reactions are dehydration reactions, indicating that the mineral assemblage observed in the charnockites is stable under conditions of CO₂-rich fluid. We conclude that the difference of mineral assemblage between charnockites and gneisses can be ascribed to the difference of fluid composition.

Keywords: charnockite, hornblende-biotite gneiss, Sri Lanka, metamorphic reaction

Mid to late Archean TTG magmatism in the eastern Madagascar; a view from whole rock geochemistry and U-Pb geochronology

ICHIKI, Takashi^{1*}, Yasuhito Osanai², Masahiro Ishikawa¹, Nobuhiko Nakano², Tatsuro Adachi²

¹Graduate School of Environment and Information Sciences, Yokohama Nat. Univ., ²Earth Sci., Kyushu Univ.

Madagascar occupies a key position in the East African Orogen for understanding the continental growth and the tectonics of collision between East Gondwanaland and West Gondwanaland. Especially in the eastern Madagascar is composed of mid Archean domain (Masora) in the east and late Archean domain (Antananarivo) in the west.

The magma genesis and timing of magmatism were studied by whole-rock chemical analysis and LA-ICP-MS U-Pb zircon age dating of granitoids in these domains.

Masora domain is divided into two parts, north and south region, defined by metamorphic grade. The rocks from the north region are only weakly deformed. The north region mainly consists of trondhjemites with subordinate amounts of metapelites including meta-BIFs, and late granitoids with mafic-ultramafic rocks. Trondhjemites in the northern Masora domain are characterized by high SiO₂ (67.80-70.98 wt.%), high Al₂O₃ (15.86-18.44 wt.%), and high Na₂O (5.35-5.98 wt.%), low TiO₂ (0.27-0.40 wt.%), Mg# (31-35), CaO (1.90-2.24 wt.%), K₂O (1.64-2.65 wt.%). Antananarivo domain is divided into two parts, north and south region, defined by lithology. The south region mainly consists of Hbl-Bt gneisses with subordinate amounts of Grt-Opx granulites, amphibolites, quartzites and metapsammites including meta-BIFs. Whole rock chemical analyses for the major and trace elements demonstrate that Hbl-Bt gneisses in the southern Antananarivo domain are of igneous origin and chemically comparable with CIPW normative tonalities. Hbl-Bt gneisses are characterized by high SiO₂ (71.00-73.16 wt.%), high Al₂O₃ (15.89-16.33 wt.%), and high Na₂O (4.41-4.67 wt.%), low TiO₂ (0.18-0.23 wt.%), Mg# (38-43), CaO (3.40-3.82 wt.%), low K₂O (1.04-1.71 wt.%). All of these granitoids (trondhjemites and Hbl-Bt gneisses) show pronounced negative Nb, Ti, P anomalies on the primitive mantle-normalized spidergram. These characteristics are comparable to Archean TTG (tonalite-trondhjemite-granodiorite) (e.g. Martin et al., 2005).

A xenocrystic zircon in a trondhjemite sample collected from the northern part of the Masora domain gives a single grain concordant age of mid Archean (ca. 3.2 Ga). Hbl-Bt gneiss sample in the southern part of the Antananarivo domain shows slightly scattered and discordant late Archean age (ca. 2.7 Ga). This new age is slightly older than reported oldest ages of Antananarivo domain (ca. 2.5 Ga; Kroner et al., 2000).

These results show that the area between mid Archean (Masora) and late Archean (Antananarivo) domains is underlain by ca. 2.7Ga tonalitic rocks. Similar magmatic age is reported from the southern India. Late Archean magmatic age of the charnockite and meta-granite (ca. 2.65-2.53) were reported from the Salem Block in the southern India (Clark et al., 2009; Sato et al., 2011), where located between the mid to late Archean Dharwar Craton (e.g. Peucat et al., 1993) to the north and late Archean Madurai Block (Plavsa et al., 2012) to the south. Hence we speculate that Madagascar and India records progressive outward continental growth by accretion of mid-to-late Archean (ca. 2.7-2.6 Ga) crust and the late Archean (ca. 2.5 Ga) crust to ca. 3.2Ga Dharwar nuclei crust. Although Tucker et al. (2011) suggested the 'Greater Dharwar Craton' model to explain the juxtaposition of Madagascar with India from the late Archean (ca. 2.5 Ga), we suggest more stepwise crustal growth in the Archean era.

Keywords: Madagascar, Archean TTG, geochronology, geochemistry