

Significance of melt inclusions in Zrn in estimating the duration of high-T metamorphism -An example of the Ryoke belt

KAWAKAMI, Tetsuo^{1*}, YAMAGUCHI, Isao¹, MIYAKE, Akira¹, MAKI, Kenshi¹, SHIBATA, Tomoyuki², Takaomi D. Yokoyama¹, HIRATA, Takafumi¹

¹Graduate School of Science, Kyoto Univ., ²Geothermal Research Lab, Kyoto Univ.

It is important to understand when and how Zrn was formed in order to interpret the U-Pb Zrn ages. Melt inclusion in Zrn is an important evidence of Zrn growth in the presence of melt. Therefore, they contribute to constrain the timing of Zrn growth in high-T metamorphic rocks. In this contribution, we report the occurrence of melt inclusions in Zrn from the migmatite of the Ryoke metamorphic belt at Aoyama area, SW Japan.

In the Aoyama area, the pelitic-psammitic schists dominate in the north, and migmatites (metatexites and diatexites) dominate in the south [1]. Zircon (Zrn) in pelitic-psammitic schists from the low-T part of the Grt-Crd zone is coarse-grained and shows almost no metamorphic overgrowth. On the other hand, Zrn in migmatites from the mid- and high-temperature part of the Grt-Crd zone has a thin, bright layer under BSE image along which tiny inclusions of several microns are aligned [cf. 2]. TEM observation of some of these tiny inclusions gave halo patterns revealing that they are the glass (formerly melt) rich in Si, Al and K. This melt inclusion alignment divides the core with various detrital ages from the rim that gives U-Pb concordia age of 93.1 +/- 2.9 Ma. The rim of the Zrn from the high-T part of the Grt-Crd zone is over 20 um in thickness, and its Th/U ratio is less than 0.02. Presence of melt inclusion alignment at the core/rim boundary shows that the melt was present when the Zrn rim grew. Therefore, the Zrn rim probably grew under the presence of melt and monazite during the Ryoke metamorphism. The melting reaction that consumes Bt + Sil + Qtz and produces Grt + Crd + Kfs +/- Ilm + melt plays an important role in the formation of melt in this area [1]. However, Bt is not an important sink of zirconium [3], and Bt breakdown cannot supply zirconium enough for new Zrn rim growth. In this case, dissolution of pre-existing Zrn is required. It would be difficult to saturate melt in terms of Zrn component by dissolving Zrn when the amount of melt is increasing. Therefore, it is likely that 93.1 +/- 2.9 Ma Zrn rim crystallized during the solidification of the melt in migmatites, possibly near the wet-solidus. Mixed analysis of U-Pb dating of Zrn core and rim reveals that similar young rim is developed, although thin, in Zrn from the mid-T part of the Grt-Crd zone. These results imply that presence of partial melts plays an important role in the dissolution and recrystallization of Zrn [e.g., 4].

Although the whole-rock zirconium content is not especially high in the pelitic-psammitic schists from the low-T part of the Grt-Crd zone, modal amount of Zrn more than 20 um in diameter is higher in them. Zrn less than 20 um in diameter is confirmed to become common in mid-T and high-T part of the Grt-Crd zone, and ca. 30 um Zrn without detrital core are rarely found in the high-T part of the Grt-Crd zone. From these observations, these tiny Zrn grains are considered to have nucleated during the Ryoke metamorphism.

On the other hand, monazite grows at amphibolite facies grade and the presence of the melt does not largely affect its recrystallization [4]. In the case of the Aoyama area, Mnz from the migmatites records the prograde growth age of 96.5 +/- 1.9 Ma during regional metamorphism [5]. Using the difference of growth timing of Mnz and Zrn, that is, Mnz being prograde and Zrn being post peak, the duration of metamorphism higher than the amphibolite facies could be constrained, and it is at least ca. 3.5 Ma in the case of the Aoyama area.

[1] Kawakami (2001) JMG 19, 61-75. [2] Cesare et al. (2003) CMP 146, 28-43. [3] Bea et al. (2006) Can Min 44, 693-714. [4] Rubatto et al. (2001) CMP 140, 458-468. [5] Kawakami & Suzuki (2011) Island Arc, 20, 439-453.

Keywords: zircon, glass inclusion, migmatite, LA-ICP-MS, partial melting

The chlorine-rich fluid activity during granulite facies metamorphism in the continental collision zone

HIGASHINO, Fumiko^{1*}, KAWAKAMI, Tetsuo¹, MADHUSOODHAN, Satish-Kumar², ISHIKAWA, Masahiro³, Kenshi Maki¹, TSUCHIYA, Noriyoshi⁴, Geoff Grantham⁵, HIRATA, Takafumi¹

¹Kyoto University, ²Shizuoka University, ³Yokohama National University, ⁴Tohoku University, ⁵Council for Geoscience

Metamorphic fluids play important roles in thermal transport, mass transfer (e.g., Helgeson, 1964), metasomatism (e.g., Meyer & Hemley, 1967) and changing stability field of mineral assemblages (e.g., Powell et al., 1991). In the granulite facies rocks, CO₂-rich fluid has been considered important and studies on Cl-rich fluid are not sufficiently available, because fluid inclusions of the Cl-rich one are less observed than CO₂-rich ones. Some of the reasons Cl-rich fluid inclusion is less found are low viscosity and low wetting angle of brines (Watson & Brenan, 1987). However, Cl-bearing brines are increasingly recognized as playing an important role in high-*T* metamorphic rocks (Newton & Manning, 2010). Using Cl concentration of minerals, it is possible to decipher the Cl-rich fluid activity and its role of them in the lower crust.

We have investigated the field distribution of Cl-rich biotite in the pelitic gneisses of the Sor Rondane Mountains, East Antarctica where Late Proterozoic to Cambrian granulites are widely exposed (Shiraishi et al., 2008). Among more than 20 samples studied, a Grt-Bt-Sil gneiss from Balchenfjella was selected as best suited sample to constrain the *P-T-t* condition of Cl-rich fluid activity. This gneiss contains garnet porphyroblasts (5-10 mm) that have P-rich core with oscillatory zoning in P. The core with oscillatory zoning is discontinuously overgrown by the P-poor rim. The discontinuous zoning in P suggests that garnet porphyroblasts have experienced resorption and overgrowth. The garnet core includes Cl-poor biotite and apatite, and those included at the rim are Cl-rich. Coarse-grained (ca. 100 μm), round zircon grains are exclusively included in the rim of the garnet porphyroblast and also present in the matrix. This mode of occurrence suggests that Cl-rich biotite and apatite and round coarse-grained zircon were formed almost simultaneously.

The *P-T* conditions of the Cl-rich biotite entrapment in the garnet rim were estimated to be ca. 800 °C and 8 kbar, and those of the peak metamorphic condition were ca. 850 °C and 11 kbar, using Grt-Bt geothermometer and GASP geobarometer (Hodges & Spear, 1982). The f_{HCl}/f_{H_2O} ratio of the fluid in equilibrium with Cl-rich biotite (Selby & Nesbitt, 2000) and apatite (Piccoli & Candela, 1994) in the garnet rim are ten times larger than that in equilibrium with Cl-poor biotite and apatite in the matrix and the garnet core. The LA-ICP-MS U-Pb dating of the coarse-grained zircon included in the garnet rim gave a concordia age of 600 ± 13 Ma. Therefore, the Cl-rich fluid infiltration took place at near the metamorphic peak condition of ca. 800 °C and 8 kbar at 600 ± 13 Ma and formed Cl-rich biotite and apatite (e.g., Sisson, 1987).

The field distribution of Cl-rich fluid activity shows somewhat linear distribution. Some of them are located near the ductile shear zones (Ishikawa et al., 2011), and suggesting its relation to high-strain zones (e.g., Kullerud et al., 2001). Regional distribution of near-peak metamorphic Cl-rich fluid activity in the Sor Rondane Mountains implies that it is one of the major phenomena in the continental collision processes.

Keywords: chlorine, fluid, biotite, apatite, Sor Rondane Mountains

High-T metamorphic rocks and exhumation process of the Tseel terrane, SW Mongolia

ULZIIBUREN, Burenjargal^{1*}, OKAMOTO, Atsushi¹, TSUCHIYA, Noriyoshi¹

¹Graduate School of Environmental Studies, Tohoku University

Tseel area, Tseel metamorphic terrane, SW Mongolia is mainly composed of metapelitic gneisses, amphibolites (sillimanite zone) and intruded by ring structure of the granitoids (sillimanite zone) and covered by sediments. Three mineral zones in the pelitic gneisses determined based on the petrographic study and mineral assemblages of each zones as follows: (1) Biotite zone (Bt + Pl + Qtz + Ms + Chl + Cal) (2) Garnet zone (Grt + Bt + Pl + Qtz +/- Chl +/- Ru + Il + Cal) (3) Sillimanite zone (Grt + Bt + Pl + Qtz + Sil + Ms + Chl +/- Crd +/- St + Ru + Il + Cal +/- Fe-Mg Amp). Biotite zone observed in most area of southern and northern parts of Tseel area. Garnet zone is determined between biotite and sillimanite zones. The sillimanite zone is observed in the central parts of Tseel area. All mineral zones observed along the foliation from E to W trends at Tseel area.

We showed the characteristic features of chemical compositions of garnet, biotite and plagioclase in the pelitic gneisses and determine the P-T paths deduced from the garnet chemical zoning using thermodynamic equilibria in Tseel area as follows: Garnet in the pelitic gneisses shows almandine-rich compositions and some samples show clear compositional zoning. Based on the Ca (grossular) composition of garnet, the sillimanite zone is subdivided into sillimanite A and B zones. Garnet in the garnet zone shows relatively homogeneous compositions in the range of (530-600 °C and 6.0-10.0 kbar). The garnet chemical compositions in the sillimanite B zone show a homogeneous composition in a range of (620-750 °C and 1.8-6.0 kbar), and sometimes contain the retrograde rim produced by post-growth diffusion. The garnet chemical zoning in the sillimanite A zone typically is divided into three zones from core to rim: zone 1 (high Ca with homogeneous), zone 2 (decrease in Ca), and zone 3 (low Ca). The compositions of zones 1 and 2 correspond to those of garnet in the garnet and sillimanite B zones, respectively, and zone 3 indicate the effect of the post-growth diffusion. The P-T conditions during garnet growth were calculated by garnet-biotite geothermometry and garnet-biotite-plagioclase-quartz geobarometry. In application of geothermobarometry to garnet zonings, we calculated P-T conditions for four cases with highest and lowest XMg, Bt and XAn, Pl, that provide the possible P-T ranges for a given garnet composition and constrain roughly the shape of P-T path. The shapes of the P-T paths obtained for individual garnets are similar among the four cases, and the differences in pressure and temperature among the four cases are within 15 °C and 2.5 kbar, respectively. The P-T conditions estimated from the garnet zone are in a range of 530-600 °C and 6.0-10 kbar, that are located at the kyanite stability field. The P-T conditions estimated from the sillimanite B zone are in a range of (570-690 °C and 1.8-8.0 kbar). The longest P-T path is obtained from the samples in the sillimanite A zone, such as sample 3001, which shows a decompression from zone 1 (530-570 °C and 6.0-9.6 kbar) to zone 3 (570-620 °C and 2.0-4.0 kbar), corresponding with a change from the kyanite stability field to the sillimanite stability field. Zone 2 shows an increase in temperature by ~40 °C and a decrease in pressure by 4-6 kbar.

The intrusions of the granitoids occurred during the exhumation of the Tseel terrane, and produced the regional anticline structures and mineral zones. The high temperatures and homogeneous garnet zoning of the sillimanite B zone were resulted from the thermal effects from the granitoids. The sillimanite A and B zones experienced the P-T conditions of the kyanite stability field, that was similar to the garnet zone. The high temperature of the garnet zone cannot be explained solely by the contact metamorphism by granitoids, but the regional high geothermal gradient is expected in the subduction zones.

Keywords: Tseel terrane, garnet, P-T paths, exhumation process

Timing of HP and HT-UHT metamorphism in the Red River shear zone, northern Vietnam: Paleogene vs. Permo-Triassic

NAKANO, Nobuhiko^{1*}, OSANAI, Yasuhito¹

¹SCS., Kyushu Univ.

The Red River shear zone is left-lateral shear zone caused by collision of the India to the Eurasian continent at the Paleogene. Although most metamorphic rocks were reset by the deformation, recent U-Pb and Th-Pb in-situ chronological investigations have suggested complex ages from 262 to 25 Ma. In this presentation, we make clear the timing of the high-grade metamorphism using HP and HT-UHT granulites.

To realize the metamorphic conditions and timing of the high-grade event, we use the fluid inclusion technique in association with in-situ U-Pb zircon age. Abundant fluid inclusions are observed in garnet, corundum, staurolite, while are rare in quartz and zircon. Raman analysis shows that all fluid inclusions are composed of CO₂. Their average densities calculated from two Raman peaks are 1.00 +/- 0.06, 1.07 +/- 0.04, 1.09 +/- 0.03, 0.29 +/- 0.07, and 1.15 +/- 0.05 g/cm³ for garnet, corundum, staurolite, quartz, and zircon, respectively. The low-density CO₂ fluids in quartz imply that garnet and corundum have grown up at the different stage than quartz. The estimating pressure-temperature condition based on the mineral paragenesis (exclude quartz) and isochemical phase diagrams using whole rock chemistries of 3 rock types (garnet-corundum-sillimanite, garnet-spinel-sillimanite, and garnet-corundum-spinel granulites), former eclogite-facies (>2.0 GPa at 800 C) metamorphism and subsequent decompression under granulite-facies condition (>1000 C at 1.5 GPa) are identified.

U-Pb zircon ages show wide range from 265 to 36 Ma, however dark luminescent core of zircon containing high-density CO₂ inclusions yields a concordia age of 257 +/- 8 Ma, clearly indicating time of high-grade metamorphic event. This simple technique of combining Raman microscopy and U-Pb age could be easily and widely applicable to evaluate the zircon age and mineral associations in considering pressure-temperature-time evolution of multi-metamorphic events.

Keywords: in-situ U-Pb dating, fluid inclusion, high-grade metamorphism, Red River shear zone, Vietnam

Monazite age of pelitic gneiss in the Lhenice shear zone (southern Bohemian Massif) and its significance

KOBAYASHI, Tomoyuki^{1*}, Simon L. Harley², HIROI, Yoshikuni¹, HIRAJIMA, Takao³

¹Department of Earth Sciences, Faculty of Science, Chiba University, ²Department of Geology and Geophysics, University of Edinburgh, ³Department of Geology and Mineralogy, Graduate School of Science, Kyoto University

Recently, multiple equilibrium stages were identified from Grt-rich gneiss at Kti? in the Lhenice shear zone, located along the western margin of the Blansky les massif (Kobayashi et al., 2011). The characteristic matrix mineral assemblage of the Grt-rich gneiss is Crd+Sil+Bt+Grt+/-Spl with Qtz+Kfs+Pl. The coarse-grained Grts are commonly composed of dusty core and clear rim. The dusty core is defined by the alignment of fluid (CO₂-N₂) and micro-sized solid inclusions (Qtz, Kfs, Pl, Rt, Ap, Mnz and Zrn) of which outline are hexagonal in shape. Some coarse-grained (> 3 mm) Grts show chemical heterogeneity both in major and minor elements; Grs-content is homogeneous and high (Xgrs = 0.27) in an apparent core of the grain and continuously decreases towards the rim (Xgrs = 0.02). However, Prp-content shows an inverse pattern against Grs-content, i.e., Prp content is low and constant (Xprp = 0.03) in the core and gradually increases towards the rim (up to Xprp = 0.28). The outline of Grs and Prp content contours show symmetrical hexagonal shapes. Phosphorous (P)-content is almost below the detection limit of EPMA in the apparent core but it is high at the margin of the grain with local development of P-poor outermost rim. The outline of P-poor core shows a hexagonal shape, similar to that of Grs and Prp content contours. The geothermobarometry, based on the mode of occurrence of constituent minerals and the zoning pattern of Grt, depicts following developing history of the host rock, such as, a prograde stage defined by the assemblage of P-poor Grt core (Grs=27) + Pl (An₁₁₋₁₅) under 1.5-2.3 GPa at 700-900 °C (Stage 1), a subsequent Grt-rim forming stage represented by P-rich Grt (Grs5) + Pl (An₁₂₋₁₉) + Ky/Sil at 730-830 °C and 1.0-1.3 GPa (Stage 2), and a following decompression stage by the outermost rim of Grt (Grs₂) + Sil + Crd +/- Spl at 740-850 °C and 0.6-0.8 GPa (Stage 3) (Kobayashi et al., 2011).

To evaluate age of multiple equilibrium stages, chemical Th-U-Pb isochron method (CHIME) Mnz age dating was carried out for Grt-rich gneiss. Mnzs which included in core of Grt show bimodal grain size; coarse-grained (1 mm in diameter) and fine-grained (10 micrometer in diameter). Mnzs which included in rim of Grt have fine-middle grained size (10 micrometer to 0.5 mm in diameter). Mnzs in matrix have middle-coarse grained size (0.5 mm to 3 mm in diameter). Most of middle-coarse grained Mnzs show a chemical zoning; relatively low Th constant in the core and high Th constant in the rim. The Mnz grains which included in the core of Grt give an age of 336±11 Ma. The Mnz grains which included in the rim of Grt give an age of 335.4±7.2 Ma. The Mnz grains in matrix give 334.9±3.9 Ma. These results suggest that the studied rock experienced very fast exhumation from stage 1 to stage 3. Furthermore, felsic volcanic rock-like inclusions (FVRLI) are found from the core and rim of coarse-grained Grt. The FVRLI mainly consists of micro to cryptocrystalline aggregate of Qtz, Pl and Kfs. The FVRLI show spherulitic, granophyric, and porphyritic textures in addition to quartz dendrites. These features of the inclusions are similar to those of "nanogranites" which are FVRLIs enclosed within Grt in high- to ultrahigh-temperature pelitic migmatites and/or granulites as reported by Cesare et al. (2009) and Hiroi et al. (2011). Cesare et al. (2009) concluded that nanogranites are the crystallized anatectic melts which were trapped by peritectic minerals growing during partial melting. The FVRLI in this study may suggest that partial melts formed during early high-pressure metamorphism stage (stage 1) and trapped by garnet have undergone nonequilibrium crystallization under specific conditions of continuous rapid cooling. The possible corresponding geotectonic process may be "vertical extrusion and horizontal channel flow" (e.g. Schulmann et al., 2008) proposed for the Himalayan-Tibetan and Variscan (Bohemian Massif) orogeny.

Keywords: Bohemian Massif, Monazite age, Grt-rich gneiss, Partial melting, Felsic volcanic rock-like inclusions (FVRLI), Rapid cooling

Thermobaric structure in low P/T type metamorphic belt -Case study of the Ryoke metamorphic belt in the eastern Yamaguc

SUGAWARA, Yu^{1*}, IKEDA, Takeshi¹

¹Kyushu University

Recent seismic studies and EM monitoring imply that there is thermal heterogeneity in present subduction zone (Nakajima et al., 2009., Ogawa et al., 2001), furthermore numerical simulation predicts time variation of these heterogeneities (Iwamori, 2000). On the other hand, by petrological method, the same phenomenon is observed in ancient crust and mantle (Baker, 1987) and the Gibbs method is enabled to estimate P-T path (Okamoto and Toriumi, 2001). In this way, seismic and petrological studies are complementary, extensive data which is corresponded to seismic research should be provided by petrological studies. In this study, we provide specific data on thermally heterogeneity by using petrological method, for the Ryoke metamorphic belt in the eastern Yamaguchi Prefecture.

Using pelitic and psammitic mineral assemblages, the study area can be divided into seven zones, i.e. chlorite, chlorite-biotite, biotite, muscovite-cordierite, K-feldspar-cordierite, garnet-cordierite, sillimanite-K-feldspar zones. The K-feldspar-cordierite, garnet-cordierite and sillimanite- K-feldspar zones continue to east part where Ikeda (1998) performed metamorphic zonation. The garnet-cordierite zone decreases its width toward west and disappears in around Hikari city. On the other hand, the sillimanite-K-feldspar zone is widely distributed in Murotsu-Kudamatsu-Oshima area.

Pressure-temperature conditions of seven samples from the sillimanite-K-feldspar zone and three samples from the garnet-cordierite were estimated by using the garnet-biotite thermometer of Hodges and Spear (1982) and the relative geothermometry of Ikeda (2004). Addition of result of Ikeda (2004) enables us to reveal the thermobaric structure of this area. Temperature increases toward southeast, exceeding 800 degree in relatively-limited area around Hizumi area, and decrease further toward south below 800 degree. In contrast, pressure increases toward southeast monotonously and it reaches 5-6kbar in the south of Hizumi district-Cape Kandori. The isotherms are oblique to isobaric lines, suggesting that the crust have thermally heterogeneity at the same depth.

Applying the concept proposed by Miyazaki (2007) to the present results suggests that rate of melt migration increases and duration of melt migration decreases in the following order: Yanai-Hizumi area which is east part, Oshima-Murotsu area which is south part, and Kudamatsu area which is west part in study area. Additionally, estimated thermobaric structure enables us to impress west and south parts were formed in same depth.

References:

- Baker,A.J.,1987. *J.Metamorphic Geol.*,5;101-118
- Ikeda,T.,1998. *J.Metamorphic Geol.*,16:39-52.
- Ikeda,T., 2004. *Contrib Mineral Petrol*, 146: 577-589.
- Iwamori,H.2000.*Earth Planet.Sci.Lett*,181,131-144.
- Hodges and Spear,1982. *Am Mineral.*,67:118-1134
- Ogawa ,Y., Mishina,M., Goto,T., Satoh,H.,Oshiman,N., Kasaya,T., Takahashi,Y., Nishitani,T., Sakanaka,S., Uyeshima,M., Takahashi,Y., Honkura,Y. and Matsushima,. 2001. *geophys. Res. Lett.*, 28, 3741 -3744.
- Okamoto,A and Toriumi,M., 2001.*Contrib.Mineral.Petrol.*,141;268-286.
- Miyazaki,K.,2007. *Island Arc.*,16,69-82.
- Nakajima,J., Hirose,F. and Hasegawa, A. 2009.*J.geophys.Res.*114, B08309.

Keywords: metamorphic rock, Low P/T metamorphic belt, Ryoke belt, subduction zone, Thermobaric structure, biotite

Dehydration breakdown of antigorite and the formation of b-type olivine CPO

NAGAYA, Takayoshi^{1*}, WALLIS, Simon¹, MICHIBAYASHI, Katsuyoshi², MIZUKAMI, Tomoyuki³

¹Earth & Planetary Sci., Nagoya Univ., ²Instit. Geosci. Shizuoka Univ., ³Nat. Sci. Tech., Kanazawa Univ.

Observations of the seismic waves in the mantle wedge (MW) above many subduction zones display a fast seismic direction perpendicular to plate motion. This cause of this anisotropy is considered by many workers to be due to the existence of b-type olivine (Ol) CPO in the MW with [100] axes concentrated perpendicular to the flow direction parallel to plate motion.

Deformation experiments show that b-type CPO patterns can be formed as a result of dislocation creep under water-present high-stress conditions⁽¹⁾ and thermomechanical models have been used to suggest that conditions necessary to form b-type Ol CPO may be achieved close to subduction boundaries⁽²⁾.

However, work on hydrated antigorite (Atg) bearing forearc mantle shows that the presence of Atg destroys any preexisting CPO and prevents strong Ol CPO from being developed⁽³⁾. The reduction in CPO strength is thought to be due to slip concentrating in the weaker Atg layers with associated grain-boundary sliding occurring between Ol and Atg. This result suggests that b-type Ol CPO fabrics are unlikely to be developed close to subduction boundaries where Atg is expected to be stable.

Nevertheless, several examples of naturally occurring b-type Ol CPO have been reported. A review of published reports shows some of these were formed at relatively high temperatures and low stress, which is incompatible with the predictions from experimental work. Natural samples also lack evidence for c-slip, which is expected for the formation of b-type Ol CPO by dislocation creep.

These considerations show that the formation of naturally occurring b-type Ol CPO is not well understood.

Here we document b-type Ol CPO formed by the topotaxial growth of Ol on Atg from the Happo-One region of the Hida Marginal belt, Japan. Before dehydration and conversion to Ol, the Atg had a strong preexisting CPO due to deformation at relatively shallow levels under low temperatures and hydrated conditions. In the Happo-One region, non-deformed secondary Ol formed in veins as a result of the dehydration of foliated Atg, due to contact metamorphism⁽⁴⁾.

The CPO of the vein Ol shows a strong b-Type fabric that is characterized by a c-axis concentration parallel to the stretching lineation and a b-axis concentration normal to the foliation. The CPO of the Atg bordering the vein shows a strong concentration of c-axes at a high angle to the foliation and a strong alignment of b-axes parallel to the lineation. Many recent studies have shown this type of Atg CPO is the most widespread in the forearc MW.

Two types of topotaxial growth relationships are known between Ol and Atg: in both cases [010]atg is parallel [001]ol but [010]ol may be parallel to either [100]atg (type 1) or [001]atg (type 2)⁽⁵⁾. The observed relationships between the Ol and Atg CPO patterns in this study imply type 2 topotaxial relationships between the two minerals.

Atg-bearing mantle is predicted to be a widespread component of forearc mantle. As this material is dragged down by the traction of the downgoing slab, it will become deformed and foliated. When this foliated antigorite schist reaches sufficiently high T and P conditions, it will undergo dehydration. Our results show that when this dehydration occurs, the newly formed Ol is likely to have a b-type Ol CPO. This topotactic Ol CPO can form in the MW away from the coldest part immediately adjacent to the subduction boundary.

The CPO formation mechanism reported here can reconcile the differences between the laboratory and natural examples of b-type Ol CPO patterns and also explain why such b-type CPO is found associated with subduction zones.

References

- (1) Katayama, I. & Karato, S. 2006, *Phys. Earth Planet. Inter.* 157 (1-2), 33-45
- (2) Kneller, E.A. et al. 2005, *Earth Planet. Sci. Lett.* 237, 781-976
- (3) Wallis, S. R. et al. 2011, *J. Geol. Soc. London, Special Publications* 360, 113-127
- (4) Nozaka, T. 2005, *J. Metamorphic Geol.* 23 711-723
- (5) Boudier, F. et al. 2010, *J. Petrol.* 51, 495-512


Keywords: topotaxy, olivine, CPO, antigorite, seismic anisotropy

Spinifex-textured olivine-enstatite rock and clinoenstatite formed by high pressure breakdown of antigorite

NISHIYAMA, Tadao^{1*}, EGUCHI Hibiki¹, YOSHIASA Akira¹, SUGIYAMA Kazumasa², ARIMA Hiroshi², YUBUTA Kunio²

¹Graduate School of Science and Technology, Kumamoto University, ²Institute for Materials Research, Tohoku University

Occurrence of spinifex-textured meta-peridotite is well known from the Higo Metamorphic Rocks (Mizuta, 1978), Central Kyushu. The rock occurs as massive sheet like bodies in pelitic and basic gneisses, and is composed mainly of elongated crystals of olivine and coarse-grained enstatite. Olivine crystals do not show preferred orientation, and is mostly altered to antigorite. Enstatite is fresh in most cases, but replaced partly by talc in some cases. Small amounts of spinel, tremolite, and anthophyllite occur as secondary minerals, therefore the rock is a metamorphosed olivine-enstatite rock or hartzburgite. Trommsdorff et al.(1998) studied a spinifex-textured olivine-enstatite rock from Spain, and discussed its origin as a breakdown product of antigorite at high pressures (2 GPa) based on its mineral assemblage and texture. They argued that the spinifex texture, either of igneous or metamorphic origin, formed under fluid (melt or metamorphic fluid)-rich environment, resulting in formation of olivine crystals elongated to [001]. Recently, Kendrick et al. (2011) analyzed fluid inclusions from this rock and clarified that they retain the composition rich in chlorine and noble gas elements originated from seawater. These features suggest that a subducting mantle was hydrated and serpentinized by seawater along fractures, which was caused by slab-bending at fore-arc. The rock from Higo has features common to those from Spain, therefore it is natural to consider that the rock formed by dehydration of antigorite by the reaction:



This reaction takes place at about 600-700 C and 2 GPa. Although most of the Higo metamorphic rocks are the product of high T/low P metamorphism of Cretaceous in age (e.g.Obata et al., 1994, Miyazaki 2004), several studies revealed that they contain some mineral assemblages showing high pressure conditions incompatible with the high T/low P metamorphism (Kano and Uruno, 1995; Karakida et al., 1989; Osanai et al., 1998; Maki et al., 2004 and 2009). Our finding shows that the spinifex-textured olivine-enstatite rock shows a pressure condition much higher than that ever known for the Higo metamorphic rocks.

Pyroxene from the olivine-enstatite rock has the composition of En90Fs10-En95Fs5 and is very poor in CaO and Al₂O₃. The pyroxene consists of ortho-enstatite (Pbca) with lamellae of clinoenstatite (P2₁/c) according to the study with a single X-ray diffractometer (Rigaku RAPID) and a transmission electron microscopy (TOPCON, EM-002B at 200 kV). Clinoenstatite has usually the same composition as orthoenstatite, but in some samples it shows slightly iron-rich composition than orthoenstatite. Clinoenstatite lamella occurs along (100) plane with a width of nanometer to several tens of micrometer vertical to the a* direction. Based on the rule of absent reflection and a model analysis of clino- and ortho- structures, the space groups, lattice parameters and structures of the two pyroxenes are confirmed.

The occurrence of clinoenstatite from the terrestrial rocks is very limited. Recently, high pressure phase of clinoenstatite (C2/c) has been found by several experiments (e.g. Kanzaki, 1991), and some authors discuss it as the possible candidate for the origin of natural clinoenstatite from peridotites and clinopyroxenites especially of the ultra-high pressure metamorphic terrane. Zhang et al.(2002) reported wide occurrence of clinoenstatite in garnet-pyroxenites from the Dabie-Sulu ultra-high pressure metamorphic terrane, and estimated the formation condition to be about 750 C and 6.7 GPa. In the case of Higo, the condition of spinifex-textured olivine-enstatite rock is about 600-700 C and 2 GPa,so it is natural to consider at present that clinoenstatite lamellae and domains formed by shear stress from orthoenstatite, however, the possibility of transition from high pressure phase still remains.

Keywords: serpentinite, spinifex texture, clinoenstatite, ultra-high pressure metamorphic rock, Higo Metamorphic Rocks

Petrogenesis and implications of jadeite-kyanite eclogite from the Iratsu body of the Sanbagawa belt, SW Japan

ENDO, Shunsuke^{1*}, TSUBOI, Motohiro²

¹Nagoya University, ²Kwansei Gakuin University

Discovery of a new occurrence of jadeite- and kyanite-bearing eclogite from an outcrop within the Iratsu body of the subduction-type Sanbagawa belt, SW Japan, allows us to assess existing solid-solution models for clinopyroxene, and obtain insights into spatial variation in P-T conditions and/or H₂O activity during the eclogite-facies metamorphism of this region. The jadeite + kyanite assemblage is stable at higher P-T conditions or lower H₂O activity compared to paragonite. There is no significant difference in bulk rock composition between the newly found jadeite-kyanite eclogite and paragonite eclogite, which is the predominant eclogite type in the Iratsu body. The jadeite-kyanite eclogite is a medium-grained massive metagabbro consisting mainly of garnet, omphacite, kyanite, quartz, epidote, phengite, subcalcic amphibole and rutile. Pre-eclogitic relics of sodic augite (Jd₇₋₂₄Acm₆₋₁₆), actinolitic hornblende, Fe-rich garnet (Alm₆₂₋₇₁Grs₁₉₋₂₅Prp₅₋₁₃Sps₃₋₄) and magnetite are sporadically preserved. Eclogitic garnet (Alm₅₄₋₆₁Grs₁₆₋₂₀Prp₂₀₋₂₅Sps₁₋₂) optically shows a dusty appearance due to abundant microscopic inclusions of kyanite, quartz, epidote, phengite, omphacite (Jd₄₀₋₅₅Acm₇₋₁₃) and impure jadeite (Jd₆₂₋₈₆Acm₀₋₇). Jadeite is exclusively present as inclusions in garnet. The miscibility gap between the ordered omphacite (P2/n) and disordered impure jadeite (C2/c) progressively narrows during garnet growth, implying the temperature of the solvus apex coincides with the thermal peak of metamorphism. The observed compositional gap and the result of garnet-clinopyroxene Fe²⁺-Mg exchange thermometry are consistent with the phase diagram calculated in the pseudo-binary augite (Di₆₆Hd₁₄Acm₂₀)-jadeite system by using the newest solid-solution model of Deiner and Powell (2012), and the calculated apex of the omphacite-jadeite solvus is at 625 deg.C. However, calculated pseudosections with XRF-derived bulk rock composition and the solid-solution model have no jadeite stability field for any reasonable values of P, T, X_{Fe³⁺} and M_{H₂O}, although the matrix assemblage is satisfactorily reproduced. This may suggest effective bulk composition around growing garnet was significantly different from the XRF-derived bulk rock composition. The localized feature of eclogite-facies equilibration is also inferred from the observed microstructures such as Omp + Qz + Amp pseudomorphs (prograde symplectites) after igneous augite. We obtain the jadeite stability field in the calculated pseudosections by subtracting augite from the XRF-derived bulk composition. High chlorine contents of amphibole (<1.5 wt% Cl) and apatite (<7.0 wt% Cl) indicate that the eclogite-facies equilibration was triggered by an influx of saline fluids. Multi-equilibrium thermobarometry for the assemblage Grt + Omp + Ky + Ph + Ep + Qz gives metamorphic P-T conditions of around 2.3 GPa and 600 deg.C. The estimated high-P conditions are also supported by high residual pressure (max. delta omega-1 = 13.3 cm⁻¹, equivalent to 0.85 GPa) of quartz inclusions in garnet. These results imply the presence of a significant metamorphic pressure gradient within the Iratsu body, and detailed baric structure of this region will be revealed by further application of the quartz-in-garnet barometry (Enami et al. 2007).

Reference: Deiner and Powell (2012) *J Metam Geol* 30, 113-130; Enami et al. (2007) *Am Mineral* 93, 1303-1315.

Keywords: eclogite, Iratsu body, jadeite, metagabbro, omphacite

Metamorphism of garnet glaucophane schists from the Bizan area, Sambagawa metamorphic belt, eastern Shikoku, Japan

Kabir Md Fazle^{1*}, Takasu Akira¹, Nakamura Mayuko¹, Kondo Yuki¹, Kainuma Masaaki¹

¹Department of Geoscience, Shimane University, Japan

The Bizan area is located in the Tokushima Prefecture of eastern Shikoku, Japan and it is mainly composed of pelitic schists, basic schists and siliceous schists with minor garnet glaucophane schists. Faure (1983) suggested that there occur a melange with tectonic blocks of serpentinite, metagabbro and garnet-amphibolite (garnet glaucophane schists in this study) along the ductile shear zone between spotted and non-spotted zones in the Bizan area. Garnet glaucophane schists consist mainly of garnet, amphibole, epidote, phengite (Si 6.41-6.80 pfu), paragonite, chloritoid, chlorite, albite, rutile, titanite, ilmenite and quartz. A schistosity is defined by preferred orientation of glaucophane, phengite and chlorite.

Garnets are almandine-rich composition and display prograde growth zoning with decreasing X_{Sps} (0.18-0.01) and increasing X_{Alm} (0.54-0.70) from the core to the rim. The cores of garnet contain inclusions of chloritoid (X_{Mg} 0.08-0.23), chlorite, epidote, phengite (Si 6.41-6.55 pfu), paragonite, titanite and quartz and polyphase inclusions of phengite + paragonite + epidote + chloritoid + chlorite + quartz assemblage. The rims of garnet contain inclusions of amphiboles (glaucophane; zoned amphibole with barroisite core and glaucophane rim), epidote and quartz and polyphase inclusions of glaucophane + epidote. Garnets are partly replaced by chlorite. Glaucophanes in the matrix contain inclusions of phengite, epidote, chlorite, titanite and quartz. Some of them are partly replaced by Wnc, Brs, Mg-Ktp and Act at their rims and cracks. Phengites (Si 6.50-6.80 pfu) in the matrix are contain inclusions of glaucophane, epidote and chlorite. Some large grains of epidotes and chlorites contain inclusions of glaucophane, phengite, epidote, titanite and quartz.

Textural relationship, mineral assemblage and thermobarometric results suggest a polyphase tectonometamorphic evolution of the garnet glaucophane schists. The mineral assemblages of polyphase inclusions within the cores of the garnet such as barroisitic amphiboles, chloritoid, chlorite, epidote, phengite, paragonite and quartz constrain the P - T conditions of a prograde stage at 450-500°C and 9-11 kbar (by THERMOCALC) at epidote-blueschist facies metamorphic conditions. The rims of the porphyroblastic garnets include inclusions of glaucophane, epidote, quartz and schistosity-forming matrix phengite suggesting the peak metamorphic mineral assemblages. THERMOCALC average P - T calculation suggests a metamorphic condition of the eclogite facies metamorphism of 550-600°C and 17-19 kbar. Porphyroblastic garnets are partly replaced by chlorite at their rims, and matrix glaucophanes are replaced by Wnc, Brs, Mg-Ktp and Act along rims and cracks suggests a retrograde metamorphism took place at the epidote-amphibolite facies to follow a clockwise decompression path.

Large grains of epidotes and chlorites in the matrix which contain peak metamorphic mineral assemblages of glaucophane, phengite, epidote and quartz suggesting another high-pressure prograde metamorphism took place. This high-pressure metamorphism can be correlated with the Sambagawa metamorphism in the Besshi area, central Shikoku (Aoya, 2001; Kabir and Takasu, 2010a, b).

The eclogite facies metamorphism followed by another high-pressure metamorphism as the Sambagawa metamorphism is first described from the garnet glaucophane schists in the Bizan area, Sambagawa metamorphic belt. The metamorphic evolution is similar to that of the eclogites in the Besshi area, central Shikoku (Kabir and Takasu, 2010a, b). Eclogite in the Sambagawa belt occurs mainly in the Besshi area, central Shikoku and slightly in the Kotsu area, eastern Shikoku. This study revealed that the occurrence of eclogites now extend to the Bizan area in eastern Shikoku.

Reference:

Aoya (2001) *J Petrology*, 42, 1225-1248; Faure (1983) *J Geol Soc of Japan*, 89, 319-329. Kabir and Takasu (2010a) *J Meta Geol*, 28, 873-893; Kabir and Takasu (2010b) *Earth Sci*, 64, 183-192.

Keywords: Sambagawa metamorphic belt, garnet-glaucophane schist, eclogite, chloritoid, Bizan area, eastern Shikoku

Metamorphic conditions of kyanite-garnet-chloritoid schists associated with eclogites in the Lake Zone, SW Mongolia

OTGONKHUU, Javkhlan^{1*}, TAKASU, Akira¹, Batulzii Dash², Md. Fazle Kabir¹

¹Geoscience Dept. Shimane University, Japan, ²Mongolian University of Science and Technology, Mongolia

The kyanite-garnet-chloritoid schists associated with eclogites from the Alag Khadny metamorphic complex, Chandman district, Lake Zone, SW Mongolia, consist of garnet, chloritoid, muscovite, phengite, chlorite, paragonite, kyanite, rutile, ilmenite, zircon, quartz and carbonaceous matter. Garnets occur as subhedral to anhedral porphyroblast up to 5 mm across and they are almandine-rich variety in composition. The garnets are zoned with inclusion-rich cores and inclusion-poor rims. The garnets display a prograde pattern of compositional zoning, X_{Sps} decreasing and X_{Prp} increasing from core to rim. The cores contain inclusions of muscovite ($Si=6.06-6.29$ cations per formula unit, *pfu*), paragonite, chlorite, chloritoid and quartz. The rims contain inclusions of kyanite ($Fe_2O_3 < 1.24$ wt%; $Cr_2O\{3\} < 0.03$ wt%), phengite ($Si=6.40-6.63$ *pfu*), chloritoid (X_{Mg} [$Mg/(Fe+Mg)]=0.08-0.18$), chlorite ($X_{Mg} < 0.42$), and quartz. A well-developed schistosity is defined by preferred orientation of chloritoid ($X_{Mg} = 0.11-0.21$), chlorite ($X_{Mg}=0.36-0.53$) and white micas [phengite ($Si=6.57-6.63$ *pfu*) and muscovite ($Si=6.23-6.34$ *pfu*)] in the matrix.

Based on the textural relationship and chemical composition of minerals, following metamorphic stages are distinguished in the kyanite-garnet-chloritoid schists, i.e. (i) pre-peak stage, (ii) peak metamorphic stage, and (iii) retrograde stage.

The porphyroblastic garnets represent a typical prograde zoning, X_{Sps} decreasing and X_{Prp} increasing from core to rim. The pre-peak stage (i) is defined by the mineral inclusions in the cores of the garnets. They are muscovite ($Si=6.06-6.29$ *pfu*), paragonite, chlorite, chloritoid ($X_{Mg}=0.08-0.13$) and quartz, and they indicate relatively low-pressure and low-temperature conditions such as the greenschist facies. The peak metamorphic stage (ii) is defined by the mineral assemblage of the inclusions in the rims ($X_{Prp} < 0.13$) of the garnets, i.e. kyanite, phengite ($Si=6.40-6.63$ *pfu*), chloritoid ($X_{Mg}=0.08-0.18$), chlorite ($X_{Mg}=0.42$), rutile and quartz, and schistosity forming minerals, i.e. chloritoid ($X_{Mg} = 0.11-0.21$), phengite ($Si=6.57-6.63$ *pfu*), and chlorite ($X_{Mg}=0.36-0.53$) coexisting with the rims of porphyroblastic garnet. THERMOCALC (V. 3.33) (Powell and Holland, 1994) calculations for the rim of the garnet coexisting minerals of kyanite, phengite, chloritoid, and chlorite yielded P-T conditions of $T=575-585^\circ C$ and $P=10-11$ kbar of high-pressure epidote-amphibolite to low-pressure eclogite facies conditions of the high-pressure intermediate type metamorphism which are distinctly lower in metamorphic pressure than accompanied eclogites ($T=590-610^\circ C$, $P=20-22.5$ kbar; Stipska et al., 2010) even though the similar temperature conditions. However, $^{40}Ar/^{39}Ar$ muscovite plateau ages of the eclogite (543 +/- 3.9 Ma) and kyanite-garnet-chloritoid schist (537 +/- 2.7 Ma) are similar (Stipska et al., 2010), indicating simultaneous exhumation of both metamorphic rocks to the crustal level. There is no evidence of high-pressure type metamorphic event similar to the eclogites in the kyanite-garnet-chloritoid schists. These fact suggest that subduction within low-geothermal gradient conditions to produce the eclogites occurred first, subsequently relatively high-geothermal gradient metamorphism for the kyanite-garnet-chloritoid schists took place, and then whole sequence of metamorphic rocks were exhume to the shallower levels at ~ 540 Ma.

Reference:

Powell, R. and Holland, T.J.B. (1994). *Amer. Miner.*, 79, 120-133

Stipska, P., Schulmann, K., Lehmann, J., Corsini, J., Lexa, O. and Tomurhuu, D. (2010) *J. of Meta. Geol.*, 28, 915-933

Keywords: Kyanite-garnet-chloritoid schist, Eclogite, Alag Khadny metamorphic complex, Lake Zone, Mongolia

Raman spectra and reflectances of carbonaceous matters showing graphitization temperatures at around 300 - 400 C

MURATA, Masami^{1*}, Wataru Sato¹, HOSHINO, Kenichi²

¹Fac. Sci., Hiroshima Univ., ²Grad. Sch. Sci., Hiroshima Univ.

The degree of graphitization of carbonaceous matter (CM) has long been investigated using X-ray diffraction analyses, optical (reflectance) studies and Raman spectroscopies to analyze thermal histories of sedimentary rocks. Hirota and Hoshino (2007) noted that the degree of graphitization may reveal much detailed thermal histories than those recorded in the mineral assemblages. Komorek and Morga (2003) also demonstrated that the reflectances of CMs increased by only several hours heating in an oven. Therefore, we may expect the CM geothermometers for analyses of short-range thermal events such as dike intrusion, vein formation and faulting.

However, the thermometer using Raman spectrum of CM proposed by Beyssac et al. (2002) can be applied to the temperature range from 640 to 330 C with poor correlations at low temperatures. On the other hand, the thermometers by reflectance of CM ('vitrinite reflectance') are probably limited below 300 C (e.g., Hashimoto et al, 2004). Therefore, we have studied detailed characteristics of Raman spectra and reflectances of CMs showing their graphitization temperatures at around 300 - 400 C to investigate a future possibility of the CM geothermometer as a tool for analyses of the short-range thermal events.

We collected pelitic rock samples of the Kuga formation of the Jurassic accretion complex from the Yasaka contact aureole, Yamaguchi Prefecture, where the Hiroshima-type granite of the Cretaceous age intruded. Takami et al. (1993) classified the contact aureole into the cordierite zone (ca. < 1 km from the contact) and the biotite zone (1 - 2 km wide). Takami and Nishimura (2000) mentioned that the apparent d002 of CM decreases sharply in the biotite zone at around 1 km from the contact.

Two types of CM can be identified under microscopic observations of the collected rock samples. Hereafter, we may call them temporarily as M-type and P-type CMs. The former shows visible reflection pleochroism as well as anisotropism, while the latter does non- or weak ones. Therefore, the P-type CM is probably collinite for so-called 'vitrinite reflectance' measurements. They can be identified clearly also by their Raman spectra. The G bands of the P-type CMs are broad, while those of the M-type are characteristically narrow and sharp. It is interesting that the P-type CM could not be observed in the samples collected from the vicinities of the contact with granite.

The R2 ratio of the Raman spectrum of the M-type CM shows a systematic change with a distance from the contact. It increases from 0.47 (= 430 C, estimated from the equation of Beyssac et al., 2002) at Loc. 03 (0.1 km away from the contact) to 0.60 (370 C) at Loc. 10 (2.4 km). On the other hand, although the ratio of the P-type CM is the highest as 0.77 (300 C) at Loc. 10, the ratio in the other samples closer to the contact does not vary with the distance and is stable as 0.75 (310 C), implying the lower limit of the ratio of the P-type CM.

The P-type CM from Loc. 10 shows the lowest reflectance as 3.2 % (= 270 C, from the equation of Barker, 1988), while those from the other localities are around 5.0 % (310 C) and show no significant variation with the distance, implying also the upper limit of the reflectance of the P-type CM. Although the reflectance of the M-type CM is slightly higher than that of the P-type CM in the sample from Loc. 10, no obvious difference between the two could be seen in the other samples.

It should be emphasized that there may be a size dependency of the M-type CM on its R2 ratio in the samples close to the boundary, that is, a smaller CM tends to show a higher temperature. Hence, it can be expected to apply the R2 ratio of the M-type CM not only as a thermometer but also as a 'chronometer' for short-range thermal events.

An estimation of a P-T condition of the vein-formation in the footwall of the Nobeoka thrust will be presented as an example of the analyses of the thermal events.

Keywords: carbonaceous matter, Raman, reflectance, geothermometer

Study of negative residual pressure of quartz in garnet: Quartz Raman spectra of high-temperature metamorphic rocks

KOUKETSU, Yui^{1*}, ENAMI, Masaki¹, IKEDA, Takeshi²

¹Graduated School of Environmental study, Nagoya University, ²Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University

The volume of quartz inclusions in garnet porphyroblasts changes with temperature and pressure until the rock reaches the surface of the earth. Positive residual pressure is preserved by the compressional stress when the volume of the included quartz becomes larger than that of the surrounding garnet, whereas negative residual pressure is preserved by the elongational stress when the volume of quartz becomes smaller than that of the garnet. The peak position of quartz Raman spectrum shifts toward high wavenumbers with increasing compressional stress and toward low wavenumbers with increasing elongational stress. Enami et al. (2007) measured the peak shift of quartz Raman spectra of several samples and estimated the residual pressure. They showed that there is a positive correlation between the residual pressure and metamorphic pressure. They also showed that there is almost no dependence of the residual pressure on temperature or on the chemical composition of garnet, and proposed that this technique can be applied to a quartz Raman barometer. Since quartz Raman barometry is independent of the thermodynamic model, this barometer is applied as one of the indices of pressure estimation of high-pressure metamorphic rocks in several studies (e.g., Kouketsu et al., 2010).

Recently, the quartz inclusions that preserve the negative residual pressure in the Higo metamorphic rocks were reported (Nishiyama and Aikawa, 2011). The simple elastic model used by Enami et al. (2007) shows almost no P - T range where the quartz inclusion acquires negative residual pressure (Van der Molen, 1981). Hence, Nishiyama and Aikawa (2011) interpreted that the negative residual pressure was acquired by the transition of the included quartz from beta- to alpha-phase.

In this study, we analyzed the rocks formed under high-temperature conditions to investigate in detail about the negative residual pressure that is not evaluated by the conventional quartz Raman barometer. We analyzed samples from Yanai Ryoke, collected from 5 different metamorphic zones and from East Antarctica. The metamorphic P - T conditions of Yanai Ryoke were estimated in detail by Ikeda (2004); Chl-Bt, Ms-Crd, and Kfs-Crd zones are stable in alpha-quartz, and Grt-Crd zone is stable in beta-quartz. The Sill-Kfs zone is near the alpha-beta transition line. The samples from East Antarctica are estimated to have formed at ultra-high temperatures.

The results of the analysis showed that the quartz included in the garnets of every sample from Yanai Ryoke have comparable values of negative residual pressure. In the sample from East Antarctica, quartz with very low residual pressure was commonly observed. This result implies that negative residual pressure is acquired when the metamorphic condition of the quartz included in the garnet is stable whether in alpha or beta phases. This result implies that the explanation provided by Nishiyama and Aikawa (2011) is insufficient. Hence, we reexamined the conventional elastic model; it became clear that the model of Van der Molen (1981) is not suitable for estimating the relation between the residual pressure and metamorphic pressure, and that the model of Zhang (1998) is more appropriate for the quartz Raman barometer. We improved the model to estimate the residual pressure by taking into consideration the pressure dependency of the bulk modulus and the temperature dependency of the thermal expansion.

The new model can explain the negative range of the quartz Raman barometer qualitatively. It was shown that the quartz acquires negative residual pressure even if it had been included in the garnet under the metamorphic conditions in a stable alpha-phase. In addition, it was also shown that the quartz included under the ultra-high temperature conditions, as in the case of the sample from Eastern Antarctica, acquires much lower values of residual pressure than that observed in the case of samples from Yanai Ryoke.

Keywords: negative residual pressure, quartz Raman barometer, high temperature metamorphism, inclusion-host system, elastic model

Correlation stress history with statistical analysis on mineral composition at small brittle fault in the borehole core

TANAKA, Nobuaki^{1*}, FUJIMOTO, Koichiro¹, SHIGEMATSU, Norio²

¹Tokyo Gakugei University, ²MEXT

The Median Tectonic Line (MTL), the largest on-land fault in Japan, has a long history of displacement, and the fault rocks deformed under variable conditions are distributed. The analysis of internal structure of the MTL, therefore, helps to improve our understandings of variable fault behavior depend on the physical conditions and development of fault zone.

AIST drilled a borehole penetrating the MTL for predicting Tonaikai-Nankai Earthquake at Iitaka, Matsusaka, Mie prefecture. The drilling length is 600m. It crosses MTL at the depth of 473.9m. Hangingwall of the MTL consists of Ryoke-derived tonalitic mylonite and footwall of the MTL consists of fractured rocks derived from Sanbagawa metamorphic rocks.

The rocks in the hangingwall experienced the four kinds of stress pattern after the mylonitization. These are stresses which caused normal faultings (vertical compressive stress (Stress-A) and North-South tensional stress (Stress-B), order of the two is unknown), North-South compressive stress and East-West compressive stress (present stress pattern) in turn with time (Shigematsu et al., oral presentation in the 117th Annual Meeting of the Geological Society of Japan, 2010). Among the 327 small brittle faults contributed to stress inversion on Shigematsu et al. (2010), 153 faults could be uniquely attributed to one of the four stress patterns, 127 faults could be attributed to two or three of them, and 47 faults belong to none of them (Tanaka et al., oral presentation in the 118th Annual Meeting of the Geological Society of Japan, 2011).

We tried to infer a suitable stress pattern each of those 127 faults from statistical analysis on the fault material on the slip surfaces of the small brittle faults.

We sampled the fault material on the 129 of those 153 faults, analyzed mineral composition of those samples by X-ray diffraction (XRD) and applied principle component analysis (PCA) to the 129 mineral composition data sets. As a result, an inverse relationship between quartz and carbonate was proved to be most significant characteristic. The fragmented wall rock is dominant in the quartz rich samples; in contrast, alteration minerals are dominant in the carbonate rich ones. Correlating among the four with removal of the carbonate rich ones, the newer (shallower) stress pattern is, the more carbonate content in fault material is. In the fault material, Stress-A contains less carbonate than Stress-B. Thus, the former may be older than the latter.

We sampled the fault material on the 113 of those 127 faults, analyzed mineral composition of those samples by XRD and applied discriminant analysis to the 113 mineral composition data sets. Assuming the stress pattern having minimum Mahalanobis' distance is valid, the 78 faults could be attributed to a suitable stress pattern. Combining the 153 faults and the 78 faults, the faults attributed to Stress-A and Stress-B are densely distributed from 140m to 250m depth, about 300m far from the MTL. In contrast, the faults attributed to the other two stresses are concentrated within 200m from the MTL. Thus, the width of the brittle deformation zone along the MTL might be narrower as regional uplifting and faulting.

Keywords: Median Tectonic Line, fault, borehole core, mineral composition, statistical analysis, stress history

Gneissose tonalites and meta-granitoids in the Hase-Ichinose district of the Ina city, central Japan

ONO, Akira^{1*}

¹None

Hiji tonalite is exposed along the Median Tectonic Line in the Hase-Ichinose district (Figures A and B). The granitoids exhibit considerable variations in chemical composition and rock texture from place to place. Considering gneissose structures and recrystallizations of minerals, the granitoids can be classified into three types: (1) massive, banding and layered tonalites (2) gneissose tonalites (3) meta-granitoids. These are (1) ordinary granitoids, (2) granitoids deformed at a final stage of the solidification of granitic magma and (3) granitoids deformed and metamorphosed under high temperatures after the solidification of granitic magma. Parts of these granitoids were metamorphosed at about 673K and the Kashio mylonites were formed. The Kashio mylonites are mainly distributed in the yellow region of Figure B.

The Hiji tonalite in the Hase-Ichinose district is characterized by the common occurrence of layered and banded granitoids. Banding structures consist of the alternation of felsic and mafic layers which are poor in continuity. Felsic and mafic layers are often folded locally in a hand specimen scale. Folding layered granitoids is also found in several outcrops. Gneissose tonalites are also common. They are often rich in quartz-feldspar veins which are folded complexly. The quartz-feldspar veins may be deformed under the existence of a small amount of granitic magma. The deformation may be continued even after the solidification of the magma. The problem is whether igneous minerals were recrystallized or not. In this regard, strong schistose texture, plastic deformation of minerals and preferred mineral arrangements are uncommon. The recrystallization under the solid-state condition appears to be slight in most cases.

Meta-granitoids in the study area are called as the Gatsuzozan metamorphic rocks [1, 2]. They are characterized by conspicuous deformations and foliations [2]. However, typical meta-granitoids are rare. Moreover, it is not easy to confirm the recrystallization of minerals because of common occurrences of relic igneous minerals. For example, we examined chemical variations of plagioclase grains for a massive tonalite and a meta-tonalite. The results are shown in Figure C. The compositional variation is small for the meta-tonalite relative to that for the tonalite. The conclusion is however obtained by neglecting a calcium-rich large plagioclase grain for the meta-tonalite.

The original rocks of some meta-granitoids are vein-rich gneissose granitoids. The conclusion is based on the study of the chemical variation of plagioclase grains for a meta-granitoid that is rich in quartz-feldspar veins. The examined rock was exposed in an outcrop near Mt. Gatsuzozan. Plagioclase crystals in the veins exhibit conspicuous chemical zonings, and the chemical variations among plagioclase grains are large. On the other hand the recrystallizations of plagioclases are considerable in the matrix. The data suggest that the quartz-feldspar veins were formed before the metamorphism.

The tectonics of the formation of meta-granitoids took place in almost all the study area. Nevertheless, typical meta-granitoids are uncommon. Complexly zoned plagioclases are common for the granitoids in the study area. The poor recrystallization may be due to the following facts. Deformations took place in limited regions. Driving force to the recrystallization was small because the recrystallization occurred slightly below the solidus temperature of granitic magma.

[1] Ono, 2008, Abst. Ann. Meet. Geol. Soc. Japan, p.243.


[2] Ono, 2009, Abst. Ann. Meet. Geol. Soc. Japan, p.259.

Keywords: gneissose granitoid, meta-granitoid, Hiji tonalite, Gatsuzozan metamorphic rocks, recrystallization, Hase-Ichinose

SMP46-P01

Room:Convention Hall

Time:May 25 15:30-17:00



P-T paths of tectonic blocks in the Kamuikotan metamorphic rocks, Etanbetsu-Horokanai district, Hokkaido

OKAMOTO, Ayumi¹, TAKESHITA, Toru^{1*}

¹Hokkaido University

The Kamuikotan metamorphic rocks are known as typical high-P/T type metamorphic rocks formed at convergent plate boundaries at Cretaceous, northern Japan. On these rocks, a number of studies were conducted in 1980s, but since then few studies have been conducted and their tectonics has not been fully interpreted. In the study area, Etanbetsu-Horokanai district, the Kamuikotan rocks are characterized by the paragenesis of lawsonite and glaucophane, and suffered the highest pressure metamorphism (Sakakibara and Ota, 1994) in the Kamuikotan rocks. In this area, while accretionary sediments suffered a high-P/T type metamorphism (prograde rocks), epidote-amphibolite and epidote-garnet amphibolites indicating medium pressure type metamorphism also occur as tectonic blocks, which later suffered the same high-P/T type metamorphism as the sediments did (retrograde rocks, Ishizuka and Imaizumi, 1980; Ishizuka et al., 1983). It is inferred that amphibolite originated from layered gabbro, which is somewhat similar to the Horokanai ophiolite. Further, although blueschist which originated from basalt is classified as prograde rocks by these authors, the amphibolite and blueschist could have been derived from the same oceanic crust, as discussed below. If so, a series of ophiolitic rocks was brought to the subduction channel by tectonic erosion, where the formation of the Kamuikotan rocks proceeded.

We have analyzed mineral assemblages in these rocks (tectonic blocks), and conducted micro-chemical analyses of compositional zoning in amphiboles from these rocks with an EPMA. Among two blueschist samples collected from the Etanbetsu pass, amphiboles from one sample (falling rocks) show a zoning where actinolite core (Al(IV)=0.17-0.33, Na(B)=0.38-0.43) is overgrown by glaucophane rim (Al(IV)=0.003-0.024, Na(B)=1.81-1.86), while ones from the other sample (outcrop rocks) almost uniformly show a composition of winchite. On the other hand, amphiboles in an epidote-amphibolite sample from the Horokanai pass have a zoning where actinolite-actinolitic hornblende core (Al(IV)=0.33-0.71, Na(B)=0.09-0.40) is overgrown by thin glaucophane rim (Al(IV)=0.009-0.079, Na(B)=1.39-1.85). Accordingly, amphiboles in the amphibolite and blueschist have a similar zoning, and thus these rocks could have been derived from the same ophiolitic rocks. For epidote constituting these amphibolite and blueschist, an interesting observation is that some epidote grains were severely fractured often forming microboudins, while the other grains show idiomorphic shape, and not fractured at all. Hence, there may be a possibility that epidote grains were formed at two distinct stages. It is anticipated that the relation between P-T path and deformation stage will be inferred in the future by analyzing pistasite component of epidote.

Keywords: Kamuikotan metamorphic rocks, high-P/T type metamorphism, tectonic blocks, tectonic erosion, compositional zoning in amphibole, pressure-temperature paths

Analyses of deformation stages and paleostress in the Kamuikotan metamorphic rocks, west of the Asahikawa-city, Hokkaido

NAKAYAMA, Takahito^{1*}, TAKESHITA, Toru¹, OKAMOTO, Ayumi¹

¹Hokkaido University

The Kamuikotan belt, distributed in the central Hokkaido, is a representative high-P/T type metamorphic belt that developed on the convergent boundary between the Eurasian Continent and the oceanic plate subducting beneath the edge of the Asian continent in the Cretaceous. In the metamorphic rocks cropping out in the Kamuikotan gorge area in the western part of Asahikawa city, structures such as foliations and folds have been developed well, and many quartz veins intruded. We measured the orientations of these, and found the existence of three deformation stages. Okamoto (2011, graduation thesis) has suggested, from analyses of orientations of foliations. The D1 is defined as the stage at which principal foliations in this area (S1) were formed. S1 was folded to form closed east-vergent folds with axes which trend N-S?NNE-SSW and plunge south at low angles (D2). Furthermore, S1 locally formed open folds with axes that are almost vertical and fold axial surface oriented in E-W. These folds overprinted D2 folds (D3). Poles of quartz veins in the outcrop where D2 folds are dominant were distributed symmetrically with respect to the D2 fold axial surface. This indicates that quartz veins intruded in a stress field that formed D2 folds. We also sampled a quartz vein with dominant D3 crenulation folds that was intruded parallel to foliations, and analyzed microstructures of quartz using three methods explained below to infer the paleostress field. We first define the sample coordinates, the X-Y-Z axes, where X trends E-W and plunge east at 15 degrees, Y trends north and plunges horizontally, and Z trends E-W and plunges east and upward at 75 degrees, respectively. Poles of deformation lamellae in quartz grains are relatively rotated towards sigma 3 direction from the direction of c-axis. Estimation of the paleo-sigma 1 and 3 directions can be possible from the distribution of arrows which connect the c-axis and lamellae pole (arrow head) orientations in grains. Arrows converged in the direction that plunge the -X axis at 20-60 degrees in the XZ plane. Kink bands tend to be formed in grains where the c-axes are oriented between the sigma 1 and 3 axes (i.e. orientations of highest resolved shear stress on slip plane), and not to be formed in grains where the c-axes are oriented to the directions of principal stress axes. Based on these facts, we can estimate the directions of principal stress axes from distributions of c-axes orientations in grains containing and not containing kink bands. The c-axes in grains with kink bands showed a girdle distribution with the angular radius of 45 degrees whose center is Y axis, and those in grains without kink bands were concentrated in the direction that plunges -X direction at 20-30 degrees in the XZ plane. Sigma 3-direction can also be estimated from distribution of poles of healed microcracks formed in quartz grains because they are ideally formed perpendicularly to the sigma 3-axis. The highest concentrated area of poles of microcracks plunges -X direction at 40 degrees in the XZ plane. Combining all these results, we could estimate that the paleostress field during the D3 period is such that the sigma 1, 2, and 3 axes trends N-S with no plunge, vertical, E-W with no plunge, respectively. From the measurement of homogenization temperature of fluid inclusions in quartz grains constituting a vein which probably formed during the D3 period, we obtained the values of temperature of 142-176 degrees. We could estimate that the pressure was 1.6-2.1 kbar when fluid inclusions were trapped, using the slope of isochore of pure water (76 degrees/kbar). This pressure value during D3 period is significantly lower than that estimated from fluid inclusions trapped during D2 period by Okamoto (2011, graduation thesis) (about 2.5 kbar). These facts indicate that an uplift of 0.5-1.0 kbar (1.5-3 km) occurred from the D2 period to the D3 period.

Keywords: Kamuikotan metamorphic rocks, deformation stage analyses, paleostress analyses, deformation microstructures in quartz, healed microcrack, fluid inclusion

Wedge extrusion followed by out-of-sequence thrusting and duplexing, and solving knocker problem, the Sambagawa HP-LT sc

OSOZAWA, Soichi^{1*}

¹Department of Earth Sciences, Graduate School of Science, Tohoku University, ²Department of Earth and Environmental Sciences, California State University, Fresno

The Sambagawa high P/T metamorphic rocks central Shikoku is regionally mapped, and characteristic major structures performed the exhumation of high grade metamorphic rock is clarified. Our geologic map is expected to be an index map available for researchers of the Sambagawa zone. As noted by Osozawa and Pavlis (2007), the most fundamental structure is the extrusional wedge, rebelled D2-1 in this paper. The S directed extruded wedge consists of a series of normal faults domain at hanging wall of the Asemigawa detachment N directed, and a series of thrust faults domain of the footwall S directed. These faults bound every metamorphic mineral zone, and the thermal culmination of the Asemigawa section is oligoclase-biotite zone, hanging wall of the detachment fault. To the W, we now confirm that the culmination contains a metamorphic ultramafic rocks- amphibolite-eclogite complex. To the E, these extrusional wedge and a series of normal and thrust faults are linearly traceable on regional map, but suddenly discontinuous to the footwall chlorite zone. The disruption is due to the D2-2 Hamegano out-of-sequence thrust, newly found in this paper, and the hanging wall consists of the high grade rocks. The OST divides the chlorite zone into the upper L tectonite consisting of varicolored mafic schist, and the other main pelitic schist regionally distributed. D2 folds at the hanging wall is disjunctive and broken by the movement of brittle OST. The OST is traceable to the W from the Asemigawa area. To the N, the Tomisato syncline is observed as mapped by previous studies, but it exists only on hanging wall, and the structure is discordant to the footwall main chlorite zone. Its S limb, hanging wall of the OST, is extrusional wedge, but the NW limb consists of stacked metamorphic mineral zones, and the syncline is asymmetric. Further to the N, OST and hanging wall duplex is involved in the W plunging D2-3 Oboke anticline. We named the D2-2 duplex stack formed near anticlinal axis, the Tomisato duplex. OST and duplex directed S, as the second expression of metamorphic exhumation, following the first wedge extrusion. The Median Tectonic Line might have played an important role for exhumation as a root, and the high grade rocks, including eclogite, is consequently rootless. The Oboke anticline contains the structurally lowest psammitic schist included in the main chlorite zone. The similar psammitic schist unit of the upper different horizon is newly recognized only at the NW limb of the anticline. The psammitic schist and the Oboke thrust is concordantly folded with the OST and duplex at the Oboke anticlinal axis. Another anticline is to the S, the Nakashichiban area, where the psammitic schist of the same horizon is once again exposed. Metamorphosed ultramafic rocks, amphibolite (mafic schist) except for eclogite, calcareous schist, and piemontite siliceous schist are distributed in every metamorphic zone. The rocks are affected by D1 stretching and W directed shear, but never tectonic melange. Therefore, ultramafic rocks were amalgamated with surrounding rocks before D1, and possible mixing mechanism is sedimentary melange. The eclogite might be derived from metamorphic sole of the Jurassic Mikabu ophiolite as the same fashion.

Keywords: Sambagawa HP-LT metamorphic rocks, exhumation, extrusional wedge, out-of-sequence thrust, duplex, eclogite

Microstructural analysis of peridotite xenoliths from Kimberley and Lesotho kimberlite pipes in South Africa

UMEGAKI, Toshiya^{1*}, Masashi Kino¹, MICHIBAYASHI, Katsuyoshi¹, Ikuo Katayama², KOMIYA, Tsuyoshi³

¹Institute of Geosciences, Shizuoka University, ²Department of Earth and Planetary Systems Science, Hiroshima University,

³Department of Earth Science & Astronomy, University of Tokyo

Kimberlite pipes carry peridotite xenoliths derived from the deep upper mantle (70-250 km) beneath Archean cratons. Such xenoliths contain well-developed deformation microstructures. This study presents microstructures of peridotite xenoliths from two kimberlite pipes in South Africa: Kimberley and Lesotho.

The peridotite xenoliths of Kimberley kimberlite pipe consist of lherzolite, harzburgite and dunite. The lherzolites consist of olivine, orthopyroxene, garnet and clinopyroxene. The harzburgites consist of olivine, orthopyroxene, garnet and a minor amount of clinopyroxene. A small amount of phlogopite occurs in both rocks. These rocks are texturally divided into two types: coarse granular (granular type) and extensively sheared peridotites having obvious foliations (foliated type). The granular type peridotites show equigranular textures with slightly curved grain boundaries. The foliated type peridotites show porphyroclastic textures, which consist of coarse-grained porphyroclasts and very fine-grained neoblast matrix. These peridotites display no visible evidence of extensive metasomatism, but show some secondary serpentine alterations along grain boundaries. Major-element compositions of minerals were determined by an electron probe microanalyzer (EPMA). Mineral chemistries for the lherzolites and the harzburgites indicate that these mantle rocks were equilibrated at similar P-T conditions regardless of their texture types. Olivine crystal-preferred orientations (CPOs) were measured in highly polished XZ thin sections using a scanning electron microscope (SEM: Hitachi S-3400N) equipped with electron back-scattered diffraction (EBSD: Oxford-HKL Channel5). The peridotite xenoliths show various olivine CPO patterns.

The peridotite xenoliths of Lesotho kimberlite pipe are grouped into two types: coarse granular type and small-grained elongated type. Olivine CPOs show intense concentrations of [010] axes normal to the foliation, with girdles of [100] and [001] axes within the plane of the foliation (AG type). In particular, the small-grained elongated peridotites show strong concentration of [010] axes. This CPO pattern may result from axial shortening. The coarse granular peridotite shows higher equilibrium temperatures than those for the small-grained elongated peridotites. It suggests that the cratonic mantle beneath Lesotho may contain a domain where the coarse peridotites alternate with the small-grained elongated peridotites at around 150 km deep.

Keywords: Kimberlite, Peridotite xenoliths, microstructure, crystal fabric analysis

Strain localization in the Konoyama mylonite zone, SW Japan

MAEDA, Sumire^{1*}, OKUDAIRA, Takamoto²

¹Faculty of Science, Osaka City University, ²Graduate School of Science, Osaka City University

In a shear zone, grain size of minerals gradually decreases towards the center of the shear zone, and fine-grained minerals developed as a localized high-strain zone at the shear zone center. The difference between grain sizes of minerals in the shear zone center and the shear zone margin resulted from the difference in (1) strain rate, (2) stress, and (3) duration of deformation. Therefore, we should clarify which is the most important factor to localize the strain in the shear zone center. However, in general, duration of deformation cannot be estimated by a conventional geochronological analysis. In shear zone rocks, if metamorphic minerals crystallized during shearing are different in different lithologies, the formation temperature of different lithologies would be different. Thus, strain localized with decreasing temperature with time. In this case, the formation period of them would be different.

In the Ryoke metamorphic belt, there are many ductile shear zones with different scale. The Konoyama mylonite zone (e.g., Takagi et al., 1988; Ishii et al., 2007) is one of the shear zones, and developed around Mt. Konoyama, Kishiwada, SW Japan. In the area, deformed granodioritic rocks from protomylonite, through mylonite, to ultramylonite can be observed. In this study, to clarify whether the formation temperature of deformed rocks is different in different lithologies, we analyzed mineral assemblages and mineral chemistry for deformed rocks with different degree of strain magnitude.

We found some interesting observations as follows: (1) There is no chlorite and muscovite in protomylonites and mylonites, but present in ultramylonites; (2) The recrystallized biotite in ultramylonites shows lower Ti and higher Al than those in protomylonites and mylonites, although the difference is small. These observations suggest that the formation temperature of ultramylonites is slightly lower than those of protomylonites and mylonites, and the formation period of ultramylonites and protomylonites/mylonites is different. This implies that strain localization in the Konoyama mylonite zone occurred retrogressively.

Ref: Takagi, H., Mizutani, T. and Hirooka, K. (1988) *Jour. Geol. Soc. Japan*, 94, 869-886; Ishii, K., Kanagawa, K., Shigematsu, N. and Okudaira, T. (2007) *Jour. Struct. Geol.*, 29, 1083-1098

Keywords: mylonite, shear zone, biotite

Petrology and geochronology of metamorphic rocks in the Bayankhongor area, the central part of Mongolia.

ADACHI, Tatsuro^{1*}, OSANAI, Yasuhito¹, NAKANO, Nobuhiko¹, OWADA, Masaaki², MADHUSOODHAN, Satish-Kumar³, SEREENEN, Jargalan⁴, Chimedtseye Boldbaatar⁵, Kazuhiro Yonemura¹, Aya Yoshimoto¹

¹Kyushu University, ²Yamaguchi University, ³Shizuoka University, ⁴Mongolian Univ. Sci. Tech, ⁵Miner. Res. Auth. Mongolia

Mongolia is situated in the Central Asian Orogenic Belt (e.g. Mossakovsky et al., 1993), which extends from the Siberian craton, Baltica craton, North China craton and Tarim craton. It is composed of subduction-accretion complexes and developed from c. 1000 Ma to c. 250 Ma (Windley et al., 2007). Metamorphic rocks are partly distributed in the central and western part of Mongolia. Metamorphic history would be a key for understanding the tectonic evolution of this area, however detailed study on metamorphic rocks has not been well-documented, including geochronology. In this study, we report petrology and U-Pb monazite and zircon ages on metamorphic rocks occurring in the Bayankhongor area.

The Bayankhongor area is situated between Gobi-Altai range and Hangay range in the central part of Mongolia. The area is mainly composed of granitic gneisses (biotite gneiss and clinopyroxene-biotite gneiss), amphibolites and pelitic gneisses. Pelitic gneisses are classified into garnet-biotite gneiss, garnet-sillimanite-biotite gneiss and garnet -cordierite-biotite gneiss. Garnet in garnet-cordierite-biotite gneiss is replaced by cordierite and biotite, indicating decompression process. Sillimanite in garnet-sillimanite-biotite gneiss is replaced by andalusite, which also indicates later stage metamorphism under the lower pressure condition. Amphibolites are classified into garnet-amphibolite, clinopyroxene-amphibolite and amphibolite. Garnet in garnet-amphibolite is replaced by plagioclase and hornblende, which qualitatively imply decompression as well as pelitic gneisses. Thus, metamorphic rocks in the Bayankhongor area widely recorded amphibolite- to upper amphibolite-facies metamorphism with subsequent decompression.

In the presentation, we show additional data including mineral chemistry, P-T estimations and U-Th-Pb in situ dating and discuss metamorphic evolution of the Bayankhongor area. Based on the interpretation of those data, we suggest its implication for the tectonic process related to the development of the Asia continent.

References

- Mossakovsky et al. (1993) Central Asian fold belt: geodynamic evolution and history of formation. *Geotectonics*, 6, 3-33.
- Nakano et al. (2010) Metamorphic rocks from the Mongol-Altai: A first report of Japan-Mongolia Joint Geological Research (2009-2013). *Journal of Geological Society of Japan*, XI-XII.
- Windley et al. (2007) Tectonic models for accretion of the Central Asian Orogenic Belt. *Journal of the Geological Society*, London, 164, 31-47.

Keywords: metamorphic rock, P-T condition, U-Pb dating, Bayankhongor Area, Mongolia

A primary and secondarily-modified microstructure of kelyphites in garnet pyroxenites from the Ronda peridotite, Spain

OBATA, Masaaki^{1*}

¹Kyoto University

Kelyphite as a breakdown product of garnet may contain in its internal microstructure thermal and recrystallization history that rocks had followed because of its relatively high reactivities coming from its fine-grained nature. I re-investigated a once-well studied and published complexly-zoned kelyphite in a garnet pyroxenite layer in the Ronda peridotite, Spain (Obata, 1994). The zoning has been reported to consist of Zone I (Opx+Pl+Sp) adjacent to garnet, Zone II (Ol+Pl+Sp) and Zone III (Ol+Pl) adjacent to primary aluminous Cpx (Obata, 1994). A steady state-reaction and diffusion model has been proposed to account for a simultaneous generation of such zoned structure by a reaction between garnet and primary Cpx (Obata, 1994) in the plagioclase-lherzolite facies condition. However more simple structured kelyphite that consists of Opx+Pl+Sp+Cpx occurs in other pyroxenite layers from the same body. An important finding in sample R127 is the occurrence of trace amount of Cpx in both Zones I and II, thereby casting a doubt to the modeling that assumed a single-stage metasomatic reaction in the plagioclase-lherzolite facies as proposed in Obata (1994). Through more careful comparative studies of these samples, the author came to a conclusion that the simple structured kelyphite is a primary one and the complex zoning that Obata described is a secondarily modified one through later fluid-aided metasomatic reactions. The zone II and III are reaction products developed between a reactant pair of Zone I and the primary Cpx. From the presence of a small amount of Cpx in Zone I, It is concluded that first kelyphitization forming Zone I took place in the seilad subfacies of the spinel-lherzolite facies (OHara, 1967) and that late modification producing Zone II and III took place in the plagioclase-lherzolite facies. It is considered from the fact that these reactions are metasomatic, some metamorphic fluids acted as agents of the reactions. Kelyphitization of garnet occurred also in the host garnet peridotites. It is therefore possible to explore the fluid activities through comparative studies of kelyphitization between different lithofacies with close spatial associations.

Keywords: kelyphite, garnet, reaction zone, fluid, Ronda peridotite

Fault kinematics along the Itoigawa - Shizuoka Tectonic Line in western Yamanashi

KAZATO, Yoshihito¹, KANAGAWA, Kyuichi^{1*}

¹Department of Earth Sciences, Chiba University

Fault kinematics of the Itoigawa - Shizuoka Tectonic Line (ISTL) in western Yamanashi is discussed based on structural observations and analyses of fault rocks.

In the eastern foothill area of Mt. Ho-oh, the ISTL forms the boundary between the Kaikoma-Ho-oh granite with an intrusion age of ca 15 Ma on the west and the sediments of the Momonoki Subgroup in the Miocene Koma Group on the east. It strikes NNW in the northern and southern subareas, but NNE in the central subarea, and dips 45-60 degrees toward W. The microstructures of the mylonitized granite near the ISTL indicate reverse faulting toward SE during its mylonitization. Based on the temperatures during its mylonitization inferred from quartz microstructures and c-axis fabrics and the cooling history of the granite pluton inferred from K-Ar ages (Sato et al., 1989), this reverse faulting likely occurred at 12-13 Ma. Cataclasites along the ISTL have structures indicative of sinistral faulting, while gouges along the ISTL show structures indicative of reverse faulting.

In the Hayakawa River area located ca 20 km southward away from the Mt. Ho-oh area, the ISTL forms the boundary between the slates of the Eocene to Miocene Setogawa Group on the west and the volcanoclastic rocks of the Kushigatayama Subgroup in the Miocene Koma Group on the east. It strikes NNW and steeply dips toward W. Asymmetric structures in the Setogawa slates near the ISTL indicate reverse faulting toward ESE during cleavage development at ca 15 Ma and the subsequent sinistral faulting after the cleavage development. Based on the temperatures during these two stages of faulting inferred from vein quartz microstructures and c-axis fabrics and the cooling history of the Setogawa slates inferred from fission-track ages of detrital zircon (Yamagiwa et al., 1997), the reverse faulting likely changed to the sinistral faulting at ca 13 Ma. Cataclasites along the ISTL have structures mostly indicative of sinistral faulting, while gouges along the ISTL show structures mostly indicative of reverse faulting. K-Ar ages of gouges (Tanaka et al., 1997) suggest that this reverse faulting occurred after 7 Ma.

In summary, three stages of faulting along the ISTL are recognized in both areas in western Yamanashi; initial reverse faulting associated with granite mylonitization and slate cleavage development, secondary sinistral faulting associated with the formation of cataclasites, and final reverse faulting associated with the formation of gouges. These three stages of faulting occurred in this sequence, and accordingly the pressure and temperature during faulting decreased. However, the ages described above suggest that the change from the initial reverse faulting to the secondary sinistral faulting occurred ca 1 Ma earlier in the Hayakawa River area than in the Mt. Ho-oh area. This is likely because the collision of the Izu arc with the Hoshu arc started from the south and propagated northward with time.

Keywords: Itoigawa - Shizuoka Tectonic Line, western Yamanashi, fault kinematics

Two burial-exhumation cycles of tectonic blocks from the northern Montague fault zone, Guatemala

ENDO, Shunsuke^{1*}, Simon R. Wallis¹, Luigi A. Solari²

¹Nagoya University, ²Universidad Nacional Autonoma de Mexico

The Motagua fault zone (MFZ) of central Guatemala is a sinistral suture between the Maya Block (North American Plate) and the Chortis Block (Caribbean Plate). High-P tectonic blocks occur in serpentinite-matrix melanges exposed immediately north and south of the MFZ. The northern MFZ (NMFZ) melange hosts warm-subduction related blocks of epidote eclogite and garnet amphibolite, whereas the southern MFZ (SMFZ) melange includes cool-subduction related blocks of lawsonite eclogite and blueschist. Both the NMFZ and SMFZ eclogites give indistinguishable Sm-Nd garnet-omphacite isochron ages of around 140-130 Ma (Brueckner et al. 2009), but K-Ar phengite ages of the NMFZ high-P blocks (77-65 Ma) are significantly younger than those of the SMFZ high-P blocks (125-116 Ma) (Harlow et al. 2004). Further petrological study especially on the NMFZ high-P blocks is needed to interpret the available age data, and hence to understand the tectonic evolution of this region. We present detailed petrological data of garnet amphibolite from the NMFZ melange exposed at Estancia de La Virgen.

Single garnet crystals in the studied sample consist of four distinct growth zones (Grt1, Grt2, Grt3 and Grt4) and grossular inclusions. Compositions of the four garnet generations are: Grt1 (Alm₄₄₋₅₂ Sp₈₅₋₁₃ Prp₅₋₆ Grs₂₈₋₄₁), Grt2 (Alm₃₇₋₄₃ Sp₈₈₋₁₆ Prp₁₋₂ Grs₄₁₋₅₁), Grt3 (Alm₄₈₋₅₃ Sp₈₃₋₄ Prp₃₋₆ Grs₃₈₋₄₁) and Grt4 (Alm₅₀₋₅₈ Sp₈₂₋₃ Prp₈₋₁₀ Grs₃₀₋₃₈). Grt1 occurs as highly corroded cores of the garnet crystals, and contains inclusions of epidote, omphacite (Jd₂₈), titanite and actinolite, indicating the first subduction-related metamorphism in the eclogite facies (M1). The second growth zone (Grt2) hosts inclusions of chlorite, hornblende/actinolite, albite/oligoclase, allanite/epidote, zoisite, K-feldspar, indicating relatively low-P/T metamorphism in the transitional greenschist/amphibolite facies (M2). Grossular (Grs₇₂₋₈₆ Adr₆₋₂₂ Sp₈₁₋₇) is exclusively present as discrete crystals included in an intermediate zone within Grt2. The third garnet generation (Grt3) is recognized as a skeletal/dendritic overgrowth on Grt2. Grt3 hosts abundant aqueous fluid inclusions in addition to an almost identical inclusion paragenesis as Grt2. Metamorphism associated with Grt3 growth (M3) could be related to a pulse-like hydrothermal event. The last garnet generation (Grt4) resulted from second subduction-related metamorphism (M4). The early phase of M4 took place in the garnet-unstable conditions, and is recorded as relic inclusions within matrix amphibole. This early M4 assemblage includes glaucophane, actinolite, pumpellyite, phengite (Si=3.58-3.70 apfu) and paragonite, indicating high-P/T type metamorphism in the transitional pumpellyite-actinolite/blueschist facies. The inclusion paragenesis within Grt4 is identical with the matrix assemblage that consists of edenite/barroisitic hornblende, albite/oligoclase, epidote, phengite (Si=3.38-3.45 apfu), chlorite, titanite and quartz. The matrix amphibole was slightly replaced first by glaucophane + actinolite and then by actinolitic hornblende during final exhumation. P-T conditions during Grt4 growth are estimated to be ~550 deg.C, 1.1 GPa, and thus the K-Ar system of phengite have been closed after the thermal peak of M4.

By combining the available information, we presume that the NMFZ high-P blocks underwent, at least, two subduction-exhumation cycles with three main tectono-metamorphic phases: 1) the first subduction-related metamorphism (~140-130 Ma), 2) subsequent thermal events at shallow crustal levels, and 3) the second subduction-related metamorphism (~77 Ma) at the transitional oceanic to continental subduction setting.

References: Brueckner et al. (2009), *EPSL* 284, 228-235; Harlow et al. (2004), *Geology* 32, 17-20.

Keywords: exhumation, polygenetic garnet, Guatemala, subduction

Geologic structure of the Sanbagawa Metamorphic Rocks in western Shikoku and deformation pattern of viscous fluids

MIYAZAKI, Kazuhiro^{1*}

¹GSJ AIST

The Sanbagawa Metamorphic Rocks are high-pressure metamorphic rocks formed at depth deeper than lower crust. During exhumation, it is expected that the metamorphic rocks should suffered large-scale deformation. I created geological map of the Sanbagawa Metamorphic Rocks in the Iyo-Nagahama district, western Shikoku, and compared the geologic structure and patterns of viscous fluid under simple shear deformation.

The Sanbagawa Metamorphic Rocks in the Iyo-Nagahama district can be divided into Uchiko and Iyo units 1). The Uchiko unit is mainly composed of pelitic schist, and the Iyo unit is mainly composed of mafic schist. The Iyo unit can be also divided into lower, middle and upper subunits. The lower subunit is composed of mafic, siliceous, pelitic and psammitic schists. The middle subunit is mainly composed of thick mafic schist. The upper subunit is composed mainly of mafic and pelitic schists. The lower and middle subunits suffered metamorphism equivalent to that of chlorite zone in central Shikoku. In both subunits, mineral assemblage of epidote + glaucophane + albite + chlorite + quartz occur in mafic schist, and no garnets appear in pelitic schist. In the upper subunit, albite porphyroblast occur in pelitic and mafic schists, and garnet appears in pelitic schist. Therefore, metamorphic grade in the Iyo unit increases toward upper structural levels. Phengite K-Ar ages obtained by this study and previous study 1) yield 78.7 ± 2.0 - 88.1 ± 2.2 Ma for the Iyo unit, and 90.6 ± 2.3 - 94.6 ± 2.4 Ma for the Uchiko unit.

An E-W trending antiform develops in the Iyo unit at central part of the district, and the schistosity dips gently to north at northern side of antiform, and dips gently to south at southern side of the antiform. Recumbent fold with a few km wavelengths is inferred by detailed mapping of red-colored siliceous schist of the middle subunit of the Iyo unit in western part of the district. Folds with sub-horizontal axial plane also develop at outcrop scales in the Iyo unit.

Pressure and temperature in high P/T metamorphic rocks, such as the Sanbagawa Metamorphic Rocks, drastically increase in perpendicular to schistosity. Formation of this type of P-T structure may require tectonic displacement under simple shear. I examined structural evolution of viscous fluids with different viscosity under simple shear deformation. The system consists of high viscosity fluid (10^{23} Pa s) and low viscosity fluid (10^{21} Pa s). Both fluids are sandwiched by solid plates. Layering structures slightly oblique to shear direction are assumed as initial configuration. Shear deformation were induced by two plate moving to the opposite direction with 1 to 10 cm/y. Results show that folds with a few cm to few km wavelengths are formed during evolution of patten under simple shear deformation. Although the comparison of the results with natural observation is insufficient, the results suggest that tectonic displacement of metamorphic rocks formed at different P-T and formation of large-scale folds may take place simultaneously under simple shear deformation.

1) Banno, Y., et al. (2010) Geology of the Ozu District. Quadrangle Series, 1:50,000, Geological Survey of Japan, AIST, 58p.

Keywords: Sanbagawa Metamorphic Rocks, Geologic structure, viscous fluids, metamorphsim, deformation, fold

Protolith and tectonic environment of blueschist from the Kurosegawa tectonic zone, South-west Japan.

YOSHIMOTO, Aya^{1*}, OSANAI, Yasuhito¹, NAKANO, Nobuhiko¹, ADACHI, Tatsuro¹, YONEMURA, Kazuhiro¹, Hideo Ishizuka²

¹Kyushu University, ²Kochi University

Blueschists from the Kurosegawa tectonic zone are distributed from central Kyushu to Kanto mountains. These rocks are usually embedded in the serpentinite melange. Blueschist is layered and can be divided into two layers; blue and green layers. Coexisting minerals of the blue layer are crossite-glaucophane, lawsonite, and less amount of sodic pyroxene. Secondary minerals that can be found in the blue layer are epidote, pumpellyite, actinolite and chlorite. Mineral assemblage of the green layer is similar to that in the blue layer. However, chlorite and actinolite of the green layer are more abundant than that of the blue layer, which might be due to hydration during retrograde metamorphism.

It is also noted that the green layer is enriched in LIL elements in comparison with the blue layer affected by the retrograde metamorphism, and then we used the data only from the blue layer for later discussion.

Based on several discrimination diagrams using HFS and REE elements of the blueschists, their protoliths can be divided into following four types of basalt: (1) N-MORB type, (2) E-MORB type, (3) OIB type, (4) T-MORB type.

Rb-Sr isotopic analysis (using blueschist samples from Kyushu, Shikoku, and Kii peninsula) yields a whole rock isochron age of 269 +/- 8Ma (SrI=0.70513 +/- 0.00018). Zircon U-Pb ages were also determined to be an inherited age (430-490Ma) of pelitic schist from Itsuki area in Kyushu and as a protolith age (480-520Ma) of gabbro from Engyoji area in Shikoku. Therefore, the Rb-Sr isochron age of 269Ma might reflect the timing of the blueschist-facies metamorphism from the Kurosegawa tectonic zone.

Results of these studies suggests that the formation process of Kurosegawa tectonic zone was possibly prior to continental collision of the North China craton and South China craton during the age of early Triassic (ca. 230-220Ma).

In addition, Nd-Sm isotopic analysis (using blueschist samples from Itsuki area in Kyushu) yields a whole rock isochron age of 804 +/- 64Ma (NdI=0.511781 +/- 0.00073). Therefore, blueschists in Kurosegawa tectonic zone were formed by subduction of the oceanic plate that was located in front of the South China craton.

Keywords: Kurosegawa tectonic zone, blueschist

An Outline of High-Pressure Metamorphic Rocks from Bantimala and Barru Complex in South Sulawesi, Indonesia

SETIAWAN, Nugroho^{1*}, OSANAI, Yasuhito¹, NAKANO, Nobuhiko¹, ADACHI, Tatsuro¹, YONEMURA, Kazuhiro¹, YOSHIMOTO, Aya¹

¹Graduate School of Social and Cultural Studies, Kyushu University

Accretionary and metamorphic complexes regarded as Cretaceous subduction complexes of Indonesia are distributed in Central Indonesia include of West and Central Java, South Kalimantan and South Sulawesi. The widely dispersed rocks of the pre-Tertiary basement in the Indonesian region comprise variably metamorphosed accretionary complexes, imbricated terranes, melange, turbidite and broken formations, and ophiolite (Parkinson et al., 1998). These accretionary and metamorphic complexes in South Sulawesi are recorded on the restricted area namely Bantimala and Barru Complex.

Bantimala Complex is a tectonic assemblage of slices and blocks consisting of sandstone, shale, conglomerate, chert, siliceous shale, basalt, ultramafic rocks, schist and schist breccia with the ages of components range from Jurassic to middle Cretaceous (Wakita et al., 1996). The metamorphic rocks are intercalated with melange deposits and mainly consist of high-pressure metabasites and low-pressure metamorphosed clastic sequence rocks. The melange includes clasts and blocks of chert, sandstone, basalt, limestone and schist embedded within a sheared shale matrix (Wakita et al., 1996). K-Ar ages of phengite on garnet-glaucophane rock yields 125-139 Ma and 107-119 Ma, 118-130 Ma on mica rich intercalated with garnet-glaucophane rock, 108-120 Ma and 109-121 Ma for mica-quartz schist intercalated with hematite bearing glaucophane schist (Wakita et al., 1994, 1996) and 134-140 Ma for eclogite (Parkinson et al., 1998). Miyazaki et al. (1996) estimate the peak pressure of the Bantimala eclogites were 18-24 kbar at 580-640 °C. Parkinson et al. (1998) suggest the pressure and temperature for eclogites and garnet-glaucophane rocks were 18-24 kbar at 580-620 °C and jadeite-garnet-quartz (coesite) rock were >27 kbar at 720-760 °C.

High-pressure metamorphic rocks in the Bantimala Complex mainly crop out as river boulders. Mafic rocks (eclogite) are the most common comparing to pelitic lithologies such as glaucophane schist and garnet-glaucophane-phengite schist. Rarely garnet-jadeite-quartz rock crops out as 3 m wide river boulders in Bantimala River. Meanwhile, the Barru Complex is located approximately 30 km north of the Bantimala area. Metamorphic rocks in this area are crop out more restricted area with most common lithologies are variably of garnetiferous quartz-mica schist and serpentized peridotite. Wakita, et al. (1994) reported phengite K-Ar age determination from quartz-mica schist of 106 Ma.

The evidence from mineral chemistry analysis, suggests that the Grt-Jd-Qtz rock passed through Jd + Qz stability field during metamorphic evolution. Peak metamorphism of eclogite is represented by garnet rim and omphacite in the matrix. Grt?Cpx thermometer (Krogh, 1988., Pattison & Newton, 1989., and Ravna, 2000) yield a temperature of 720-820 °C for pressure of 25 kbar on the sample of eclogite 110310T03F. The occurrence of glaucophane inclusions in the core and mantle of garnet indicate the former stability field in the blueschist area. Retrogression observed in the eclogites is represented by replacement of omphacite by actinolite-tremolite-winchite-barroisite amphiboles.

Moreover on the retrogression stage, three types of eclogite are recognized. All of the types are contain garnet and omphacite bearing assemblages. The difference between each type is secondary phase minerals that present during retrogression. Type I of eclogite is dark green eclogite with rich of omphacite in the matrix and less of amphibole or minimally affected by the secondary hydration. Type II of eclogite is bluish colored rich of glaucophane and less of omphacite in the matrix. Omphacite grains in this eclogite are restricted as inclusions on garnet or other minerals. Type III of eclogite is light green-bluish colored dominated by mostly amphiboles in the matrix that ranges in composition from actinolite to barroisite. In these rocks, omphacite is partially to completely replaced by amphiboles.

Keywords: high-pressure metamorphic rocks, eclogite, Bantimala Complex, Barru Complex, South Sulawesi, Indonesia

Metamorphic conditions of blueschists and greenschists in the Toudaoqiao area, in Inner Mongolia, NE China

ZHAO, Limin^{1*}, TAKASU, Akira¹, LI, Weimin², Liu Yongjiang²

¹Department of Geoscience, Shimane University, Japan, ²Jilin University, P.R.China

Blueschists and related metamorphic rocks are exposed in the Toudaoqiao area of the Honghuaerji-Yimin district in Inner Mongolia, northeastern China (Ye et al., 1994). These metamorphic rocks occur along Tayuan-Xiguitu fault, located between the Ergun block and the Xing'an block, and those are situated in the eastern parts of the Central Asian Orogenic Belt (Sengor et al., 1993).

Zhao et al. (2011) reported that blueschists occur together with greenschists, pelitic schists and siliceous schists in this area. The blueschist consists of amphibole (Gln, Mg-rbk and Win), epidote, albite, phengite (Si=6.7-7.0 p.f.u.), quartz, titanite and hematite, and the metamorphic conditions are in the epidote-glaucophane facies (380-450°C, 11-14kbar).

In this study we describe barroisite-bearing blueschists and greenschists. The barroisite-bearing blueschists consist mainly of amphibole (Mg-rbk, Gln, Win, Ktp and Brs), epidote, albite, phengite (Si=6.7-7.2 p.f.u.), chlorite, quartz, calcite and titanite with minor amounts of hematite. The amphibole is of subhedral prismatic crystal with size up to 0.2 mm long. It contains inclusions of amphibole (Ktp and Brs) as a relic of the precursor metamorphic event. The amphibole is zoned with winchite core ($Na_B=0.54-1.48$ p.f.u.) to magnesioriebeckite rim ($Na_B=1.52-1.81$ p.f.u.), and it is partly replaced by chlorite, calcite, titanite and quartz. The zoning texture suggests the prograde stage is in the winchite stability field followed by the magnesioriebeckite/glaucophane stability field. The peak metamorphic stage is characterized by the rim of amphibole (Mg-rbk/Gln) and coexisting minerals (epidote+albite (An<0.38)+phengite (Si=7.2 p.f.u.)+hematite+titanite+calcite), suggesting the epidote-glaucophane schist facies metamorphic event. The retrograde stage is represent of chlorite, calcite, quartz and titanite, which replace the amphibole, probably suggests the greenschist metamorphic event. The peak stage mineral assemblage is same as the previously described blueschist, and the metamorphic condition are T=390-450°C, P=13-15kbar.

The greenschists consist mainly of amphibole (Act, Mg-hbl, Ts, Ed and Hs), epidote, chlorite, albite, phengite (Si=6.7-6.9 p.f.u.), quartz, calcite and titanite with minor amounts of hematite. The amphibole is mainly classified as actinolite ($Na_B<0.34$ p.f.u.), and it is of euhedral to subhedral prismatic crystal, with size up to 0.5 mm long. It is commonly zoned with magnesiohornblende (Ti=0.14-0.26 p.f.u.) core and actinolite ($Na_B<0.16$ p.f.u.) rim, and its core contains inclusions of amphibole (Ts, Ed, Hs and Mg-hbl (Ti<0.10 p.f.u.)) and titanate. Actinolite ($Na_B=0.10-0.34$ p.f.u.) and chlorite occur along cleavages of the zoned amphibole. The core of the amphibole and its inclusions (Ts, Ed, Hs and Mg-hbl) as relic of the precursor high temperature metamorphic event. The peak metamorphism is represented by actinolite+chlorite+epidote+albite (An<0.2)+phengite (Si=6.9 p.f.u.)+quartz+hematite, suggesting low-temperature and high-pressure metamorphic condition.

Reference

Blundy and Holland, 1994, *Miner and Petrol.*, 116, 433-447; Evans, 1990, *Lithos*, 25, 3-23; Holland, T.J.B., 1983, *Contr. Miner. Petr.*, 116, 433-447; Massonne and Schreyer, 1987, *Contr. Miner. Petr.*, 96, 212-224; Otsuki and Banno, 1990, *J. Metamorphic Geol.*, 8, 425-439; Sengor et al., 1993, *Nature*, 364, 299-307; Ye et al., 1994, *M-SGT Geology Issue*, 73-83; Zhao et al., 2011, *JPEU Conf.*

Keywords: high P/T metamorphism, blueschist, greenschist, Tayuan-Xiguitu fault, NE China

Geochemistry of the HP-UHP metamorphic rocks, Makbal complex, northern Kyrgyz Tien-Shan

BAKIROV, Azamat¹, Kadyrbek Sakiev¹, TAGIRI, Michio², Apas Bakirov¹, TAKASU, Akira^{3*}, OROZBAEV, Rustam⁴, Asel Togonbaeva¹, Jun-Ichi Kimura⁵

¹Institute of Geology, Kyrgyz National Academy of Science, ²Hitachi City Museum, Miyatacho, ³Department of Geoscience, Shimane University, ⁴Department of Geology and Mineralogy, Kyoto University, ⁵Institute for Research on Earth Evolution, JAMSTEC

Geochemistry of the HP-UHP metamorphic rocks, Makbal complex, northern Kyrgyz Tien-Shan

A.A. Bakirov¹, K.S. Sakiev¹, M. Tagiri², A.B. Bakirov¹, A. Takasu³, R.T. Orozbaev⁴, A.A. Togonbaeva¹ and J.-I. Kimura⁵

The Makbal UHP complex occurs in the western part of Kyrgyz Range of the Northern Tien-Shan and consists of quartzite and muscovite-quartzite, garnet-muscovite-quartz and chlorite-muscovite-quartz schists, marbles, amphibolites with relicts of eclogites and UHP talc-garnet-chloritoid schists (Tlc-Grt-Cld). The coesite has been found in quartzites and Tlc-Grt-Cld schists, and they experienced UHP metamorphism ($P > 29$ kbar and $T < 600$ °C), whereas the eclogites and garnet amphibolites are experienced mainly HP metamorphism with different peak P-T conditions (Tagiri et al., 2010). UHP Tlc-Grt-Cld schists are conformably intercalated in thick quartzite layers. Eclogites occur as imbricated lenses within these host UHPM rocks.

A phengite K-Ar age of 509±13 Ma (Tagiri et al., 2010), CHIME monazite age of 481±26 Ma (Togonbaeva et al., 2009) and 498±7 Ma for the UHP Tlc-Grt-Cld schist and paragonite K-Ar age of 482±17 Ma (Tagiri et al., 1995) and zircon SHRIMP U-Pb ages 509±7 Ma and 502±10 Ma (Konopelko et al., 2011) has been reported for the eclogite. The granitic rocks discordantly intruded in the Makbal complex are dated as 399±10 Ma (Tagiri et al. 2010).

We carried out whole-rock geochemistry analyses on major, trace and REE abundances for the eclogites, Qtz-Grt-carbonate rocks, Tlc-Grt-Cld schists, quartzites, garnetite and one clinopyroxene-biotite rich rocks (lamproite?) from the Makbal Formation. The classification diagrams, using major and trace elements, classify the eclogites as tholeiitic basalts. Tlc-Grt-Cld schists, garnetite and Qtz-Grt-Carb rocks plot in the same field. In the conventional tectonic discrimination diagrams, the eclogites plot in the field of MORB and VAB. Similarly, the Tlc-Grt-Cld schists and Qtz-Grt-Carb rocks show also same compositions in these diagrams as eclogites. However, Tlc-Grt-Cld schist have very low content of CaO (<2 wt%), Na₂O (<0.1 wt %) and relatively high MgO (<18 wt %), which is not comparable with basaltic composition. Togonbaeva et al. (2010) pointed out the similarities of the eclogite and Tlc-Grt-Cld schists compositions, and they suggested that the Tlc-Grt-Cld schist is the mixture between eclogite protoliths (basaltic) and quartzites (pelitic) compositions.

Chondrite and primitive mantle normalized diagrams of REE and some trace element abundances show relatively similar pattern for the eclogites, Qtz-Grt-Carb rocks and Tlc-Grt-Cld schists, except some REE enrichments in Tlc-Grt-Cld schists. Eclogites and Tlc-Grt-Cld schists are depleted in incompatible elements (Rb, Ba, Sr, K, Ca), however enriched in Pb.

Here, we propose the possible metasomatic origin of Tlc-Grt-Cld schist in the Makbal complex, similarly to whiteschists formation in Dora-Maira Massif (DMM), Western Alps. Two main genetic hypotheses were proposed for the origin of whiteschists in DMM, i.e. sedimentary (highly Mg-meta-evaporite; Schreyer, 1977) vs. metasomatic (Mg-rich fluid into shear zones; Compagnoni & Hirajima, 2001; Schertl & Schreyer, 2008). We suggest that the origin of Tlc-Grt-Cld schist could be related to Mg-metasomatism on the basaltic protoliths of eclogites, at least prior to peak UHP conditions. This process can be responsible for the modification of some major and trace elements in the eclogites to more Mg-rich and Ca- and Na-poor, similarly to whiteschist in DMM. The metasomatic origin supported also by field relations of these rocks and their similar geochronological data.

References: Tagiri et al. (1995), Island Arc; Tagiri et al. (2010), JMPS; Togonbaeva et al. (2009), JMPS, Togonbaeva et al. (2010) JAMS Annual Meeting, Matsue; Compagnoni & Hirajima (2001), Lithos; Konopelko et al. (2011), Gondwana Research; Schreyer (1977), Tectonophysics; Schertl & Schreyer (2008), Eur. Jour. Miner.

Keywords: geochemistry, UHP metamorphism, metasomatism, Makbal, Tien-Shan, Kyrgyzstan

Grain-boundary diffusion coefficient based on variation of quartz grain sizes in metacherts around a contact aureole

OKUDAIRA, Takamoto^{1*}, Hikaru Bando¹

¹Dept. Geosci., Osaka City Univ.

Grain-boundary diffusion coefficient is the most important parameter for considering metamorphic and deformation processes. Grain-boundary (or bulk) diffusion coefficients have been estimated experimentally (Farver and Yund, 1990, 2000) or naturally (Joesten, 1983). According to Farver and Yund (2000), the temperature dependence of silicon (bulk) diffusion in the novaculite is described by the Arrhenius parameters: $D_0 = 3.7 \times 10^{-10}$ m²/s and $Q = 137 \pm 18$ kJ/mol, and $D_0 = 6.2 \times 10^{-9}$ m²/s and $Q = 178 \pm 38$ kJ/mol for the hydrothermal and dry experiments, respectively. On the other hand, variation of the quartz grain sizes in nodular chert in the Christmas Mountains contact aureole was matched by a normal grain growth model in which the temperature dependence of the Arrhenius function along temperature-time history calculated by an one-dimensional thermal modeling. Assuming the grain boundary width of 1 nm, the data permit an estimate of the coefficient for the grain-boundary diffusion of oxygen in quartz as $D = 8.07 \times 10^{-10} \exp(-210 \times 10^3/RT)$. Therefore, there is significant difference between the coefficients for a quartz aggregate estimated experimentally and naturally.

In this study, we evaluate the grain-boundary diffusion coefficient by the same scheme of Joesten (1983). We collected metachert samples from the contact aureole around the Hanase-Bessho quartz diorite (Kiji et al., 2000) at the Hanase Pass, Kyoto. Photomicrographs are taken under optical microscope and scanning electron microscope. The grain boundaries are traced using drawing software (Canvas), and grain sizes as equivalent diameter of circles are estimated by ImageJ. The quartz grain sizes vary systematically with the distance from the quartz diorite. We calculated temperature-time history using a one-dimensional thermal model. Based on Arrhenius plot of the observed quartz grain sizes and calculated temperatures, we estimated the activation energy 140 ± 16 kJ/mol that is comparable with the value for silicon bulk diffusion estimated by Farver and Yund (2000) for hydrothermal experiment. Furthermore, using the value estimated here, we obtained the value of $D_0 = 5 \times 10^{-12}$ m²/s.

References: Kiji, M., Ozawa, H. and Murata, M. (2000) Cretaceous adakitic Tamba granitoids in northern Kyoto, San'yo belt, Southwest Japan. *Jour. Mineral. Petrol. Sci.*, 29, 136-149.

Keywords: grain-boundary diffusion coefficient, quartz, normal grain growth, contact aureole, thermal model, grain size

Unstable boundary formation behind the growth front in sealing of open crack

TORIUMI, mitsuhiro^{1*}, Mayuko Fukuyama²

¹JAMSTEC, ²Akita University

Unstable boundary formation behind the growth front in sealing of open cracks

Mitsuhiro Toriumi * and Mayuko Fukuyama**

* IFREE, JAMSTEC

** IES, Academia Sinica/ Univ Akita

Plate boundary quartzite metamorphic rocks contain abundant sealed open cracks and shear cracks that are occupied mainly by quartz with minor amount of albite and chlorite. Apparently the sealed cracks can be classified into two types: one is the comb type and another is the granular type. The latter displays rapid increase of grain size toward the center of the sealed cracks and the grain size proportional to the width of the sealed cracks as indicated by Toriumi and Hara (1995).

The growth front in the sealing of the cracks is considered to be parallel to the wall surface and it faces fluid phase in the cracks. The boundary behind the growth front is characterized by wavy interface, showing the various wavelengths of periodicity. This type of unstable boundary should be formed by the instability of growth front during precipitation of constituent minerals in active open cracks. The wavelength of the unstable boundary increases with increasing grain size. The comb type sealed cracks show narrow wavelength but the granular type ones do wide wavelength. In this study, the relationship between the unstable boundary and the instability of growth front during the sealing of cracks.

Crack geometries and deformation by the crack-seal mechanism in the Sambagawa metamorphic belt, Toriumi, M and E. Hara, *Tectonophysics* 245, 249-261, 1995.

Keywords: unstable boundary, crack sealing, instability of growth, plate boundary metamorphism

Fabric analyses of glaucophane and lawsonite in low-grade blueschist from Diablo Range, California

KIM, Daeyeong^{1*}, KATAYAMA, Ikuo¹, MICHIBAYASHI, Katsuyoshi², TSUJIMORI, Tatsuki³

¹Department of Earth and Planetary Systems Science, Hiroshima University, ²Institute of Geosciences, Shizuoka University,

³Pheasant Memorial Laboratory, Institute for Study of the Earth's Interior, Okayama University

Deformational microstructures of glaucophane and lawsonite in a lawsonite blueschist from New Idria serpentinite body, Diablo Range are studied to understand rheological behaviors of subducting oceanic crust. Glaucophane deforms by recovery and dynamic recrystallization possibly accommodated by dislocation creep, based on developments of crystal-preferred orientations (CPOs), small grain size and irregular or curved grain boundary. Euhedral or subhedral grains with angular or straight grain boundary of lawsonite suggest its deformation mechanism as rigid body rotation. Both minerals in glaucophane-rich layer (GR) show stronger CPOs, higher aspect ratios and lower angle to foliation (stronger shape-preferred orientation, SPOs) than those in lawsonite-rich area (LR), denoting that strain is mainly localized into the GR. In addition, stronger fabrics are observed in the GR rather than in the LR on the basis of fabric analyses ($M=0.20$, $J=18.0$ for glaucophane and $M=0.21$, $J=9.6$ for lawsonite in the GR, and $M=0.18$, $J=16.0$ for glaucophane and $M=0.15$, $J=7.8$ for lawsonite in the LR). All results of this study therefore indicate that rheological behaviors of subducting oceanic crust are mainly controlled by glaucophane rather than lawsonite.

Keywords: relative contrast, glaucophane, lawsonite, blueschist, subducting oceanic crust

Origin of compositional gradient recorded in metamorphic reaction rims

IKEDA, Takeshi^{1*}

¹Kyushu University

Localized occurrence of product minerals in reaction rims strongly suggests that the formation of the rim has been rate-controlled by diffusion. It has been a general consensus that the disequilibrium state during the rim formation was quenched, and that the chemical potential gradients were preserved in the rims, which was recognized as presence of compositional gradients of product minerals. This study propose alternative model to explain the compositional gradient occurring in the metamorphic coronas. In retrograde stages, migration distance (mean free path) decreases with descending temperature. The metamorphic reaction would cease when the migration distance sinks below the spatial distance between the reactant minerals. Each of reactant minerals that are chemically isolated could attain the local equilibrium. That is, the diffusion profiles maintained during the reaction would modify to homogenize after the reaction ceased. The continuous reaction produces the product minerals that change their composition progressively during descending temperature. Migration of components within crystal is much slower than that along grain boudary by a factor of 5 to 6. This suggests that compositional heterogeneity in product minerals would be likely to preserve even if local equilibrium was maintained among minerals within the migration distance.

Keywords: reaction rim, compositional gradient, fractional crystallization

Raman spectra of carbonaceous material in low-grade thermally-metamorphosed accretionary complex

HIDEKI, Mukoyoshi^{1*}, HIROSE, Takehiro², YAMAMOTO, Yuzuru³, SAKAGUCHI, Arito³

¹Marine Works Japan Ltd., ²JAMSTEC Kochi, ³JAMSTEC

Sedimentary rocks contain a trace amount of initially poorly ordered carbonaceous material (CM), which transforms into well-ordered graphite with increasing metamorphic grade. Previous study has demonstrated that peak metamorphic temperature (T) can be estimated by an area ratio R2 (=D1/[G+D1+D2]) of peaks recognized in Raman spectra of CM. This geothermometer can be used at temperature range of 330-650°C (e.g., Bayssac et al., 2002; Aoya et al., 2010). Herein, we present Raman spectra from a suite of samples with different metamorphic temperatures which are estimated by a vitrinite reflectance method: Miocene Hota complex (50°C) (Yamamoto et al., 2005), Cretaceous Shimanto complex (150°C and 230°C) (Mukoyoshi et al., 2006) and Jurassic Ashio complex (300°C). First-order Raman spectrum of CM often decomposed into four peaks of a Raman shift (G peak at about 1580cm⁻¹, D1 peak at about 1350cm⁻¹, D2 peak at about 1620cm⁻¹, D3 peak at about 1500cm⁻¹). In our amorphous CM (coal) samples we recognized other three peaks on the D1 peak around 1150 cm⁻¹, 1220 cm⁻¹ and 1450 cm⁻¹. These peaks has been also reported in (e.g., Bar-Ziv et al., 2000; Zaida et al., 2007; Potgieter-Vermaak et al., 2011).

The first-order Raman spectrums of our coal samples, in particular low-temperature samples, are hard to fit with decomposed four peaks using the LabSpec program due to the influence of faint shoulders on D1. Therefore, there is no clear correlation between T and average R2 ratio in each sample. However, the Raman spectrums can be fit with the above seven peaks. The correlation between R2 and T can be described by the following exponential equation:

$$T (^{\circ}\text{C}) = 8.6 * \exp(7.0 * R2) \quad (R^2 = 0.98)$$

In addition, when we use an area ratio of D1/[decomposed seven peaks] which is referred to as R6 in this study, the correlation between R6 and T is given by

$$T (^{\circ}\text{C}) = 10.9 * \exp(11.9 * R6) \quad (R^2 = 0.99)$$

These correlations can be used for a potential geothermometer for low-grade metamorphosed sediments, in the temperature range of 50-300°C.

Keywords: raman spectroscopy, vitrinite reflectance, carbonaceous material, accretionary complex, geothermometry

Structural change of carbonaceous material inferred from Raman spectra in low- to high-temperature metamorphic rocks

KOUKETSU, Yui^{1*}, MIZUKAMI, Tomoyuki², MORI, Hiroshi¹, ENDO, Shunsuke¹, Mutsuki Aoya³, HARA, Hidetoshi³, Daisuke Nakamura⁴, WALLIS, Simon¹

¹Graduate School of Environmental Studies, Nagoya University, ²College of Science and Engineering, Kanazawa University, ³Institute of Geology and Geoinformation, National Institute of Advanced Industrial Science and Techn, ⁴Department of Earth Sciences, Faculty of Science, Okayama University

We report the structural change of carbonaceous material (CM) from the samples metamorphosed at temperatures from 165 to 655 °C based on Raman spectroscopic analysis. 10 samples were selected from contact and regional metamorphic rocks of the Kasuga, Daimonji, and Shirataki areas and 9 samples were collected from accretionary complexes of the Kitagawa, Kure, and Nobeoka areas.

The shapes of CM Raman spectra show a change from broad and complex to sharp and simple ones with increasing metamorphic temperature. The D ("Defect")-bands dominate in the low temperature range and the G ("Graphite")-band increases in importance with temperature increase. These changes can potentially be used to construct a CM Raman geothermometer. However, there is no study defining a methodology for decomposing the CM Raman spectrum that is applicable to low to high temperature samples, and it is difficult to discuss the sequence of the development of CM. In particular, there has been little discussion of how peak decomposition of CM Raman spectra in the low temperature range should be carried out. In this study, several band combinations for the Raman spectrum of CM metamorphosed at the low temperature range were tested and parameters for each band were compared.

These studies show that the Raman spectrum of CM can be divided into four groups by the metamorphic temperature range: low-grade CM (150-280 °C), medium-grade CM (280-400 °C), high-grade CM (400-650 °C), and ordered graphite (> 650 °C).

Low-grade CM: The Raman spectrum of low-grade CM is best considered to consist of four D-bands (D1-, D2-, D3-, and D4-bands). The G-band is not clearly distinguishable in this temperature range. The shape of low-grade CM Raman spectra is qualitatively similar to those for amorphous carbon.

Medium-grade CM: The G-band first appears clearly in samples metamorphosed at temperatures of around > 280 °C. The D4-band disappears at around 340 °C and the prominence of the D3-band gradually decreases with increasing temperature. The peak positions of G-, D1-, and D2-bands and intensity ratio with D1- and D2-bands show changes in the temperature range of medium-grade CM. This observation indicates that CM transforms from amorphous carbon to disordered graphite in this temperature range.

High-grade CM and ordered graphite: Most band parameters become almost constant in the Raman spectra of high-grade CM. These changes suggest that the transition of CM from amorphous carbon to disordered graphite is complete at around 400 °C. A lot of CM grains in the highest temperature sample metamorphosed at 655 °C show spectra with only a G-band present and no recognizable D1- or D2-bands. This result indicates that CM grain reaches fully-ordered graphite at temperatures around 650 °C.

The above results are an important first step in establishing a reliable CM Raman geothermometer in low to high temperature range covering the transition of CM.

Keywords: Carbonaceous material, Raman spectroscopy, geothermometer, amorphous carbon, graphite

Low grade graphitization of deformation in Regional metamorphism

NAKAMURA, Yoshihiro^{1*}, AKAI Junji¹, TOYOSHIMA, Tsuyoshi²

¹Department of Geology, Sci,Niigata Univ, ²Grad. Sch. Sci. & Tech., Niigata Univ.

Graphitization in regional metamorphism is affected by various factors; thermal, pressures, time, shear stress, precursor materials, hydrothermal and catalyst(Large, 1994; Itaya,1980). Especially, it is well known that elastic deformation in graphitization is the one of the most important factors for acceleration of graphitization(Bustin,1995).

This study aims to verify how tectonic deformation influences graphitization process in the upper sequence of Hidaka metamorphic rocks (Zone1-2). The upper sequence of Hidaka Metamorphic belt is located in East side of Hidaka Mountains, Hokkaido, Japan. These metamorphic rocks are divided into metamorphic zoning from Zone1(Ms-Chl)to Zone2(Ms-Bt) by Osanai(1987) and ten stage(D0-D9) based on the nature and sequence of deformation by Koyasu(2006MS).

Degree of graphitization in deformed rocks that record elastic deformation (D2-D7 stage) was no variation, although these rocks with brittle fracturing (D8-D9 stage) compounded to low grade graphitization with bimodal distribution in Koikakusyu-satunai river route. High-grade crystalline graphite is distributed in matrix of metamorphic textures, on the other hand, low-grade crystalline graphite is distributed in vein like texture in Raman mapping.

These poorly crystalline graphite may deformed under the influence of brittle deformation process in later stage rather than in ductile deformation of peak metamorphic temperature. Moreover, these processes occurred in small crashed zones.

Bimodal distribution of graphitization in thin section scale may be common process in regional metamorphism. Relationship between deformation and graphitization will be discussed.

Keywords: Graphitization, Hidaka metamorphic belt, Micro-raman spectroscopy, XRD, HRTEM