

Calcification mechanisms in foraminifera and proxy incorporation Calcification mechanisms in foraminifera and proxy incorporation

Bijma Jelle^{1*}, Nehrke Gernot¹, Raitsch Markus¹, ド・ノイエ レナート², Funcke Antje¹, ケウル ニイナ³
Jelle Bijma^{1*}, Gernot Nehrke¹, Markus Raitsch¹, Lennart Jan de Nooijer², Antje Funcke¹, Nina Keul³

¹ アルフレッドウェゲナー研究所, ²NIOZ, ³Lamont-Doherty Earth Observatory

¹AWI, ²NIOZ, ³Lamont-Doherty Earth Observatory

Calcifying organisms, such as pteropods, bivalves, corals and foraminifera provide a rich resource for pale-oceanographers and climatologists because their geochemical make-up (proxies) can be used to reconstruct past ocean history and evolution during and after natural carbon perturbations. However, it has been shown for all geochemical proxies that the main assumption of only one environmental variable controlling a target proxy is too simple. Empirical calibrations introduce a lot of uncertainty because the mechanisms of proxy incorporation are not well understood. The major problem is that the calcification mechanisms are still a black box. In this presentation I will review our current understanding of calcification and proxy incorporation in foraminifera.

キーワード: 石灰化

Keywords: calcification, foraminifera

有孔虫が石灰化する時の環境変動
Environmental variation during foraminiferal calcification

豊福 高志^{1*}
Takashi Toyofuku^{1*}

¹ 海洋研究開発機構
¹JAMSTEC

Foraminiferal calcification process will be indicated with pH and Calcium variations.

キーワード: 有孔虫, 石灰化

蛍光色素を用いた共生藻をもつ有孔虫の室形成過程の解明

Distribution of intracellular pH and algal endosymbionts during chamber formation in symbiont-bearing reef foraminifers

藤田 和彦^{1*}, 由地 蛍¹, 豊福 高志²

Kazuhiko Fujita^{1*}, YUJI, Kei¹, Takashi Toyofuku²

¹ 琉球大学 理学部, ² 海洋研究開発機構

¹Univ. Ryukyus, ²Biogeos, JAMSTEC

Calcification mechanisms of algal symbiont-bearing reef foraminifers have not yet been well understood. In particular, it is cytologically not clear how the photosynthesis by algal endosymbionts enhance the calcification of host foraminifers. We visualized the distribution of intracellular pH and algal endosymbionts using a fluorescence probe HPTS as well as chlorophyll autofluorescence. High pH vesicles were gradually stored around symbionts before chamber formation. Our observations suggest that the photosynthesis by symbionts enhances the production of calcite needles or carbonate pools in vesicles by removing CO₂ around them.

Autoclave aquaria allow for high-pressure culture experiments on deep-sea benthic foraminifera
Autoclave aquaria allow for high-pressure culture experiments on deep-sea benthic foraminifera

Wollenburg Jutta^{1*}
Jutta Wollenburg^{1*}

¹ アルフレッドウェゲナー研究所
¹AWI

Our understanding of palaeodeep-water circulation modes and deep-water renewal in the world's oceans is essentially based on isotopical and geochemical ratios recorded in tests of calcareous deep-water benthic foraminifera that precipitate their shells in a constant ratio to the surrounding water mass. However, as more field data are assembled it appears that the documentation of deep-water in benthic foraminiferal shells is not always straightforward. Therefore, culture experiments on deep-sea benthic foraminifera are needed to verify the established palaeodeep-water proxies. However, to our knowledge, barophilic species like *F. wuellerstorfi*, the most trusted species for reconstructions of palaeodeep-water circulation and ventilation, neither formed new chambers nor reproduced in mesocosms kept at 1 bar. To accommodate this problem we have developed facilities and procedures that allow long-lasting high-pressure culture experiments on undepressurized deep-sea sediments and associated fauna and flora. In this presentation I will describe the experimental set-up and present results from methane-seepage simulations.

キーワード: 圧力, 飼育, 有孔虫
Keywords: pressure, culture, foraminifera

飼育実験手法を基にした溶存酸素濃度の浮遊性有孔虫への影響評価
Laboratory culture experiments: effect of dissolved oxygen concentration on planktonic foraminifera

黒柳 あずみ^{1*}, da Rocha Regine E.², Bijma Jelle², Spero Howard³, 川幡 穂高¹

Azumi Kuroyanagi^{1*}, Regine E. da Rocha², Jelle Bijma², Howard Spero³, hodaka kawahata¹

¹ 東京大学 大気海洋研究所, ² アルフレッドウェゲナー極域海洋研究所, ³ カリフォルニア大学 デービス校

¹ Atmosphere and Ocean Research Institute, The University of Tokyo, ² Alfred-Wegener-Institut für Polar- und Meeresforschung,

³ University of California Davis

Planktonic foraminifera shows significant species turnover ratio related Cretaceous oceanic anoxic event (OAEs), however, the direct effect of dissolved oxygen (DO) concentration on planktonic foraminifera remain obscure. Although culture experiments can investigate quantitatively the relationship between foraminiferal ecology and environmental parameters, DO-controlling experiment has never been conducted because of the difficulty of observation and/or control of dissolved oxygen under the modern ocean condition. In the present study, we cultured two planktonic foraminifera species (*Orbulina universa* and *Globigerina bulloides*) under six different DO conditions. Both species have extremely high tolerance to low DO than we expected before, and it suggests that at least "dysoxic" condition (more than 0.7 mg O₂ l⁻¹) could not be a direct cause of the extinction of planktonic foraminifera at OAEs. Their high tolerance to extremely low DO would be caused by the evolutionary descendant of benthic foraminifera. Final shell weight increased with DO despite almost the same culture duration among treatments, thus it suggests foraminiferal fossil shell weight could reflect the DO conditions.

キーワード: 浮遊性有孔虫, 飼育実験, 溶存酸素濃度

Keywords: planktonic foraminifer, culture experiment, dissolved oxygen concentration

亜北極・北極 *Emiliana huxleyi* のココリス形態の培養温度に対する応答 Subarctic and Arctic *Emiliana huxleyi* coccolith morphological responses to the growth temperatures

猿渡 和子^{1*}, 佐藤真奈美¹, 原田尚美², 鈴木石根¹, 白岩善博¹

Kazuko Saruwatari^{1*}, Manami Satoh¹, Naomi Harada², Iwane Suzuki¹, Yoshihiro Shiraiwa¹

¹ 筑波大学生命環境科学, ² 海洋研究開発機構

¹Faculty of Life and Environmental Sciences, University of Tsukuba, ²Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology

Coccolithophore is a key calcified phytoplankton for biogeoscience because of their specific products such as calcified scales, coccolith, and unsaturated ketones, alkenone. As a biomarker, the chemical compositions of alkenones are often used to estimate paleo ocean temperatures and to identify the changes of the marine environments (e.g., Harada et al., 2012). In particular, the polar region is an important area to recognize the changes of the global environments and the sea-ice melting is known to be accelerated by the global warming. MIRAI subarctic expeditions have been continued to investigate the subarctic to Arctic environment and concurrently the culture experiments of marine organisms including coccolithophores have been performed to predict the future changes of marine ecosystem. In this study, two *Emiliana huxleyi* strains, MR57N and MR70N, isolated by MIRAI subarctic expedition in 2010 were cultured at the different temperatures (5, 10, 15 and 20°C) and salinities (25, 32, and 35 per mill) and investigated from the viewpoint of the coccolith morphology. As the results, these strains showed the similar growth properties that the growth rate increased with raising the temperature. According to SEM observations and the image analyses, the size (length of distal shield: LDS) and the numbers of distal shield elements of the coccolith decreased with raising the temperature. The central area of the coccoliths was also changed from grill structures to completely calcified structures. Moreover, the cell size of *E. huxleyi* decreased with raising the temperature. Thus, the subarctic and Arctic *E. huxleyi* strains showed the correlations between cell sizes and coccolith morphometric parameters with variable central area morphology depending on the growth temperatures. For the salinity experiments, there is almost no variation on the growth properties and a little variation on the morphometrical parameters. These results imply that the subarctic and arctic *E. huxleyi* strains can maintain enough to grow in the temperature range between 5 to 20°C and the salinity range between 25 to 35 per mill, suggesting that these strains can ideally survive under the warmer and/or the less salinity environments. The coccolith morphologies and the parameters vary in response to the growth temperature but not to the salinity. This relationship was applied to the natural plankton samples reported by Bollmann et al. (2009) and the part of their data was correlated with sea water temperatures but the data shallower than 15 m depth was not. This may imply the importance of the light intensity and the further culture experiments with the different light intensity are required.

Harada et al. (2012) *Global Biogeochemical Cycles* 26, GB2036.

Bollmann et al. (2009) *EPSL* 284, 320.

放散虫 *Spongotrochus glacialis* にみられる 2 隠蔽種の形態的特徴及び好適環境要因
Morphological and ecological characters of two cryptic genetic types in the radiolarian
Spongotrochus glacialis.

石谷 佳之^{1*}, 氏家 由利香², 瀧下 清貴¹
Yoshiyuki Ishitani^{1*}, Yurika Ujii², Kiyotaka Takishita¹

¹ 海洋研究開発機構 深海生態系研究チーム, ² 高知大学海洋コア総合研究センター

¹Japan Agency for Marine-Earth Science and Technology, ²Center for Advanced Marine Core Research, Kochi University

The discovery of cryptic species often helps in grasping the true diversity and detailed ecological preferences of pelagic planktons. Radiolarians are a good environmental proxy for paleontology because of their worldwide oceanic distribution and the good preservation of their shells. However, cryptic species have not yet been recognized in this planktonic protist group. In the present study, we focused on the morphospecies *Spongotrochus glacialis* with specimens collected from different layers in the Equatorial-Subtropical Pacific. A molecular phylogeny based on the internal transcribed spacer of rDNA (ITS-rDNA) sequences recovered two clades of cryptic species (types I and II) from this single morphospecies. These two distinct types were separately distributed, either in the oligotrophic surface water (type I) or below the chlorophyll maximum layer (type II). Moreover, the types showed morphological differences in the shells. Our morphometric analyses established lengths of spines as a morphological criterion to distinguish between the two types: type I with longer spines and type II with shorter spines. The length of spines is apparently associated with the habitat of each type. Type I with longer spines could be suitable for extending flagella and floating on the surface, whereas type II with shorter spines are appropriate to protrude only short flagella and dwell in deep water. Such morphological and ecological features at the cryptic species level of Radiolaria could provide new proxies for paleoceanographic studies.

キーワード: 放散虫, 隠蔽種, ITS rDNA, 形態計測

Keywords: Radiolaria, cryptic species, ITS rDNA, morphometric

南シナ海北部の環境DNA中の放射虫類18S rDNAの系統解析

The Phylogenetic Affiliations of Radiolarian-like Environmental 18S rDNA Genes from the Northern South China Sea

李 緑硯^{1*}, 遠藤一佳¹

Luyan Li^{1*}, Endo Kazuyoshi¹

¹ 東京大学

¹The University of Tokyo

To determine the molecular taxonomic affiliations of radiolarian-like environmental 18S rDNA genes from the northern South China Sea, we constructed phylogenetic trees by the 18S rDNA sequences of identified radiolarian species, correlating environmental sequences from other sea areas with those full length 18S rDNA gene of representative clones from the northern South China Sea. In our previous work, a great number of our radiolarian-like sequences from the South China Sea were placed within the RAD-III, named by Not et al. (2007) of environmental sequences from the Sargasso Sea size-fractionated samples (< 2 micrometre). Now we found 18S rDNA sequences of two identified species *Arachnosphaera myriacantha* and *Astrosphaera hexagonalis*, collected from Okinawa Island by Yuasa et al. (2009) also belongs to RAD-III. There is a unique clade composed solely of environmental clone sequences formed basal to the Taxopodida. It does not belong to RAD IV or V, still remains unknown. Our *Nassellaria*-like environmental clone from the northern South China Sea was joined with the group of described *Plagoniid* species. And the clones from Cariaco Basin, Caribbean clustered with *Pterocorythid* and *Theoperid* groups. Comparison of our sequences with two recent survey of *Acantharia* 18S rDNA, the position of northern South China Sea clones was still difficult to distinguish. It belonged to the clade composed of subgroups *Arthracanthida*, *Sphaenacanthida* and RAD I (Unidentified Clade 1). These ambiguities may be because the genetic diversity of radiolarian in warm waters sea areas is still poorly known.

シアノバクテリア化石の顕生代海洋における pH プロキシとしての可能性 Fossil cyanobacteria as a potential pH proxy for Phanerozoic ocean

白石 史人^{1*}

Fumito Shiraishi^{1*}

¹ 白石史人

¹SHIRAISHI, Fumito

Fossil cyanobacteria appeared in the most part of Phanerozoic era, and its abundance changed significantly through time. Their fossilization related to calcification in the living state, which requires optimum chemistry in the ambient water. Therefore, the fossil record of cyanobacteria potentially reflects past ocean chemistry. Recent study revealed that cyanobacterial calcification depends mainly on pH, dissolved inorganic carbon, Ca^{2+} concentration (Shiraishi 2012, GCA). Using well-established proxies of Ca^{2+} and partial pressure of CO_2 , oceanic pH was estimated from the fossil record of calcified cyanobacteria. Estimated range and trend of pH are similar to those of previous studies, but exhibited more frequent changes. In a future study, it is necessary to understand the relationship between CaCO_3 saturation state achieved by photosynthesis and calcification amount, in order to increase the reliability of estimation.

Benthic foraminifera as Quaternary palaeoclimate proxies in the New Caledonia Basin Benthic foraminifera as Quaternary palaeoclimate proxies in the New Caledonia Basin

Briony Mamo^{1*}, Glenn Brock², Elsie Gretton³

Briony Mamo^{1*}, Glenn Brock², Elsie Gretton³

¹JAMSTEC, ²Macquarie University, ³Boggabri Coal

¹JAMSTEC, ²Macquarie University, ³Boggabri Coal

In 2001, gravity core GC4 was extracted from the New Caledonia Basin and the uppermost 141 cm, representing approximately the last 140 000 years, was investigated for chemical isotopic, carbonate, non-carbonate and trace element signatures. In this study, sub samples of the same uppermost 141 cm of GC4 were picked and sorted for all foraminiferal taxa. A total of 161 species of benthic foraminifera were obtained from the core and 46 species occurred in sufficient numbers to investigate changing patterns and trends in biodiversity and relative abundance. All changes can be related to broad scale oceanic and palaeoclimatic fluctuations during the last 140 000 years. Within GC4, two distinct foraminiferal assemblages were detected using Bray-Curtis cluster analysis and Multidimensional Scaling (MDS) analysis. Factors that influence the foraminiferal assemblages include environmental setting, SSP, oxygen levels at the sediment-water interface and transportation by oceanic current systems operating within the region.

キーワード: foraminifera

Keywords: foraminifera

南日本で採集した化石シャコガイ殻のSr/Ca比を用いた4.6kaの日射量の復元
Daily light cycle reconstructed by Sr/Ca in a fossil giant clam, *Tridacna gigas*, living in
4.6 ka, southern Japan

堀 真子^{1*}, 白井厚太郎¹, 高畑 直人¹, 佐野 有司¹, 小林夏子², 岡本 和明², 渡邊 剛³

Masako Hori^{1*}, Kotaro Shirai¹, Naoto Takahata¹, Yuji Sano¹, Natsuko Kobayashi², Kazuaki Okamoto², Tsuyoshi Watanabe³

¹ 東京大学大気海洋研究所, ² 埼玉大学教育学部, ³ 北海道大学理学部

¹Atmosphere and Ocean Research Institute, the University of Tokyo, ²Faculty of Education, Saitama University, ³Faculty of Science, Hokkaido University

Giant clams are long-lived bivalvia living in a shallow tropical ocean. Their aragonite shells are potential paleo-environmental archives, i.e., oxygen stable isotopic values record paleo-seawater temperatures. Generally, trace elements in calcium carbonates are also considered as a paleo-thermometer. However, the case of Tridacninae may not follow the traditional temperature dependent fractionation, but potentially reflects insolation. This is explained by "trans-calcification", which is one of photosynthetically induced calcification mechanisms. Enzymatic reaction pumping Ca^{2+} to host liquid of calcification results in relatively depleted Sr/Ca ratio in day time when photosynthesis activates. Following this hypothesis, we have reconstructed the past daily light cycle.

A fossil *T. gigas*, Stg04-b was collected at Ishigaki-jima Island in Okinawa Prefecture, Japan (N24 20'0.4" E124 09'22"). Two horizons of the specimen were dated by carbon-14 method at the accelerator mass spectrometry (AMS) center of Yamagata University. The micro analyses were performed for the outer layer of 1.4 cm thickness using an electron probe micro-analyzer (EPMA), JXA8900, and a micrometer-scale secondary ion mass spectrometer (NanoSIMS), CAMECA NS50 at the Atmosphere and Ocean Research Institute (AORI), the University of Tokyo. As a result, we found clear Sr lamination parallel to the daily increment of about 20 μm thickness. The hourly insolation (P) was calculated from the difference of Sr/Ca ratio ($\Delta\text{Sr}/\text{Ca}$) following the equation: $\Delta\text{Sr}/\text{Ca} = -a \times P$, where the parameter, a , is defined by analysis of modern *T. maxima* [1]. Using the technique, we distinguished the sunny and cloudy seasons in 4.6 ka.

Reference

[1] Sano, Y., Kobayashi, S., Shirai, K., Takahata, N., Matsumoto, K., Watanabe, T., Sowa, K., Iwai, K. 2012, Past daily light cycle recorded in the strontium/calcium ratios of giant clam shells. Nature Commun. 3, DOI: 10.1038/ncomms1763

Keywords: giant clam, Sr/Ca, NanoSIMS

「汚物だめ保存仮説」: 例外的に保存されたオルステンタイプ化石の化石化のメカニズムについて

The cesspool preservation hypothesis as a key to preservation of exceptionally well preserved orsten-type fossils

田中 源吾^{1*}, 前田晴良², 松島義章³

Gengo Tanaka^{1*}, Haruyoshi Maeda², Yoshiaki Matsushima³

¹群馬県立自然史博物館, ²九州大学博物館, ³神奈川県立生命の星・地球博物館

¹Gunma Museum of Natural History, ²Kyushu University Museum, ³Kanagawa Prefectural Museum of Natural History

最近, 神奈川県追浜公園にて掘削された完新世のコア試料のうち, ペレットを大量に含む層準より, 例外的に軟体部が保存された介形虫(節足動物)化石が発見された(Tanaka et al., 2012)。

同様な軟体部が保存された節足動物の化石は, カンブリア紀後期のオルステン石灰岩中에서도発見されている(Maeda et al., 2011)。そのような「汚物だめ」は, 集積したペレットによって作りだされた局所的な磷酸塩の濃集による例外的な磷酸塩化によるものであると考えられる。この「汚物だめ保存仮説」はカンブリア紀後期以降の海成堆積物においてみられる, 例外的に保存の良い化石の化石化の解明につながると考えられる。

キーワード: 化石, カンブリア紀, 汚物だめ, 節足動物, 古生物学, 完新世

Keywords: Fossil, Cambrian, Cesspool, Arthropoda, Paleontology, Holocene