

初期カンブリア紀の高い大陸風化速度 Intensive continental weathering rate in the Lower Cambrian: evidenced from Sr isotope ratios preserved in the strata at

澤木 佑介^{1*}, 小宮 剛², 丸山 茂徳¹
Yusuke Sawaki^{1*}, Tsuyoshi Komiya², Shigenori Maruyama¹

¹ 東京工業大学, ² 東京大学

¹Tokyo Tech, ²The University of Tokyo

One of the most important times in biological evolution was from the first appearance of soft-bodied animals and possibly Metazoan animals in the late Neoproterozoic to the sudden diversification of animals with mineralized skeletons in the Cambrian. Neoproterozoic to Cambrian fossiliferous succession is well exposed in South China (e.g., Luo et al., 1984); it has been much investigated using biostratigraphy, sequence stratigraphy and chemostratigraphy (e.g. Condon et al., 2005; Zhou and Xiao, 2007; Zhu et al., 2007). We carried out on-land drilling of the Ediacaran to Cambrian sedimentary succession in Three Gorges, South China. The drill-sampling allows us to minimize the effect of secondary alteration and oxidation on the surface and to make a very continuous chemostratigraphy at intervals of centimeters. Now, members of our group at Tokyo Tech and the Univ. of Tokyo have analyzed several kinds of isotope ratios to estimate paleo-environment; including carbon, oxygen, nitrogen and strontium isotope ratios.

The weathering influx from continents is thought to be a major influence on the change in composition of ancient seawater and on biological evolution. Its flux can be estimated from the $87\text{Sr}/86\text{Sr}$ ratio of carbonate rocks. We present a new detailed chemostratigraphy of $87\text{Sr}/86\text{Sr}$ in the Three Gorges region in South China. The result shows that $87\text{Sr}/86\text{Sr}$ ratios had decreased from ca. 0.709 around the Precambrian/Cambrian boundary to ca. 0.7085 at the Atdabanian stage. Subsequently, $87\text{Sr}/86\text{Sr}$ ratios recovered to ca. 0.709 at the end of Early Cambrian. We calculated secular variation of weathering flux using the analyzed data, suggesting intensive continental weathering rate in the Lower Cambrian.

Keywords: $87\text{Sr}/86\text{Sr}$, continental weathering, Early Cambrian, Three Gorges, drill-core

三疊紀、ジュラ紀深海堆積物中に保存された白金族元素濃度異常 Platinum group element anomalies in the Triassic-Jurassic deep-sea sediments

藤崎 渉^{1*}, 澤木 佑介¹, 山本 伸次², 横山 哲也¹, 丸山 茂徳¹

Wataru Fujisaki^{1*}, Yusuke SAWAKI¹, Shinji YAMAMOTO², Tetsuya YOKOYAMA¹, Shigenori MARUYAMA¹

¹ 東京工業大学大学院理工学研究科地球惑星科学専攻, ² 東京大学大学院総合文化研究科広域科学専攻

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²Department of Earth Science and Astronomy, University of Tokyo

One of the biggest mass extinctions in the Phanerozoic occurred at the Triassic-Jurassic (T-J) boundary (e.g., Sepkoski, 1984). The large magmatic activity associated with the breakup of Pangaea and the initial stage of rifting in the Central Atlantic Magmatic Province (CAMP) are characteristic across the T-J boundary (McHone, 2000; Nomada, 2007). So, these magmatic activities likely changed the climate and caused biotic crisis at the T-J boundary (e.g., Deenen et al., 2010). On the other hand, Olsen and others (2002) suggested that bolide impact triggered the climatic change and mass extinction, based on an Ir anomaly preserved in the Newark rift basin. In addition, it is also proposed that the encounter with dark clouds and supernova explosion caused extreme environmental change on the earth surface (the mass extinction and glaciation; Kataoka et al., 2012). However, the possibilities of these hypotheses remain controversial because of insufficient geological evidence.

In this research, we present secular variation of platinum group elements (PGEs) concentration in the Triassic-Jurassic succession in Inuyama, central Japan. Previous Ir anomalies have been reported from sediments with high sedimentation rate, around the T-J boundary (Olsen et al., 2002; Hori et al., 2007). However, sediments with low sedimentation rate are suitable for the PGEs analyses. In the Inuyama, the depositional rate of the shale part in bedded chert is about one to two orders slower than those of the chert (Hori et al., 1993). Therefore, the best target for the PGEs analyses is the shale part in bedded chert preserved in the accretionary complex.

We conducted geological survey at the Inuyama section, because of good exposure of the T-J boundary. We developed detailed geological map of the study area and collected rock samples bed-by-bed to determine the secular variation of PGEs concentrations. In particular, we collected about 250 samples from shale part in the bedded chert and analyzed the PGEs concentrations of 20 shale samples across the T-J boundary. For whole-rock analyses of PGEs, all shale samples were powdered in an Alumina planetary mill. After chemical separation from coexisting matrix elements using a chromatographic technique, PGEs concentrations were analyzed by coupled plasma mass spectrometry (ICP-MS) at Tokyo Institute of Technology. The PGEs concentrations were determined by isotopic dilution method. The results show that Ir concentration reach ca. 1 ppb just above the T-J boundary. As compared to the previous works (Olsen et al., 2002; Hori et al., 2007), the Ir anomaly is the highest across the T-J boundary and attributes to the difference of their depositional rate and/or sampling resolution. Olsen and others (2002) suggested the possibility of volcanic or impact events for origin of the Ir anomaly. The Ir anomaly in this research also may be associated with impact event, despite the lack of shocked quartz and other index of impact origin (e.g., Grieve et al., 1996). In order to recognize the correlation between PGEs concentrations and CAMP, we need additional stratigraphic and isotopic analyses.

In this presentation, we would like to discuss the origin of the Ir anomaly and its relation to evolution of life.

Reference

- Deenen, M. H. L. (2010). *Earth and Planetary Science Letters*, 5, 291, pp. 113-125.
Grieve, R. A. F. (1996). 2, *Significance in geoscience: Meteoritics*, 31, pp. 6?35.
Hori, R. (1993). *Island Arc*, 3, pp. 170-180.
Hori, R. (2007). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 5, 244, pp. 391-406.
Kataoka, R. (2012). *New Astronomy*, 21, pp. 50-62.
McHone, J. G. (2000). *Tectonophysics*, 5, 316, pp. 287?296.
Nomada, S. (2007). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 5, 244, pp. 326-344.
Olsen, P. (2002). *The Geological Society of America Special Paper*, pp. 505-522.
Sepkoski, J. J. (1984). *Paleobiology*, 10, pp. 246-267.

Keywords: the Triassic-Jurassic boundary, deep-sea sediments, platinum group elements

深海性堆積物中の白金族元素濃度を利用した宇宙起源物質の検出 Detection of cosmogenic material in deep-sea sediments based on platinum group element (PGE) abundances

浅沼 尚^{1*}, 藤崎 渉¹, 澤木 佑介¹, 丸山 茂徳¹

Hisashi Asanuma^{1*}, Wataru Fujisaki¹, Yusuke Sawaki¹, Shigenori Maruyama¹

¹ 東京工業大学地球惑星科学専攻

¹Tokyo Institute of Technology

Snowball Earth events are widely recognized to have occurred in both the Palaeoproterozoic and Neoproterozoic. All present-day animal phyla appeared following the Marinoan-Snowball Earth, which is the last recorded global glaciation. The primary objective of this research is to determine the cause of this Snowball-Earth event, which is likely associated with the evolution of life. Three models that attempt to explain the onset and termination of Snowball-Earth events are: (1) episodic decrease of greenhouse gases, (2) changes of the albedo of the Earth accompanied by the arrangement of the continents, and (3) an increase in cosmic-ray bombardment to the Earth due to Starbursts in the Milky Way Galaxy or transects of Earth through nebula.

Based on recent investigations, the effects of cosmic fluxes on the Earth are estimated here through the measurement of the abundances of platinum group elements (PGE) in sediments.

Pelagic sediments composed of interlayered shale and/or mudstones are optimal for PGE-abundance analysis because of their low sedimentation rate. Pelagic sediments used for PGE-abundance analysis in this investigation are comprised of bedded shales and/or mudstones sampled from the accretionary complex of Anglesy-Llyen, U.K. The Anglesy-Llyen reportedly was formed by an accretionary orogeny in the Neoproterozoic. During the geological survey, samples were acquired from a pelagic sedimentary sequence, which records late Cryogenian to early Cambrian sedimentation. A relatively high PGE concentration and its flat C1-normalized PGE pattern indicate possible high cosmic fluxes on the Earth during the emplacement of this sequence when compared to average upper continental crust composition.

Keywords: PGEs, deep-sea sediment, Cambrian

超海洋中央部起源の石灰岩における G-L 境界大量絶滅事件周辺の高分解能化学層序 High-resolution litho- and chemo- stratigraphy across the Middle-Late Permian boundary in the mid-oceanic limestone

小福田 大輔^{1*}, 磯崎行雄¹, 猪郷久義², 可児智美³, 石村豊穂⁴

Daisuke Kofukuda^{1*}, ISOZAKI, Yukio¹, IGO, Hisayoshi², KANI, Tomomi³, ISHIMURA, Toyoho⁴

¹ 東京大学総合文化研究科, ² 自然史科学研究所, ³ 熊本大学自然科学研究科, ⁴ 産業技術総合研究所

¹The University of Tokyo, Komaba, ²Institute of Natural History, ³Kumamoto University, ⁴AIST

The extinction around the Guadalupian-Lopingian boundary (G-LB) recorded the first major biodiversity drop during the Permian. In order to clarify the relevant global environmental changes immediately before the G-LB, high-resolution chemostratigraphy of $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{13}\text{C}_{\text{carb}}$ was analyzed for the Middle Permian paleo-atoll limestone at Akasaka and Ishiyama in central Japan.

Both limestones were derived from a paleo-atoll complex deposited primarily in the low latitude a mid-Panthalassa. At both sections, the Middle Permian black limestone is overlain by the Upper Permian light gray limestone. An interval of white-black striped limestone occurs between the two, and its top marks the G-LB horizon. Fusuline biostratigraphy indicates that the black limestone belongs to the Yabeina Zone (Capitanian; late Guadalupian), whereas the light gray limestone to the Codonofusiella-Reichelina Zone (Wuchiapingian; early Lopingian). The "barren interval" between them is divided into the lower 1) black limestone, and the upper 2) striped limestone without smaller foraminifer and gastropod.

The lithofacies of the black limestone and of the striped limestone indicates that the depositional setting was subtidal zone and intertidal zone, respectively.

At the Akasaka Limestone, a very thin (<5 mm-thick) light greenish non-carbonate layer occurs between the striped limestone and the light gray limestone, which was once reported as a felsic tuff. This layer is enriched in elements such as Al, Fe, K and Cr. The source was not yet identified, however, this layer is possibly correlated with the Wangpo Bed, i.e., the G-LB marker in South China.

This study confirms that $^{87}\text{Sr}/^{86}\text{Sr}$ ratios stayed extremely low around 0.7068 throughout the Capitanian and increased rapidly to 0.7074 at the G-LB, and that the "Permian minimum" has persisted throughout the entire Capitanian for more than 5 million years. The $\delta^{13}\text{C}_{\text{carb}}$ values stayed extremely high, +6 permil, throughout the Capitanian, and dropped to +2 permil at the G-LB. This records, no doubt, the Capitanian Kamura event of the very high primary productivity of surface ocean. These isotope profiles are correlated well with those of the coeval Iwato Formation in Kyushu, thus likely reflecting the general trend of the low latitude mid-superocean seawater.

The change in lithofacies towards the G-LB recorded that the depositional setting reached the shallowest during the deposition of striped limestone. This implies that sea level became shallower towards the G-L boundary. In accordance with the lowest sea level around the G-L boundary, this may suggest that global cooling has appeared immediately before the G-L boundary and possibly caused the end-Capitanian extinction.

キーワード: G-L 境界, 赤坂石灰岩, 石山石灰岩, 炭素同位体, Sr 同位体

Keywords: G-L boundary, Akasaka Limestone, Ishiyama Limestone, carbon isotope, strontium isotope

パンサラサ海中央部の海山起源ペルム紀石灰岩の岩相・生・C-Sr同位体層序
Litho-, bio-, and C, Sr isotope stratigraphy of the Middle Permian mid-Panthalassan
paleo-atoll carbonates

西田 昂広^{1*}, 磯崎 行雄¹, 可児 智美², 鈴木 淳³, 石村 豊穂³

Takahiro Nishida^{1*}, Yukio Isozaki¹, Tomomi Kani², Atsushi Suzuki³, Toyoho Ishimura³

¹ 東京大学大学院総合文化研究科, ² 熊本大学理学部, ³ 産総研 地質情報研究部門

¹Department of Earth Science & Astronomy, The University of Tokyo, ²Faculty of Science, Kumamoto University, ³AIST Geological Survey of Japan

The Wordian-Capitanian (Middle-Upper Guadalupian, Permian) Iwato Formation in East Kyushu is composed of shallow marine carbonates in an accreted paleo-atoll complex primarily developed on a paleo-seamount in the low-latitude domain (12 degree S) of mid-Panthalassa. By analyzing the Wordian interval, we could newly reconstruct a continuous C-isotope profile of the upper half of Guadalupian, and discuss the environmental changes in the low-latitude mid-superocean. We confirmed that the high C isotope interval (> 5 permil) had started already in the Wordian. The onset of the Kamura cooling event likely started much earlier than previously believed, and affected the diversity of the Guadalupian biota.

キーワード: ペルム紀, 岩戸層, 古海山, パンサラサ海, 上村事件, フズリナ

Keywords: Permian, Iwato Formation, paleo-seamount, Panthalassa, Kamura event, fusuline

南部北上帯中部ペルム系岩井崎石灰岩最上部の詳細層序 Detailed stratigraphy of the uppermost Iwaizaki limestone (Middle Permian) in the South Kitakami beltin

林 隆太郎^{1*}, 磯崎 行雄¹
Ryutaro Hayashi^{1*}, Yukio Isozaki¹

¹ 東京大学大学院総合文化研究科

¹Department of Earth Science & Astronomy, The University of Tokyo

南部北上帯のペルム系岩井崎石灰岩は陸棚層の浅海成石灰岩からなる。フズリナ生層序に基づくと、その上半部は中期ペルム紀後期キャピタン世の年代を持つ。しかし、その最上部の層厚約30 mの区間の詳細な岩相や年代はよくわかっていない。本発表では、その部分の詳細な岩相層序について報告し、特にキャピタン世を示すフズリナ *Lepidolina multiseptata* の産出上限周辺での環境変化について考察する。

キーワード: ペルム紀, 岩井崎石灰岩, 南部北上帯, フズリナ, 軟体動物, キャピタン世

Keywords: Permian, Iwaizaki limestone, South Kitakami belt, fusuline, mollusc, Capitanian

中部ペルム系岩井崎石灰岩（南部北上帯）の古地理学的位置 Paleogeographic position of the Permian Iwaizaki limestone in South Kitakami belt

磯崎 行雄^{1*}, 林 隆太郎¹
Yukio Isozaki^{1*}, Ryutaro Hayashi¹

¹ 東京大学大学院総合文化研究科

¹Department of Earth Science & Astronomy, The University of Tokyo

The Guadalupian (Middle Permian) Iwaizaki limestone in the South Kitakami belt, NE Japan, represents an isolated block of ancient continental margin that features non-metamorphosed Lower Paleozoic to Mesozoic shallow marine sequences. We detected for the first time a bivalve assemblage that features alatoconchidae from the upper part of the Iwaizaki limestone. According to the 10 previous reports from the world, the occurrence of alatoconchidae is strictly limited to low-latitude, i.e. paleo-equatorial domains. Their unique habitat in shallow warm-water, oligotrophic setting was likely related to photosymbiosis. Together with large-tested fusuline (e.g., *Lepidolina*) and rogoose coral, the Capitanian tropical trio from Iwaizaki positively indicates the intimate connection between South Kitakami belt and South China during the Permian. This further suggests that the eastern extension of South China continues all the way through the main part of Japan up to NE Japan, and that the South Kitakami belt represents its eastern extremity ever confirmed.

キーワード: ペルム紀, 古地理, 南部北上帯, パンサラサ海, 南中国, ガダルーブ世

Keywords: Permian, paleogeography, South Kitakami belt, Panthalassa, South China, Guadalupian