

## GELCA 結合モデルを利用した二酸化炭素の10年間インバージョン A decadal inversion of carbon dioxide using the Global Eulerian-Lagrangian Coupled Atmospheric model (GELCA)

白井 知子<sup>1\*</sup>, 石澤みさ<sup>1</sup>, Ruslan Zhuravlev<sup>2</sup>, Alexander Ganshin<sup>2</sup>, 佐伯田鶴<sup>1</sup>, Belikov Dmitry<sup>1</sup>, 小田知宏<sup>3</sup>, 齊藤誠<sup>4</sup>, Vinu Valsala<sup>5</sup>, Maksyutov Shamil<sup>1</sup>  
Tomoko Shirai<sup>1\*</sup>, Misa Ishizawa<sup>1</sup>, Ruslan Zhuravlev<sup>2</sup>, Alexander Ganshin<sup>2</sup>, Tazu Saeki<sup>1</sup>, Dmitry Belikov<sup>1</sup>, Tomohiro Oda<sup>3</sup>, Makoto Saito<sup>4</sup>, Vinu Valsala<sup>5</sup>, Shamil Maksyutov<sup>1</sup>

<sup>1</sup> 国立環境研究所, <sup>2</sup> ロシア中央高層気象台, <sup>3</sup> 国立環境研究所 (現所属: 米 CSU/NOAA ESRL), <sup>4</sup> 国立環境研究所 (現所属: 仏 LSCU), <sup>5</sup> 国立環境研究所 (現所属: インド熱帯気象学研究所)

<sup>1</sup>National Institute for Environmental Studies, <sup>2</sup>Central Aerological Observatory, <sup>3</sup>NIES, now at CSU/NOAA ESRL, <sup>4</sup>NIES, now at LSCU, <sup>5</sup>NIES, now at ITTM

A decadal estimate of global CO<sub>2</sub> flux distribution for the period of 2001-2010 was conducted using an atmospheric inversion modeling system called GELCA (Global Eulerian-Lagrangian Coupled Atmospheric model) with Kalman smoother inversion technique. The use of Lagrangian particle dispersion model (LPDM) to simulate the transport in the vicinity of the observation points enables us to avoid numerical diffusion from which Eulerian models suffer, and is suitable to represent observations at high spatial and temporal resolutions. An Eulerian model is run to generate the global background concentrations to be used as the boundary conditions for an LPDM that performs backward simulations from each receptor point (observation location). In GELCA, National Institute for Environmental Studies-Transport Model (NIES-TM) version 8.1i was used as an Eulerian global transport model coupled with FLEXPART version 8.0 as a LPDM. Two-day backward transport by FLEXPART was combined with the background CO<sub>2</sub> levels 2 days prior to the observations simulated by NIES-TM. The meteorological data for driving both models was taken from JMA Climate Data Assimilation System (JCDAS) with a spatial resolution of 1.25° x 1.25° and a temporal resolution of 6 hours. Our prior CO<sub>2</sub> fluxes consist of the following four types: daily terrestrial biospheric fluxes generated by the VISIT model (Vegetation Integrative Simulator for Trace gases); monthly oceanic fluxes generated by an ocean pCO<sub>2</sub> data assimilation system; monthly biomass burning emissions taken from the Global Fire Emissions Database (GFED), version 3.1; and monthly fossil fuel emissions combining the high-resolution Open source Data Inventory of Anthropogenic CO<sub>2</sub> emission (ODIAC) version 3.0 dataset. We employed a Kalman Smoother inversion technique with fixed lag of 3 months, solving for 42 land and 22 ocean regions.

The purpose of the present study is to evaluate the performance of the GELCA inversion system with rather long period (10 years) CO<sub>2</sub> flux estimation and to examine the impact of observation network. We tested several different sets of observation datasets starting by using the NOAA flask network ground based observations as a control case. The sensitivity of the inversion to the choice of CO<sub>2</sub> observation dataset was discussed using the footprint of each observation dataset. The CO<sub>2</sub> flux estimate was examined in terms of observation network/coverage and also compared with previous studies.

キーワード: CO<sub>2</sub>, 炭素循環, 逆計算, 結合モデル

Keywords: CO<sub>2</sub>, sources/sinks, inverse modeling, coupled model