

Recent variation of West Siberian wetland CH₄ fluxes estimated from atmospheric CH₄ Recent variation of West Siberian wetland CH₄ fluxes estimated from atmospheric CH₄

金 憲淑¹, 笹川 基樹¹, 町田 敏暢¹, Dmitry Belikov¹, 佐伯 田鶴¹, 伊藤 昭彦¹, Maksyutov Shamil^{1*}
Heon-Sook Kim¹, Motoki Sasakawa¹, Toshinobu Machida¹, Dmitry Belikov¹, Tazu Saeki¹, Akihiko Ito¹, Shamil Maksyutov^{1*}

¹ 国立環境研究所

¹National Institute for Environmental Studies

The world's largest extent of wetlands occurs in West Siberia, where wetlands account for 27% of the area of West Siberia (Peregon et al. 2009). The vast wetlands emit methane (CH₄) to the air and the magnitude depends highly on soil temperature and the water table. A large increase of atmospheric CH₄ was observed globally in 2007 and Siberian wetland emission enhanced by high temperature was mentioned as one of main contributor to the increase at northern high-latitudes (Dlugokencky et al. 2009; Bloom et al. 2010). This study shows the year-to-year variation of CH₄ emissions from West Siberian wetlands estimated from atmospheric CH₄ observed by a tower network (JR-STATION: Japan-Russia Siberian Tall Tower Inland Observation Network) and aircraft over Siberia, using inverse model of atmospheric CH₄ transport based on a fixed-lag Kalman smoother. We also use flask sampling and continuous measurement data of atmospheric CH₄ archived at WDCGG (World Data Centre for Greenhouse Gases) in flux estimates. Interannually varying CH₄ emissions are used to calculate CH₄ transport with NIES transport model including chemical sink rates developed in TransCom-CH₄ project (Patra et al. 2011): wetland and rice paddy emissions and soil sinks simulated with a process-based biogeochemical model (VISIT), biomass burning emissions of GFED v3.1, anthropogenic emissions of EDGAR v4.2, and interannually repeating termite emissions of GISS.

Annual mean of Siberian wetland CH₄ flux was estimated to be 6.9 +/- 1.1 Tg/yr in 2006-2010 and high wetland flux was concentrated between (57.5N, 65.0E) and (67.5N, 90.0E) in West Siberia (called WL area), occupying 57% (3.9 +/- 0.2 Tg/yr) of the estimated Siberian wetland flux. The annual mean of WL area was very close to that for VISIT emission (3.8 Tg/yr), but a larger year-to-year variation was estimated in wetland flux of WL area (0.8 Tg/yr). A higher wetland flux of WL area was estimated in 2007 and 2008, but lower in 2006 and 2010. The enhanced wetland fluxes in 2007 and 2008 coincided with higher surface air temperature of NCEP/NCAR and greater precipitation of GPCP than those means in 1991-2010 over WL area and explained high CH₄ concentration observed in May-Sep 2007 and 2008 at Demyanskoe and Karasevoe near extensive wetlands in WL area. The year-to-year variation of observed CH₄ concentration was well reproduced with inverse model-estimated fluxes, showing high positive correlation between observed and predicted CH₄ concentrations ($r = 0.85$ and 0.98 at Demyanskoe and Karasevoe, respectively). In WL area, we found a high positive correlation of annual mean of inverse model-estimated wetland flux with annual mean surface air temperature ($r = 0.89$) and liquid water equivalent thickness of GRACE ($r = 0.92$), but relatively low correlation ($r = 0.37$) for precipitation.

Acknowledgment: We thank to WDCGG data contributors for atmospheric CH₄ observations and TransCom-CH₄ project for developing a simulation setup.