Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



ACG35-19

会場:301A

森林性ネズミ2種の生育地分割パターンと寒冷地特異的繁殖行動 Habitat partitioning between sympatric Japanese wood mice

鈴木 亮^{1*}, 坂本信介², 鈴木智之³ Ryo Suzuki^{1*}, Shinsuke Sakamoto², Satoshi Suzuki³

¹ 筑波大学菅平高原実験センター,² 宮崎大学フロンティア科学実験総合センター,³ 東京大学大学院農学生命科学研究科 ¹Sugadaira Montane Research Center, University of Tsukuba, ²Divisions of Bio-Resources, Department of Biotechnology, Frontier Science Research Center, Universit, ³The University of Tokyo Forests, Graduate School of Agricultural and Life Sciences

Habitat partitioning can maintain the coexistence of species with very similar ecological traits. We studied habitat partitioning between terrestrial and semi-arboreal congeneric rodents (Apodemus speciosus and A. argenteus) that often coexist despite asymmetry in their competitive abilities. To understand seasonal and habitat variation in their partitioning, we evaluated seasonal variation in food resources, habitat use, and habitat similarity between the species in a site comprising a mixture of grassland, pine forest, and mixed forest. Food resources were available on the ground in all vegetation types in spring and autumn, but were severe in summer. Apodemus speciosus was observed in all types of vegetation on the ground. In contrast, A. argenteus was observed on the ground and on trees in pine forest, especially areas where the understory is covered by dwarf bamboo.

Habitat similarity tests revealed that habitat partitioning between two Apodemus species may vary seasonally, and the relationship depends on habitat structure. This study suggests that the mechanism of habitat partitioning between terrestrial and semi-arboreal rodents in temperate forest is more complex than previously recognized.

This study additionally found that breeding seasons of the two species in this cold region were observed on summer from June to October, whereas breeding seasons of them are observed on spring and autumn in many temperate regions in Japan.

キーワード:共存メカニズム,水平生育地分化,ニッチ分割,垂直生育地分化

Keywords: coexistence mechanisms, horizontal partitioning, niche division, vertical partitioning