Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



AGE04-08

会場:102B

時間:5月20日16:45-17:00

下方浸透促進による効果的な放射性降下物の地表からの削減 Radioactive fallout removal from the surface soils by enhancing vertical transport.

宮本珠未 1 , 森 也寸志 1* , 稲生栄子 2 , 末継 淳 1 , 登尾浩助 3 Tamami Miyamoto 1 , Yasushi Mori 1* , Eiko Inao 2 , Atsushi Suetsugu 1 , Kosuke Noborio 3

Fukushima nuclear power plant damaged by the East Japan Great Earthquake caused radioactive fallout around the Tohoku region. Because radioactive fallout was positively charged, it was reported to being adsorbed to soil surface. Surface soil scraper and deep plowing would be, therefore, effective for the removal of radioactive materials. However, these techniques were available for flat and wide land like school yard or farm land. Field survey revealed that correlation between EC and radioactive dosage were significant, which meant radioactive fallout did not immediately adsorb to soil surface but stayed as exchangeable ion for a while and was transported with surface water.

Thus we applied artificial macropores to effectively remove radioactive fallout from the surface soil. Artificial macropore filled with bamboo fiber was made in soils (Field: d=1cm length=50cm, Lab: d=0.6cm, length=20cm). Zeolite was placed at the bottom of the macropores (Field: 5cm, Lab: 3cm) to absorb transported Cesium. Four treatment was prepared such as macropore, macropore with ammonium sulfate, no macropore and no macropore with ammonium sulfate. 400mm artificial rainfall was applied in one month.

Results showed artificial macropore effectively transported radioactive Cesium to deeper profile while ammonium sulfate enhanced the amount of Cesium delivered to the deeper profile. No radioactive Cesium was observed from the drainage water.

Field experiment showed significant reduction in macropore plots than no-macropore plots, however, it is not significant for ammonium sulfate plots.

キーワード: 放射性降下物, 下方浸透, マクロポア Keywords: radioactive fallout, infiltration, macropore

¹ 岡山大学, ² 宮城県農業・園芸総合研究所, ³ 明治大学

¹Okayama University, ²Miyagi Prefectural Institute of Agriculture and Horticulture, ³Meiji University