Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



AHW02-P06

会場:コンベンションホール

時間:5月21日18:15-19:30

## ベトナムにおける近年の秋季降水量の数十年変動 Recent Interdecadal Variations of Autumnal Precipitation in Vietnam

井上 知栄<sup>1\*</sup>, 松本 淳<sup>2</sup>, 遠藤 伸彦<sup>1</sup> Tomoshige Inoue<sup>1\*</sup>, Jun Matsumoto<sup>2</sup>, ENDO, Nobuhiko<sup>1</sup>

## 1海洋研究開発機構,2首都大学東京

<sup>1</sup>Japan Agency for Marine-Earth Science and Technology, <sup>2</sup>Tokyo Metropolitan University

In some parts of Southeast Asia, (e.g., central Vietnam), a large amount of rainfall occurs in boreal autumn to early winter (September-December: SOND). In the present study, we investigate interdecadal variations of rainfall in SOND in Vietnam and its vicinity for the period 1961-2010, based on rain gauge observational data obtained from the Southeast Asian countries. As a result, it is very obvious that rainfall have increased [decreased] to the south [north] of 17N along the coastal area of Vietnam. There are many stations with statistically significant decrease over the Red River Delta region (north of 20N; denoted as region A hereafter). In this region, SOND rainfall has decreased since late 1980s. In northern part of central Vietnam (17-20N; denoted as region B), the rainfall decrease has been observed since late 1990s. In southern part of central Vietnam (12-17N; denoted as region C), on the other hand, SOND rainfall has largely increased since late 1990s. From comparison of seasonal marches of rainfall over the 3 regions between 25-year averages of 1961-85 and 15-year averages of 1996-2010, we find that the amount of rainfall in region A has clearly decreased in August-October, indicating the recent earlier withdrawal of summer rainy season. In region B, rainfall has decreased during a whole rainy season in boreal autumn. In region C, on the other hand, the rainfall increase is very obvious and long-lasting during a whole period of August to December. Atmospheric circulation changes based on some gridded datasets suggest that recent stronger lower-tropospheric cyclonic circulation over the southern part of the South Family South China Sea might be responsible for the recent increasing trend of rainfall over region C, though the significance of the atmospheric changes varies between the gridded datasets.

キーワード: 気候変動, ベトナム, 降水量

Keywords: climate variations, Vietnam, precipitation