Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



AHW29-P03

会場:コンベンションホール

## 弱風時の水面上のバルク輸送係数 - 広域大気循環の影響? Bulk coefficients over a water surface under calm wind condition: an influence of largescale atmospheric circulation?

魏 忠旺<sup>1\*</sup>, 杉田 倫明<sup>1</sup> Zhongwang Wei<sup>1\*</sup>, Michiaki Sugita<sup>1</sup>

<sup>1</sup> 筑波大・生命環境・地球

<sup>1</sup>Geoscience, Life & Env Sci., Univ.Tsukuba

Over the Lake Kasumigaura, 10 Hz fluxes data have been obtained since June of 2007, and characteristics of the bulk transfer coefficients of momentum Cdn and of water vapor and heat have been studied with emphasis on the possible influence of the large-scale convection to surface fluxes. Increase of Cdn when mean wind speed u becomes smaller in the range of u < 4 m/s was found. By applying the cospectral and ogive analyses, a possible cause of this increase was the influence of convective circulation, as manifested by the mismatch of characteristic time scale of the flux (2nd moment) and mean wind speed (1st moment). Possible solutions to this the mismatch of time scale under weak wind speed was studied which include adaptation of the effective wind speed Ue instead of the traditional vector-averaged mean wind speed. However, complete solution is not likely available since this problem arises from ill-defined nature of Cdn under light wind condition.

Keywords: Lake Kasumigaura, Bulk coefficients, ogive, large-scale convection, fluxes