Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

会場:301A

BAO01-15

時間:5月21日16:45-17:00

Amazonis and Elysium basins and Their Link, Marte Vallis (AME), Tharsis/Elysium Corridor, Mars Amazonis and Elysium basins and Their Link, Marte Vallis (AME), Tharsis/Elysium Corridor, Mars

Dohm James^{1*}, 丸山 茂徳¹, 宮本 英昭² James Dohm^{1*}, Shigenori Maruyama¹, Hideaki Miyamoto²

¹ 東京工業大学 地球生命研究所,² 東京大学総合研究博物館

¹Earth-Life Science Institute, Tokyo Institute of Technology, ²The University Museum, The University of Tokyo

The geologic provinces of Mars, as identified through a synthesis of geologic, paleohydro-logic, topographic, geophysical, spectral, and elemental information [1], are windows into its evolution, with the youthful province, the Tharsis/Elysium corridor [2-4], recording hydrologic, tectonic, and volcanic activity, including fissure-fed eruptions and shield-volcano field development, within at least tens of millions of years.

Geologically recent activity in the Tharsis/Elysium corridor region is particularly high-lighted in the Amazonis and Elysium basins and their link, Marte Vallis (hereafter referred to as AME). A youthful shield-volcano field with at least seven gentle sloping shield volcanoes and fissure-fed vent structures have been identified, mapped, and characterized in the western part of the Elysium. In addition, the lava flows on and near the margins of the shield volcanoes display crenulated lava flow margins, possibly marking lava-water-ice interactions. Both the shield volcanoes and pristine lavas located within AME, which are marked by a paucity of relatively small (~few-km-diameter) superposed impact craters and modified by faults and fractures and valleys, point to geological and hydrological activity on Mars in recent geologic time, making AME a significant target for future reconnaissance, including testing the hypotheses of whether Mars is geologically, hydrologically, and biologically active.

Specifically, the following questions might be addressed through international reconnaissance missions to AME, which would include instrument suites with optimal geologic, geochemical, geophysical (including seismic), environmental, and biological capabilities, including whether Mars: (1) is geologically and hydrologically active, (2) contains salty groundwater and magma at relatively shallow depths, (3) has sustained elevated heat flow, (4) records seismic activity, and (5) comprises fossilized and/or extant life.

References

- [1] Dohm, J.M. et al., (2013?in press), Nova Science Publishers, Inc.
- [2] Dohm, J.M., et al., 2008, Planetary and Space Science 56, 985?1013, doi:10.1016/j.pss.2008.01.001
- [3] Dohm, J.M., (2004) Planetary and Space Science 52, 189?198, doi:10.1016/j.pss.2003.08.013.
- [4] Miyamoto, H., (2004) Journ. of Geophy. Res. 109, E06008, doi:10.1029/2003JE002234.