

## 系外惑星のバイオマーカーのための光合成光吸収についての理論的研究 Theoretical investigation on the absorption spectrum of photosystem for the biomarker of extrasolar planets

田口 真彦<sup>1</sup>, 小松勇<sup>1</sup>, 佐藤皓允<sup>1</sup>, 庄司 光男<sup>1\*</sup>, 栢沼愛<sup>2</sup>, 神谷克政<sup>1</sup>, 白石賢二<sup>1</sup>, 矢花一浩<sup>1</sup>, 梅村雅之<sup>1</sup>  
Masahiko Taguchi<sup>1</sup>, Yu Komatsu<sup>1</sup>, Akimasa Sato<sup>1</sup>, Mitsuo Shoji<sup>1\*</sup>, Megumi Kayanuma<sup>2</sup>, Katsumasa Kamiya<sup>1</sup>, Kenji Shiraishi<sup>1</sup>,  
Kazuhiro Yabana<sup>1</sup>, Masayuki Umemura<sup>1</sup>

<sup>1</sup> 筑波大学数理物質科学研究科, <sup>2</sup> 筑波大学システム情報研究科

<sup>1</sup>Graduate School of Pure and Applied Science, University of Tsukuba, <sup>2</sup>Department of Computer Science, University of Tsukuba

Over 800 extrasolar planets have been discovered, and more than 20,000 candidates have been detected. Planets in habitable zone have been observed, and the discovery of Earth-like planets is expected. Great attentions have been paid to the detection of life in extrasolar planets. For the detection, various indices have been proposed as biomarkers. One of the indicators is red edge[1], which is a characteristic steep gradient observed in the near-infrared region of around 750 nm in plant's reflection spectra [2]. In fact, red edge can be observed in the reflection spectrum of the Earth via the Moon (earthshine [3]). Since red edge is affected by many factors, its precise predictions is not simple. However, a leaf chlorophyll absorption is thought to be the major factor of red edge [2].

On Earth, photosynthetic organisms have evolved through the collection of sunlight. On the other hand for the extrasolar planets, whose surrounding space environment has different spectrum from their primary star, photosystems should be different in many parts, such as pigment types and arrangements even though the environment is similar to the Earth.

Before predictions of biomarkers of extrasolar planets, we examine the basic characters of chlorophylls in photosystems. It is also important for the study of the diverse photosystems on Earth. Chlorophylls are concentrated in a chloroplast, and form pigments-protein complexes in the photochemical systems.

The purpose of this research is to characterize the absorption spectrum of chlorophylls in a photosystem. First of all, we calculated the absorption spectrum of the pigment in methanol using DFT based polarization continuum model (PCM) method, and confirmed the validity of our calculation method. Then, quantum mechanics/molecular mechanics (QM/MM) calculations were performed for the absorption spectrums of the photosystem. Each chlorophyll was included in the QM region. We found that the absorption wavelengths are shifted about +10 nm by the effects of the protein environment. Similar influence was observed by the effect of amino acid coordination to the central Mg ion in the chlorophyll. These calculated results indicate a fine modulation character of the adsorption wavelength for the photosystem. This character is important for photosystems in extrasolar planets as well as in extreme conditions on Earth.

[1] N. Y. Kiang et al., *Astrobiology*, 7 (1), 252, 2007

[2] S. Seager et al., *Astrobiology*, 5 (3), 372, 2005

[3] L. Arnold et al., *Astronomy&Astrophysics*, 392, 231, 2002

キーワード: バイオマーカー, 系外惑星, 光合成, 量子古典混合計算, 光吸収

Keywords: biomarker, extrasolar planet, photosystem, QM/MM, absorption spectrum