

飼育実験手法を基にした溶存酸素濃度の浮遊性有孔虫への影響評価
Laboratory culture experiments: effect of dissolved oxygen concentration on planktonic foraminifera

黒柳 あずみ^{1*}, da Rocha Regine E.², Bijma Jelle², Spero Howard³, 川幡 穂高¹
Azumi Kuroyanagi^{1*}, Regine E. da Rocha², Jelle Bijma², Howard Spero³, hodaka kawahata¹

¹ 東京大学 大気海洋研究所, ² アルフレッドウェゲナー極域海洋研究所, ³ カリフォルニア大学 デービス校

¹ Atmosphere and Ocean Research Institute, The University of Tokyo, ² Alfred-Wegener-Institut für Polar- und Meeresforschung,

³ University of California Davis

Planktonic foraminifera shows significant species turnover ratio related Cretaceous oceanic anoxic event (OAEs), however, the direct effect of dissolved oxygen (DO) concentration on planktonic foraminifera remain obscure. Although culture experiments can investigate quantitatively the relationship between foraminiferal ecology and environmental parameters, DO-controlling experiment has never been conducted because of the difficulty of observation and/or control of dissolved oxygen under the modern ocean condition. In the present study, we cultured two planktonic foraminifera species (*Orbulina universa* and *Globigerina bulloides*) under six different DO conditions. Both species have extremely high tolerance to low DO than we expected before, and it suggests that at least "dysoxic" condition (more than $0.7 \text{ mg O}_2 \text{ l}^{-1}$) could not be a direct cause of the extinction of planktonic foraminifera at OAEs. Their high tolerance to extremely low DO would be caused by the evolutionary descendant of benthic foraminifera. Final shell weight increased with DO despite almost the same culture duration among treatments, thus it suggests foraminiferal fossil shell weight could reflect the DO conditions.

キーワード: 浮遊性有孔虫, 飼育実験, 溶存酸素濃度

Keywords: planktonic foraminifer, culture experiment, dissolved oxygen concentration