

南日本で採集した化石シャコガイ殻のSr/Ca比を用いた4.6kaの日射量の復元
Daily light cycle reconstructed by Sr/Ca in a fossil giant clam, *Tridacna gigas*, living in
4.6 ka, southern Japan

堀 真子^{1*}, 白井厚太郎¹, 高畑 直人¹, 佐野 有司¹, 小林夏子², 岡本 和明², 渡邊 剛³

Masako Hori^{1*}, Kotaro Shirai¹, Naoto Takahata¹, Yuji Sano¹, Natsuko Kobayashi², Kazuaki Okamoto², Tsuyoshi Watanabe³

¹ 東京大学大気海洋研究所, ² 埼玉大学教育学部, ³ 北海道大学理学部

¹Atmosphere and Ocean Research Institute, the University of Tokyo, ²Faculty of Education, Saitama University, ³Faculty of Science, Hokkaido University

Giant clams are long-lived bivalvia living in a shallow tropical ocean. Their aragonite shells are potential paleo-environmental archives, i.e., oxygen stable isotopic values record paleo-seawater temperatures. Generally, trace elements in calcium carbonates are also considered as a paleo-thermometer. However, the case of Tridacninae may not follow the traditional temperature dependent fractionation, but potentially reflects insolation. This is explained by "trans-calcification", which is one of photosynthetically induced calcification mechanisms. Enzymatic reaction pumping Ca^{2+} to host liquid of calcification results in relatively depleted Sr/Ca ratio in day time when photosynthesis activates. Following this hypothesis, we have reconstructed the past daily light cycle.

A fossil *T. gigas*, Stg04-b was collected at Ishigaki-jima Island in Okinawa Prefecture, Japan (N24 20'0.4" E124 09'22"). Two horizons of the specimen were dated by carbon-14 method at the accelerator mass spectrometry (AMS) center of Yamagata University. The micro analyses were performed for the outer layer of 1.4 cm thickness using an electron probe micro-analyzer (EPMA), JXA8900, and a micrometer-scale secondary ion mass spectrometer (NanoSIMS), CAMECA NS50 at the Atmosphere and Ocean Research Institute (AORI), the University of Tokyo. As a result, we found clear Sr lamination parallel to the daily increment of about 20 μm thickness. The hourly insolation (P) was calculated from the difference of Sr/Ca ratio ($\Delta\text{Sr}/\text{Ca}$) following the equation: $\Delta\text{Sr}/\text{Ca} = -a \times P$, where the parameter, a , is defined by analysis of modern *T. maxima* [1]. Using the technique, we distinguished the sunny and cloudy seasons in 4.6 ka.

Reference

[1] Sano, Y., Kobayashi, S., Shirai, K., Takahata, N., Matsumoto, K., Watanabe, T., Sowa, K., Iwai, K. 2012, Past daily light cycle recorded in the strontium/calcium ratios of giant clam shells. Nature Commun. 3, DOI: 10.1038/ncomms1763

Keywords: giant clam, Sr/Ca, NanoSIMS