

## 酸素量で制御された後生動物の初期進化と絶滅 Early metazoan evolution and extinction controlled by oxygen levels

海保 邦夫<sup>1\*</sup>, 大庭雅寛<sup>1</sup>, 菊池みのり<sup>1</sup>, 千馬 直登<sup>1</sup>, 静谷 あてな<sup>1</sup>, 山田 憲司<sup>1</sup>, チェン・ツォン チャン<sup>2</sup>, トン ジンナン<sup>2</sup>, ポール ゴーザン<sup>3</sup>, 高橋 聡<sup>1</sup>  
Kunio Kaiho<sup>1\*</sup>, Masahiro Oba<sup>1</sup>, Minoru Kikuchi<sup>1</sup>, Naoto Senba<sup>1</sup>, Atena Shizuya<sup>1</sup>, Kenji Yamada<sup>1</sup>, Zhong-Qiang Chen<sup>2</sup>, Jinnan Tong<sup>2</sup>, Paul Gorjan<sup>3</sup>, Satoshi Takahashi<sup>1</sup>

<sup>1</sup> 東北大学大学院理学研究科地学専攻, <sup>2</sup> 中国地質大学, <sup>3</sup> ワシントン大学

<sup>1</sup>Institute of Geology and Paleontology, Tohoku University, <sup>2</sup>China University of Geosciences, <sup>3</sup>Washington University

Organic-molecular dissolved-oxygen index above and below storm wave base, from Cryogenian to Cambrian marine sedimentary rocks in Australia and China records three rises in dissolved oxygen levels. The first rise in dissolved oxygen levels coincides with molecular diversification of animals in the early Ediacaran, the second rise with appearance of large Ediacaran animal fossils, its drop with extinction of Ediacaran metazoa, and the third rise is coeval the explosion of Cambrian metazoa. Our evidence for widespread dissolved-oxygen changes synchronizing with macroevolution and extinction suggests that the global dissolved oxygen level in the sea had controlled the evolution of metazoans during the Neoproterozoic-Phanerozoic transition. The first rise in dissolved oxygen levels in the early Ediacaran is newly found and consistent with molecular diversification of animals in the early Ediacaran.

キーワード: 酸素, 進化, 絶滅

Keywords: oxygen, evolution, extinction