

## 砕屑物のNd-Pb同位体比を用いたアラスカ氷河発達史の解析 History of Alaskan mountain glaciers analyzed by detrital Nd and Pb isotopes

堀川 恵司<sup>1\*</sup>, 小野寺丈尚太郎<sup>2</sup>, 関宰<sup>3</sup>, 坂本竜彦<sup>2</sup>

Keiji Horikawa<sup>1\*</sup>, Jyonaotaro Onodera<sup>2</sup>, Osamu Seki<sup>3</sup>, Tatsuhiko Sakamoto<sup>2</sup>

<sup>1</sup> 富山大学大学院理工学研究部, <sup>2</sup> 海洋研究開発機構, <sup>3</sup> 北海道大学低温科学研究所

<sup>1</sup> Graduate School of Science and Engineer, University of Toyama, <sup>2</sup> JAMSTEC, <sup>3</sup> Institute of Low Temperature Science, Hokkaido Univ

Changes in atmospheric CO<sub>2</sub>, ocean circulation, the extent of the cryosphere, or a combination of these boundary conditions are believed to have caused the Late Pliocene climate Transition (LPT, ~3.6 to 2.6 million years ago), characterized by development of more glaciated conditions. Yet, cryospheric evolution in northern high latitudes is not constrained well enough to properly understand the effect of high latitude cryosphere dynamics on LPT. Here, we present 4.1-million-year neodymium (Nd) and lead (Pb) isotope records of detrital sediments?tracers of sediment provenance?from the Bering Sea (Hole U1341B). We show that Bering Sea detrital sediments represented a mixture of sediments from the Aleutian and Alaskan sectors. During glacial and deglacial periods the contribution of Alaskan-origin clastics exceeds 50% due to melting of Alaskan glacier. We also found a stepwise increase (>~10%) in Alaskan-origin clastic supply beginning at ~4.1 Ma, implying that the northern high latitude cryosphere had started to grow 1 Ma before the major onset of Northern Hemisphere Glaciation. The early growth of Alaskan glaciers might be a result of combined effects of decreased atmospheric pCO<sub>2</sub> and enhanced moisture supply due to stagnation of Pacific meridional circulation induced by the Panamanian seaway closure during the early Pliocene.

キーワード: ネオジウム同位体, 鉛同位体, ベーリング海, 雪氷圏, 鮮新世温暖期