

海洋堆積物中のアミノ酸の起源：アミノ酸と光合成色素の化合物レベル窒素同位体組成からの推定

The sources of amino acids in marine sediments estimated from nitrogen isotopic composition of amino acids and chlorin

山口 保彦^{1*}, 菅 寿美², 高野 淑識², 力石 嘉人², 小川 奈々子², 横山 祐典³, 大河内 直彦²

Yasuhiko T. Yamaguchi^{1*}, Hisami Suga², Yoshinori Takano², Yoshito Chikaraishi², Nanako O. Ogawa², Yusuke Yokoyama³, Naohiko Ohkouchi²

¹ 東京大学大学院理学系研究科, ² 海洋研究開発機構 海洋・極限環境生物圏領域, ³ 東京大学大気海洋研究所

¹Department of Earth and Planetary Science, The University of Tokyo, ²Institute of Biogeosciences, JAMSTEC, ³Atmosphere and Ocean Research Institute, The University of Tokyo

Amino acids are the building blocks of proteins and peptides and key compounds in microbial metabolisms. Amino acids represent one of major fractions of sedimentary organic matter and are important in undergoing organic matter mineralization in marine sediments. However, our understanding is still limited about the biogeochemical dynamics of amino acids in marine sediments, in part because currently available methods are not sufficient to constrain sources and transformation processes of amino acids in sediments.

Here, as a new method to estimate sources of amino acids in marine sediments, this study reports down-core profiles of compound-specific nitrogen isotopic composition ($\delta^{15}\text{N}$) of total hydrolysable amino acids (THAA) in marine sediments of the Japan Sea (a surface sediment and a 7-m-long piston core; ca. 46,500 years). The $\delta^{15}\text{N}$ profiles of THAA were compared with a down-core $\delta^{15}\text{N}$ profile of chlorin pigment (pyropheophytin a), which reflects the $\delta^{15}\text{N}$ values of organic matter produced by photosynthetic organisms in the past ocean.

Significant correlations were observed between $\delta^{15}\text{N}$ of amino acids and chlorin pigment in the piston core samples ($r^2 = 0.87$ for phenylalanine, 0.78 for glutamic acid, 0.77 for alanine, and 0.62 for glycine; $n = 13$). This result suggests that the major source of THAA is organic matter produced by the organisms in the past ocean (i.e., necromass) and that contribution of in situ sedimentary microbial production to THAA is less than 15% below 1 m depth in the core. The offset values of $\delta^{15}\text{N}$ between amino acids and chlorin pigment in the sediments of 1-7 mbsf (e.g., $\delta^{15}\text{N}$ -phenylalanine - $\delta^{15}\text{N}$ -chlorin = +7.3 permil in average) suggest that the source organisms of THAA contain not only photosynthetic algae and animals but also heterotrophic or chemoautotrophic microbes in the past ocean (water column and surface sediments).

キーワード: 有機物, 窒素同位体, アミノ酸, 光合成色素, 海洋堆積物, 古海洋

Keywords: Organic matter, Nitrogen isotope, Amino acids, Photosynthetic pigments, Marine sediments, Paleoceanography