

Magnetic anisotropies for tsunami deposits: Application to the 3.11 Magnetic anisotropies for tsunami deposits: Application to the 3.11

昆周作^{1*}, 中村 教博¹, 後藤 和久², 菅原 大助², Catherine Chague-Goff³, 飯嶋 耕崇¹, Jamaes Goff³
Shusaku Kon^{1*}, Norihiro Nakamura¹, Kazuhisa Goto², Daisuke Sugawara², Catherine Chague-Goff³, Yasutaka Iijima¹, Jamaes Goff³

¹ 東北大学理学研究科地学専攻, ² 東北大学災害科学国際研究所, ³ Environmental and Earth Sciences, University of New South Wales

¹ Graduate School of Science, Tohoku University, ² International Research Institute of Disaster Science (IRIDeS), Tohoku University, ³ Environmental and Earth Sciences, University of New South Wales

Tsunami deposits consist of well-sorted fine sand intercalating with non-marine black organic mud. It is difficult to reveal a transport direction of the deposit if the deposit showed no sedimentary fabrics, such as ripples. The proxy of anisotropy of magnetic susceptibility (AMS) appears to be a promising tool for the study of flow fabrics in recent-tsunami deposits such as Sumatra tsunami (Wassmer et al. 2010). The AMS fabric might allow us to reconstruct transport directions of unconsolidated tsunami sediments during emplacement because AMS provides a cryptic alignment of ferromagnetic and paramagnetic minerals. Such cryptic minerals, such as magnetite or phyllosilicate minerals, would behave as a different emplacement mode in a different hydrodynamic condition. In the AMS fabrics of volcanic rocks, there are large discrepancies between the magnetic lineation and the framework-forming silicate linear fabric. This suggests that the uncorroborated use of bulk AMS to detect flow fabric in tsunami deposits has risks. In this article, we show that the anisotropy of anhysteretic remanent magnetization (AARM) may resolve the difficulties. The combination of inundation eye-witness, SEM, AMS, and AARM confirms the flow pattern of recent- and paleo-tsunami deposits from the geoslicer sampleing at Rikuzen-Takata city, Japan during 2011, 11th March Tohoku tsunami. We determined if the sandy deposits are of tsunami from these magnetic anisotropies.

キーワード: 津波堆積物, 古地磁気, 帯磁率異方性, 非履歴性残留磁化

Keywords: tsunami deposits, paleomagnetism, anisotropy of magnetic susceptibility, anisotropy of anhysteretic remanent magnetization